



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 02:18 PM UTC

PDB ID : 4W9U / pdb_00004w9u
Title : Crystal Structure of an Acyl-CoA dehydrogenase from Brucella melitensis
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2014-08-27
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	
1	B	395	
1	C	395	
1	D	395	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 11315 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-CoA dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	362	Total	C	N	O	S	0	0	0
			2630	1660	467	487	16			
1	B	376	Total	C	N	O	S	0	1	0
			2809	1776	495	520	18			
1	C	365	Total	C	N	O	S	0	1	0
			2678	1689	477	496	16			
1	D	375	Total	C	N	O	S	0	1	0
			2844	1796	504	527	17			

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	66	Total O 66 66	0	0
3	B	103	Total O 103 103	0	0
3	C	77	Total O 77 77	0	0
3	D	84	Total O 84 84	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Acyl-CoA dehydrogenase




• Molecule 1: Acyl-CoA dehydrogenase




• Molecule 1: Acyl-CoA dehydrogenase



• Molecule 1: Acyl-CoA dehydrogenase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.02Å 106.66Å 99.39Å 90.00° 108.12° 90.00°	Depositor
Resolution (Å)	47.23 – 2.40 47.23 – 2.40	Depositor EDS
% Data completeness (in resolution range)	88.7 (47.23-2.40) 88.7 (47.23-2.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.41 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1769)	Depositor
R, R_{free}	0.180 , 0.239 0.183 , 0.241	Depositor DCC
R_{free} test set	2806 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtriage
Anisotropy	0.226	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11315	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/2678	0.89	5/3627 (0.1%)
1	B	0.59	0/2863	0.90	5/3866 (0.1%)
1	C	0.54	0/2730	0.91	5/3693 (0.1%)
1	D	0.57	0/2898	0.88	8/3904 (0.2%)
All	All	0.56	0/11169	0.89	23/15090 (0.2%)

There are no bond length outliers.

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	188	HIS	N-CA-C	7.37	123.18	113.30
1	C	10	ASP	CA-C-N	6.96	126.78	119.05
1	C	10	ASP	C-N-CA	6.96	126.78	119.05
1	C	188	HIS	N-CA-C	-6.94	99.94	110.14
1	B	173	SER	CA-C-N	-6.86	112.74	119.87
1	B	173	SER	C-N-CA	-6.86	112.74	119.87
1	D	109	TYR	CA-C-N	-6.47	112.33	119.19
1	D	109	TYR	C-N-CA	-6.47	112.33	119.19
1	A	10	ASP	CA-C-N	6.34	126.09	119.05
1	A	10	ASP	C-N-CA	6.34	126.09	119.05
1	D	186	ALA	N-CA-C	-5.97	101.57	110.23
1	C	173	SER	CA-C-N	-5.87	113.76	119.87
1	C	173	SER	C-N-CA	-5.87	113.76	119.87
1	A	173	SER	CA-C-N	-5.78	113.86	119.87
1	A	173	SER	C-N-CA	-5.78	113.86	119.87
1	D	10	ASP	CA-C-N	5.77	125.89	119.32
1	D	10	ASP	C-N-CA	5.77	125.89	119.32
1	B	51	ASP	CA-C-N	5.65	125.32	119.56
1	B	51	ASP	C-N-CA	5.65	125.32	119.56
1	B	371	THR	N-CA-C	5.62	117.21	111.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	173	SER	CA-C-N	-5.38	114.28	119.87
1	D	173	SER	C-N-CA	-5.38	114.28	119.87
1	D	177	ASP	N-CA-C	-5.11	107.17	113.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2630	0	2448	15	0
1	B	2809	0	2717	25	0
1	C	2678	0	2509	15	0
1	D	2844	0	2789	28	0
2	A	8	0	12	0	0
2	C	4	0	4	0	0
2	D	12	0	18	0	0
3	A	66	0	0	2	0
3	B	103	0	0	1	0
3	C	77	0	0	0	0
3	D	84	0	0	1	0
All	All	11315	0	10497	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:349:GLY:HA2	1:C:371:THR:HG23	1.76	0.66
1:B:156:LYS:HB2	1:B:187:ALA:HB2	1.80	0.63
1:D:141:PRO:HB3	1:D:168:MET:HE2	1.81	0.61
1:D:160:GLY:HA2	1:D:232:GLU:HG3	1.83	0.60
1:C:15:GLU:OE1	1:C:26:ARG:NH1	2.32	0.60
1:C:188:HIS:O	1:C:190:ASN:N	2.35	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:64:LEU:HD13	1:C:103:GLN:HG2	1.84	0.58
1:A:66:VAL:HG13	1:A:77:ALA:HB3	1.85	0.58
1:A:68:LEU:HD12	1:A:108:MET:HE2	1.86	0.56
1:D:25:ILE:HG21	1:D:81:ALA:HB1	1.87	0.56
1:A:188:HIS:N	1:A:189:ASP:HA	2.20	0.56
1:B:209:ILE:HG23	1:B:221:GLY:HA2	1.88	0.55
1:B:371:THR:HG23	1:C:349:GLY:HA2	1.88	0.55
1:B:25:ILE:HG21	1:B:81:ALA:HB1	1.88	0.55
1:C:70:GLU:HG3	1:C:76:ASN:HB2	1.89	0.54
1:B:68:LEU:HD11	1:B:108:MET:HB3	1.88	0.54
1:A:349:GLY:HA2	1:D:371:THR:HG23	1.90	0.54
1:A:290:LEU:HD23	1:C:290:LEU:HD23	1.90	0.53
1:B:68:LEU:HD12	1:B:108:MET:HE2	1.90	0.53
1:B:65:GLY:HA3	1:B:108:MET:HE2	1.90	0.52
1:B:65:GLY:HA3	1:B:108:MET:CE	2.40	0.52
1:D:232:GLU:O	1:D:235:ILE:HG22	2.09	0.52
1:C:114:TYR:HB2	1:C:236:LEU:HD13	1.91	0.52
1:A:343:GLN:HG2	3:A:533:HOH:O	2.09	0.51
1:D:153:ARG:NH2	1:D:190:ASN:OD1	2.44	0.51
1:D:168:MET:HG3	1:D:169:TRP:N	2.26	0.51
1:A:33:ALA:HB1	1:A:92:ILE:HD13	1.93	0.50
1:D:266:ASP:OD2	1:D:270[A]:ARG:NH1	2.44	0.50
1:B:168:MET:SD	1:B:222:GLU:HG2	2.52	0.49
1:C:68:LEU:HD12	1:C:108:MET:HE2	1.94	0.49
1:D:20:GLU:OE1	1:D:23:ARG:NH2	2.41	0.49
1:B:168:MET:HE3	1:B:169:TRP:CE2	2.48	0.49
1:D:65:GLY:HA3	1:D:108:MET:CE	2.43	0.48
1:D:182:TRP:CE3	1:D:245:PRO:HD3	2.49	0.48
1:B:108:MET:HE3	1:B:128:VAL:HG22	1.96	0.47
1:D:108:MET:HE3	1:D:128:VAL:HG22	1.96	0.47
1:A:181:VAL:HG11	1:A:225:MET:HE1	1.97	0.47
1:C:107:VAL:HG11	1:C:134:GLY:HA3	1.97	0.47
1:B:291:TYR:CZ	1:B:347:MET:HE2	2.51	0.46
1:B:172:ASN:HA	1:B:219:ILE:HG23	1.97	0.46
1:A:316:PHE:CE2	1:A:321:MET:HG3	2.51	0.46
1:A:291:TYR:CZ	1:A:347:MET:HE2	2.51	0.46
1:D:65:GLY:HA3	1:D:108:MET:HE1	1.98	0.46
1:D:343:GLN:HG2	3:D:530:HOH:O	2.16	0.45
1:A:11:PRO:HD2	3:A:565:HOH:O	2.16	0.45
1:B:106:LEU:HB3	1:B:136:PHE:HB2	1.98	0.45
1:B:124:LEU:HD23	1:B:124:LEU:HA	1.74	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:141:PRO:HG3	1:B:168:MET:HG3	1.99	0.45
1:C:291:TYR:CZ	1:C:347:MET:HE2	2.52	0.45
1:C:68:LEU:HD11	1:C:108:MET:HB3	1.99	0.44
1:D:106:LEU:HB3	1:D:136:PHE:HB2	1.98	0.44
1:A:186:ALA:O	1:A:187:ALA:HB3	2.18	0.44
1:D:291:TYR:CZ	1:D:347:MET:HE2	2.51	0.44
1:A:338:LEU:HD11	1:A:367:GLU:HG3	1.99	0.44
1:C:19:THR:O	1:C:23:ARG:HG3	2.18	0.44
1:D:186:ALA:O	1:D:187:ALA:HB3	2.17	0.44
1:D:140:GLU:HB3	1:D:141:PRO:HD2	2.00	0.44
1:B:290:LEU:HD23	1:D:290:LEU:HD23	2.00	0.43
1:D:33:ALA:HB1	1:D:92:ILE:CG1	2.49	0.43
1:B:87:ARG:NH1	3:B:469:HOH:O	2.47	0.43
1:B:261:LEU:HD13	1:B:306:ILE:HG13	2.01	0.43
1:D:153:ARG:HH12	1:D:186:ALA:HB2	1.84	0.42
1:B:371:THR:CG2	1:C:349:GLY:HA2	2.49	0.42
1:D:50:THR:HG21	1:D:219:ILE:HG22	2.01	0.42
1:B:85:VAL:HG12	1:B:100[B]:MET:HE1	2.01	0.42
1:B:316:PHE:CE2	1:B:321:MET:HG3	2.54	0.42
1:C:243:LYS:HE2	1:C:243:LYS:HB3	1.79	0.41
1:D:83:GLY:HA3	1:D:258:TRP:CD1	2.55	0.41
1:D:123:TYR:CE2	1:D:196:ILE:HG21	2.55	0.41
1:A:15:GLU:OE2	1:A:23:ARG:HD2	2.21	0.41
1:B:182:TRP:CZ3	1:B:245:PRO:HD3	2.56	0.41
1:D:108:MET:HE3	1:D:128:VAL:CG2	2.50	0.41
1:D:182:TRP:CZ3	1:D:245:PRO:HD3	2.55	0.41
1:B:100[A]:MET:HE3	1:B:100[A]:MET:HB2	1.82	0.40
1:D:253:ARG:HE	1:D:373:GLU:HB3	1.86	0.40
1:A:90:GLU:HG3	1:A:94:SER:HA	2.03	0.40
1:D:15:GLU:OE2	1:D:23:ARG:HD2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/395 (89%)	345 (98%)	7 (2%)	0	100	100
1	B	367/395 (93%)	357 (97%)	10 (3%)	0	100	100
1	C	356/395 (90%)	346 (97%)	9 (2%)	1 (0%)	36	50
1	D	366/395 (93%)	356 (97%)	9 (2%)	1 (0%)	36	50
All	All	1441/1580 (91%)	1404 (97%)	35 (2%)	2 (0%)	48	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	143	ALA
1	C	373	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/313 (77%)	240 (100%)	1 (0%)	84	92
1	B	274/313 (88%)	274 (100%)	0	100	100
1	C	248/313 (79%)	246 (99%)	2 (1%)	73	86
1	D	283/313 (90%)	281 (99%)	2 (1%)	76	88
All	All	1046/1252 (84%)	1041 (100%)	5 (0%)	81	91

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	VAL
1	C	25	ILE
1	C	41	VAL
1	D	171	SER
1	D	232	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	343	GLN
1	C	364	GLN
1	D	273	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	402	-	3,3,3	0.59	0	2,2,2	0.04	0
2	EDO	D	402	-	3,3,3	0.49	0	2,2,2	0.32	0
2	EDO	D	401	-	3,3,3	0.49	0	2,2,2	0.25	0
2	EDO	C	401	-	3,3,3	0.82	0	2,2,2	0.31	0
2	EDO	A	401	-	3,3,3	0.41	0	2,2,2	0.47	0
2	EDO	D	403	-	3,3,3	0.42	0	2,2,2	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	402	-	-	0/1/1/1	-
2	EDO	D	402	-	-	0/1/1/1	-
2	EDO	D	401	-	-	0/1/1/1	-
2	EDO	C	401	-	-	1/1/1/1	-
2	EDO	A	401	-	-	0/1/1/1	-
2	EDO	D	403	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	401	EDO	O1-C1-C2-O2
2	D	403	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	362/395 (91%)	0.12	12 (3%) 49 45	25, 49, 88, 122	0
1	B	376/395 (95%)	-0.27	8 (2%) 63 59	22, 36, 64, 95	1 (0%)
1	C	365/395 (92%)	-0.04	10 (2%) 56 52	21, 44, 80, 110	1 (0%)
1	D	375/395 (94%)	-0.28	10 (2%) 56 52	21, 36, 65, 116	1 (0%)
All	All	1478/1580 (93%)	-0.12	40 (2%) 56 52	21, 40, 80, 122	3 (0%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	148	ALA	4.6
1	B	356	GLU	4.4
1	A	218	SER	4.0
1	D	356	GLU	4.0
1	D	143	ALA	3.4
1	D	145	SER	3.3
1	A	372	TYR	3.3
1	C	350	GLY	3.0
1	A	141	PRO	3.0
1	D	142	ASP	2.9
1	B	142	ASP	2.9
1	C	209	ILE	2.9
1	C	50	THR	2.7
1	C	349	GLY	2.6
1	C	374	GLY	2.5
1	B	357	TYR	2.5
1	D	349	GLY	2.5
1	C	4	ALA	2.5
1	A	282	ASN	2.4
1	D	372	TYR	2.4
1	B	148	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	373	GLU	2.4
1	D	50	THR	2.3
1	C	189	ASP	2.3
1	B	349	GLY	2.3
1	A	196	ILE	2.2
1	A	51	ASP	2.2
1	D	149	GLY	2.2
1	A	158	ASP	2.2
1	C	346	ASP	2.2
1	D	186	ALA	2.2
1	B	211	GLY	2.2
1	B	141	PRO	2.2
1	B	3	ARG	2.1
1	A	52	PRO	2.1
1	A	3	ARG	2.1
1	A	166	SER	2.0
1	C	164	SER	2.0
1	A	187	ALA	2.0
1	A	209	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	D	403	4/4	0.83	0.14	55,58,59,62	0
2	EDO	D	401	4/4	0.84	0.17	60,61,61,63	0
2	EDO	C	401	4/4	0.85	0.18	44,48,50,62	0
2	EDO	A	401	4/4	0.87	0.12	51,57,59,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	D	402	4/4	0.90	0.10	40,41,45,51	0
2	EDO	A	402	4/4	0.92	0.08	36,38,39,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.