



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 07:39 AM UTC

PDB ID : 6W9T / pdb_00006w9t
Title : Crystal structure of Neisseria meningitidis ClpP protease complex with small molecule activator ACP1-06
Authors : Mabanglo, M.F.; Houry, W.A.
Deposited on : 2020-03-23
Resolution : 1.64 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	199	9% 80% 11% 10%
1	B	199	7% 78% 13% 10%
1	C	199	10% 78% 13% 10%
1	D	199	11% 79% 12% 9%
1	E	199	8% 71% 16% 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	199	12% 77% 12% • 10%
1	G	199	3% 72% 15% • 13%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 20410 atoms, of which 9866 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	180	Total	C	H	N	O	S	0	0	0
			2820	887	1415	239	271	8			
1	B	180	Total	C	H	N	O	S	0	0	0
			2829	893	1419	239	270	8			
1	C	180	Total	C	H	N	O	S	0	0	0
			2820	887	1415	239	271	8			
1	D	182	Total	C	H	N	O	S	0	0	0
			2864	902	1437	244	273	8			
1	E	174	Total	C	H	N	O	S	0	0	0
			2731	859	1368	233	263	8			
1	F	179	Total	C	H	N	O	S	0	0	0
			2810	887	1408	238	269	8			
1	G	174	Total	C	H	N	O	S	0	0	0
			2731	859	1368	233	263	8			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	165	ASP	GLY	conflict	UNP A0A0Y5K536
B	165	ASP	GLY	conflict	UNP A0A0Y5K536
C	165	ASP	GLY	conflict	UNP A0A0Y5K536
D	165	ASP	GLY	conflict	UNP A0A0Y5K536
E	165	ASP	GLY	conflict	UNP A0A0Y5K536
F	165	ASP	GLY	conflict	UNP A0A0Y5K536
G	165	ASP	GLY	conflict	UNP A0A0Y5K536

- Molecule 2 is POTASSIUM ION (CCD ID: K) (formula: K) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	K	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total K 1 1	0	0
2	C	1	Total K 1 1	0	0
2	D	1	Total K 1 1	0	0
2	E	1	Total K 1 1	0	0
2	F	1	Total K 1 1	0	0
2	G	1	Total K 1 1	0	0

- Molecule 3 is N-{2-[(2-chlorophenyl)sulfanyl]ethyl}-2-methyl-2-[[5-(trifluoromethyl)pyridin-2-yl]sulfonyl]propanamide (CCD ID: KHS) (formula: C₁₈H₁₈ClF₃N₂O₃S₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms								ZeroOcc	AltConf
3	F	1	Total	C	Cl	F	H	N	O	S	0	1
			94	36	2	6	36	4	6	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	94	Total O 94 94	0	0


Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	104	Total 104	O 104	0	0
4	C	86	Total 86	O 86	0	0
4	D	107	Total 107	O 107	0	0
4	E	93	Total 93	O 93	0	0
4	F	99	Total 99	O 99	0	0
4	G	121	Total 121	O 121	0	0





- Molecule 1: ATP-dependent Clp protease proteolytic subunit



- Molecule 1: ATP-dependent Clp protease proteolytic subunit



- Molecule 1: ATP-dependent Clp protease proteolytic subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	97.33Å 119.15Å 127.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	97.33 – 1.64 97.33 – 1.64	Depositor EDS
% Data completeness (in resolution range)	98.2 (97.33-1.64) 98.8 (97.33-1.64)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.16 (at 1.63Å)	Xtriage
Refinement program	PHENIX 1.17.1 _3660	Depositor
R, R_{free}	0.237 , 0.282 0.242 , 0.287	Depositor DCC
R_{free} test set	2010 reflections (1.10%)	wwPDB-VP
Wilson B-factor (Å ²)	21.9	Xtriage
Anisotropy	0.455	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	20410	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KHS, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/1424	0.72	0/1917
1	B	0.60	1/1430 (0.1%)	0.70	1/1925 (0.1%)
1	C	0.55	1/1424 (0.1%)	0.69	1/1917 (0.1%)
1	D	0.50	0/1447	0.68	0/1947
1	E	0.53	0/1382	0.75	2/1859 (0.1%)
1	F	0.64	3/1422 (0.2%)	0.76	2/1914 (0.1%)
1	G	0.60	1/1382 (0.1%)	0.78	3/1859 (0.2%)
All	All	0.57	6/9911 (0.1%)	0.73	9/13338 (0.1%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	85	MET	SD-CE	-11.99	1.49	1.79
1	B	85	MET	SD-CE	-9.72	1.55	1.79
1	C	188	GLU	CD-OE1	-9.33	1.07	1.25
1	G	181	MET	SD-CE	-6.29	1.63	1.79
1	F	188	GLU	CD-OE1	5.84	1.36	1.25

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	85	MET	CG-SD-CE	11.23	125.60	100.90
1	G	173	ARG	CG-CD-NE	8.23	130.11	112.00
1	B	85	MET	CG-SD-CE	8.09	118.70	100.90
1	C	188	GLU	CG-CD-OE2	7.00	134.49	118.40
1	E	181	MET	CG-SD-CE	6.50	115.21	100.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1405	1415	1415	19	1
1	B	1410	1419	1419	24	0
1	C	1405	1415	1415	18	0
1	D	1427	1437	1437	20	0
1	E	1363	1368	1368	28	1
1	F	1402	1408	1408	21	0
1	G	1363	1368	1368	21	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
3	F	58	36	0	4	0
4	A	94	0	0	1	0
4	B	104	0	0	2	0
4	C	86	0	0	1	1
4	D	107	0	0	1	0
4	E	93	0	0	1	0
4	F	99	0	0	4	0
4	G	121	0	0	3	0
All	All	10544	9866	9830	136	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 136 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:8:VAL:N	4:D:401:HOH:O	2.07	0.86
1:F:67:TYR:OH	3:F:301[A]:KHS:OAR	1.95	0.84
1:D:194:GLN:NE2	1:D:202:LEU:O	2.13	0.82
1:B:194:GLN:OE1	4:B:401:HOH:O	1.99	0.81
1:E:187:LYS:HD2	1:E:195:ILE:HD12	1.64	0.80

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:SER:O	4:C:485:HOH:O[2_545]	2.11	0.09
1:E:166:ARG:HH22	1:G:31:GLU:OE1[3_545]	1.52	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	174/199 (87%)	171 (98%)	3 (2%)	0	100	100
1	B	174/199 (87%)	171 (98%)	3 (2%)	0	100	100
1	C	174/199 (87%)	171 (98%)	3 (2%)	0	100	100
1	D	176/199 (88%)	170 (97%)	4 (2%)	2 (1%)	11	1
1	E	170/199 (85%)	168 (99%)	2 (1%)	0	100	100
1	F	173/199 (87%)	168 (97%)	4 (2%)	1 (1%)	21	7
1	G	170/199 (85%)	167 (98%)	2 (1%)	1 (1%)	21	7
All	All	1211/1393 (87%)	1186 (98%)	21 (2%)	4 (0%)	36	20

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	202	LEU
1	F	200	ALA
1	D	200	ALA
1	G	200	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	155/168 (92%)	155 (100%)	0	100	100
1	B	155/168 (92%)	155 (100%)	0	100	100
1	C	155/168 (92%)	155 (100%)	0	100	100
1	D	157/168 (94%)	157 (100%)	0	100	100
1	E	149/168 (89%)	149 (100%)	0	100	100
1	F	154/168 (92%)	154 (100%)	0	100	100
1	G	149/168 (89%)	149 (100%)	0	100	100
All	All	1074/1176 (91%)	1074 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	46	ASN
1	D	121	ASN
1	D	194	GLN
1	F	46	ASN
1	F	121	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 7 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	KHS	F	301[B]	-	29,30,30	1.78	7 (24%)	32,45,45	3.36	6 (18%)
3	KHS	F	301[A]	-	29,30,30	1.83	7 (24%)	32,45,45	3.34	6 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KHS	F	301[B]	-	-	19/31/34/34	0/2/2/2
3	KHS	F	301[A]	-	-	17/31/34/34	0/2/2/2

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	301[A]	KHS	CAQ-NAS	5.26	1.46	1.33
3	F	301[B]	KHS	CAQ-NAS	5.07	1.45	1.33
3	F	301[A]	KHS	CAZ-SAV	4.30	1.83	1.77
3	F	301[B]	KHS	CAZ-SAV	3.68	1.82	1.77
3	F	301[A]	KHS	OAM-SAL	3.35	1.47	1.44

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	301[B]	KHS	OAK-SAL-OAM	-15.48	99.63	119.06
3	F	301[A]	KHS	OAK-SAL-OAM	-15.47	99.64	119.06
3	F	301[B]	KHS	SAL-CAJ-NAI	6.33	120.31	114.40
3	F	301[A]	KHS	SAL-CAJ-NAI	5.70	119.72	114.40
3	F	301[A]	KHS	CAH-CAJ-NAI	-4.23	120.89	125.39

There are no chirality outliers.

5 of 36 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	F	301[A]	KHS	NAS-CAT-CAU-SAV
3	F	301[A]	KHS	CAQ-CAO-SAL-OAM
3	F	301[A]	KHS	CAP-CAO-SAL-OAM
3	F	301[A]	KHS	CAP-CAO-SAL-OAK
3	F	301[A]	KHS	CAP-CAO-SAL-CAJ

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	301[B]	KHS	2	0
3	F	301[A]	KHS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	180/199 (90%)	0.48	17 (9%) 14 16	16, 25, 65, 85	0
1	B	180/199 (90%)	0.64	14 (7%) 19 22	19, 30, 53, 79	0
1	C	180/199 (90%)	0.96	19 (10%) 11 12	21, 35, 61, 81	0
1	D	182/199 (91%)	0.75	22 (12%) 8 10	17, 31, 68, 96	0
1	E	174/199 (87%)	0.63	15 (8%) 16 18	19, 30, 61, 87	0
1	F	179/199 (89%)	0.84	23 (12%) 7 9	20, 32, 73, 106	0
1	G	174/199 (87%)	0.20	6 (3%) 48 52	15, 25, 49, 63	0
All	All	1249/1393 (89%)	0.65	116 (9%) 14 16	15, 30, 61, 106	0

The worst 5 of 116 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	96	CYS	7.6
1	A	8	VAL	7.3
1	C	11	VAL	7.0
1	D	22	PHE	6.8
1	F	11	VAL	6.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	KHS	F	301[A]	29/29	0.65	0.25	53,71,89,93	47
3	KHS	F	301[B]	29/29	0.65	0.25	24,73,88,94	47
2	K	C	301	1/1	0.97	0.10	35,35,35,35	0
2	K	D	301	1/1	0.97	0.06	30,30,30,30	0
2	K	B	301	1/1	0.98	0.05	29,29,29,29	0
2	K	E	301	1/1	0.98	0.06	26,26,26,26	0
2	K	G	301	1/1	0.99	0.03	21,21,21,21	0
2	K	A	301	1/1	0.99	0.03	22,22,22,22	0
2	K	F	302	1/1	0.99	0.05	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around KHS F 301 (A):

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around KHS F 301 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K E 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K G 301:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around K A 301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.