



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 10:53 AM UTC

PDB ID : 8W9Z / pdb_00008w9z
EMDB ID : EMD-37386
Title : The cryo-EM structure of the Nicotiana tabacum PEP-PAP
Authors : Wu, X.X.; Zhang, Y.
Deposited on : 2023-09-06
Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14081 (2.50 - 3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	
1	a	337	
2	B	1070	
3	C	688	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	c	1388	
5	D	892	
6	E	860	
7	F	682	
8	G	266	
9	H	531	
10	I	486	
11	i	648	
12	J	507	
13	K	331	
14	L	303	
15	M	178	
15	m	178	
16	N	770	
17	O	167	
18	P	143	

2 Entry composition [i](#)

There are 21 unique types of molecules in this entry. The entry contains 60525 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	293	Total	C	N	O	S	0	0
			2202	1397	382	411	12		
1	a	318	Total	C	N	O	S	0	0
			2441	1555	424	452	10		

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	909	Total	C	N	O	S	0	0
			6820	4328	1216	1253	23		

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit gamma.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	592	Total	C	N	O	S	0	0
			4431	2832	783	799	17		

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit beta”.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	c	1174	Total	C	N	O	S	0	0
			8086	5072	1482	1503	29		

- Molecule 5 is a protein called PAP1(pTAC3).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	D	645	Total	C	N	O	S	0	0
			5088	3228	878	956	26		

- Molecule 6 is a protein called Pentatricopeptide repeat-containing protein At1g74850, chloroplastic-like.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	E	700	Total	C	N	O	S	0	0
			4534	2823	822	866	23		

- Molecule 7 is a protein called Protein PLASTID TRANSCRIPTIONALLY ACTIVE 10-like.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	F	544	Total	C	N	O	S	0	0
			4233	2678	756	780	19		

- Molecule 8 is a protein called superoxide dismutase.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	G	218	Total	C	N	O	S	0	0
			1727	1122	291	309	5		

- Molecule 9 is a protein called Protein PLASTID TRANSCRIPTIONALLY ACTIVE 12-like.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	H	260	Total	C	N	O	S	0	0
			2110	1338	371	393	8		

- Molecule 10 is a protein called Fructokinase-like 1, chloroplastic.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	I	374	Total	C	N	O	S	0	0
			2897	1856	512	517	12		

- Molecule 11 is a protein called Fructokinase-like 2, chloroplastic.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	i	383	Total	C	N	O	S	0	0
			2983	1902	508	557	16		

- Molecule 12 is a protein called Protein PLASTID TRANSCRIPTIONALLY ACTIVE 14-like isoform X2.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	J	422	Total	C	N	O	S	0	0
			3368	2157	578	612	21		

- Molecule 13 is a protein called PAP8(pTAC6).

Mol	Chain	Residues	Atoms					AltConf	Trace
13	K	214	Total	C	N	O	S	0	0
			1779	1127	307	337	8		

- Molecule 14 is a protein called superoxide dismutase.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	L	234	Total	C	N	O	S	0	0
			1475	943	261	270	1		

- Molecule 15 is a protein called Thioredoxin-like protein CITRX1, chloroplastic.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	M	116	Total	C	N	O	S	0	0
			935	598	150	180	7		
15	m	109	Total	C	N	O	S	0	0
			851	545	135	164	7		

- Molecule 16 is a protein called UDP-N-acetylmuramoyl-L-alanyl-D-glutamate--2, 6-diaminopimelate ligase-like.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	N	537	Total	C	N	O	S	0	0
			2982	1814	579	583	6		

- Molecule 17 is a protein called Protein PLASTID TRANSCRIPTIONALLY ACTIVE 7-like isoform X1.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	O	88	Total	C	N	O	S	0	0
			692	434	124	131	3		

- Molecule 18 is a protein called PAP13(pTAC18).

Mol	Chain	Residues	Atoms					AltConf	Trace
18	P	105	Total	C	N	O	S	0	0
			887	576	148	158	5		

- Molecule 19 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
19	B	1	Total	Zn	0
			1	1	

- Molecule 20 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
20	C	1	Total	Mg	0
			1	1	


- Molecule 21 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		AltConf
21	G	1	Total	Fe	0
			1	1	
21	L	1	Total	Fe	0
			1	1	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase subunit alpha




- Molecule 1: DNA-directed RNA polymerase subunit alpha




- Molecule 2: DNA-directed RNA polymerase subunit beta






• Molecule 3: DNA-directed RNA polymerase subunit gamma


Chain C: 69% 16% 14%



• Molecule 4: DNA-directed RNA polymerase subunit beta"


Chain c: 73% 11% 15%






• Molecule 5: PAP1(pTAC3)

Chain D: 60% 12% 28%




• Molecule 6: Pentatricopeptide repeat-containing protein At1g74850, chloroplastic-like

Chain E: 65% 16% 19%



● Molecule 7: Protein PLASTID TRANSCRIPTIONALLY ACTIVE 10-like

Chain F: 69% 11% 20%




● Molecule 8: superoxide dismutase

Chain G: 72% 10% 18%




- Molecule 9: Protein PLASTID TRANSCRIPTIONALLY ACTIVE 12-like

Chain H: 46% . 51%





- Molecule 10: Fructokinase-like 1, chloroplastic


Chain I: 68% 9% 23%




- Molecule 11: Fructokinase-like 2, chloroplastic

Chain i:  50% 8% 41%







- Molecule 12: Protein PLASTID TRANSCRIPTIONALLY ACTIVE 14-like isoform X2




- Molecule 13: PAP8(pTAC6)




- Molecule 14: superoxide dismutase






- Molecule 15: Thioredoxin-like protein CITRX1, chloroplastic

Chain M: 54% 10% 35%






- Molecule 15: Thioredoxin-like protein CITRX1, chloroplastic

Chain m: 53% 7% 39%



- Molecule 16: UDP-N-acetylmuramoyl-L-alanyl-D-glutamate--2, 6-diaminopimelate ligase-like

Chain N: 65% 5% 30%



HIS

- Molecule 17: Protein PLASTID TRANSCRIPTIONALLY ACTIVE 7-like isoform X1

Chain O: 41% 12% 47%

MET
ALA
SER
THR
LEU
THR
PHE
SER
GLY
PHE
SER
THR
LEU
GLN
THR
PHE
PRO
LYS
TLE
ALA
ALA
VAL
GLU
ASN
THR
LYS
ARG
VAL
ASN
PHE
SER
TLE
ARG
SER
GLN
SER
ALA
SER
ASN
SER
LYS
ASN
GLU
SER
GLY
GLY
ARG
ARG
TLE
TRP
ARG
ARG
LYS
LEU
THR
LYS
LYS

ASP
GLU
THR
LEU
ASP
ALA
LYS
MET
GLU
R54
I55
P56
R57
L58
Q61
I65
G68
G69
K70
L71
L72
T73
H74
N85
V90
I93
E96
N104
A110
R121
E131
A132
D137
P141
GLY
PRO
GLY
TYR
MET
LYS
ASP
GLU
LEU
LEU

● Molecule 18: PAP13(pTAC18)

Chain P: 59% 15% 27%

MET
ALA
SER
PHE
TLE
THR
MET
PRO
ALA
LEU
SER
TYR
LEU
SER
THR
ASN
THR
SER
SER
LEU
GLU
ARG
THR
ASN
PHE
ARG
PRO
SER
SER
ARG
GLY
TYR
GLN
GLY
VAL
ARG
ALA
R38
E41
V50
R51
V52
D59
R60
L61
V66
W72
K73
W83
L88
V96

V99
Q104
R105
F106
M107
V110
A111
G112
L125
Y135
R138
D142
SER

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	410910	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.856	Depositor
Minimum map value	-0.729	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.050	Depositor
Recommended contour level	0.1	Depositor
Map size ($\text{\AA}$)	440.0, 440.0, 440.0	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.1, 1.1, 1.1	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FE, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.32	0/2237	0.39	0/3034
1	a	0.34	0/2487	0.43	2/3374 (0.1%)
2	B	0.22	2/6945 (0.0%)	0.35	3/9418 (0.0%)
3	C	0.16	0/4518	0.36	2/6148 (0.0%)
4	c	0.38	0/8213	0.49	10/11205 (0.1%)
5	D	0.29	1/5191 (0.0%)	0.37	3/7029 (0.0%)
6	E	0.55	0/4604	0.73	20/6285 (0.3%)
7	F	0.40	0/4339	0.55	7/5885 (0.1%)
8	G	0.55	0/1777	0.53	2/2417 (0.1%)
9	H	0.26	0/2167	0.34	0/2942
10	I	0.43	0/2970	0.41	2/4026 (0.0%)
11	i	0.40	0/3056	0.44	0/4141
12	J	0.31	1/3454 (0.0%)	0.38	2/4689 (0.0%)
13	K	0.18	0/1824	0.28	0/2468
14	L	1.11	2/1514 (0.1%)	1.33	22/2100 (1.0%)
15	M	0.60	0/951	0.60	3/1286 (0.2%)
15	m	0.51	0/865	0.68	3/1175 (0.3%)
16	N	0.08	0/3011	0.23	0/4163
17	O	0.09	0/702	0.24	0/945
18	P	0.09	0/915	0.28	0/1241
All	All	0.39	6/61740 (0.0%)	0.49	81/83971 (0.1%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	249	CYS	C-O	6.53	1.31	1.24
14	L	155	PHE	CA-C	-5.46	1.45	1.52
2	B	76	TYR	CA-C	-5.30	1.46	1.52
2	B	77	SER	CA-C	-5.16	1.46	1.52
12	J	399	TYR	CA-C	-5.05	1.46	1.52

The worst 5 of 81 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	L	152	PHE	N-CA-C	-11.11	92.30	109.52
6	E	364	ASP	N-CA-C	-10.35	100.96	114.31
6	E	365	GLN	N-CA-C	-10.15	100.30	111.36
14	L	114	PRO	N-CA-C	9.89	122.77	110.70
4	c	401	PRO	N-CA-C	9.74	122.58	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2202	0	2108	23	0
1	a	2441	0	2390	29	0
2	B	6820	0	6495	77	0
3	C	4431	0	4202	86	0
4	c	8086	0	7028	99	0
5	D	5088	0	4957	77	0
6	E	4534	0	3525	68	0
7	F	4233	0	3825	48	0
8	G	1727	0	1648	15	0
9	H	2110	0	1974	11	0
10	I	2897	0	2752	25	0
11	i	2983	0	2896	30	0
12	J	3368	0	3240	38	0
13	K	1779	0	1730	31	0
14	L	1475	0	1046	15	0
15	M	935	0	934	14	0
15	m	851	0	837	7	0
16	N	2982	0	1849	18	0
17	O	692	0	660	18	0
18	P	887	0	826	11	0
19	B	1	0	0	0	0
20	C	1	0	0	0	0
21	G	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
21	L	1	0	0	0	0
All	All	60525	0	54922	651	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 651 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:D:379:ARG:HH21	5:D:477:ARG:HH21	1.02	0.93
5:D:379:ARG:NH2	5:D:477:ARG:HH21	1.74	0.85
5:D:379:ARG:HH21	5:D:477:ARG:NH2	1.84	0.74
2:B:625:GLN:HE22	2:B:646:GLY:H	1.35	0.73
8:G:235:PHE:HA	8:G:239:LEU:HB2	1.70	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/337 (86%)	277 (96%)	12 (4%)	0	100	100
1	a	314/337 (93%)	303 (96%)	11 (4%)	0	100	100
2	B	897/1070 (84%)	862 (96%)	35 (4%)	0	100	100
3	C	584/688 (85%)	555 (95%)	29 (5%)	0	100	100
4	c	1160/1388 (84%)	1124 (97%)	36 (3%)	0	100	100
5	D	639/892 (72%)	620 (97%)	19 (3%)	0	100	100
6	E	698/860 (81%)	680 (97%)	18 (3%)	0	100	100
7	F	542/682 (80%)	529 (98%)	13 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
8	G	216/266 (81%)	211 (98%)	5 (2%)	0	100	100
9	H	258/531 (49%)	254 (98%)	4 (2%)	0	100	100
10	I	372/486 (76%)	349 (94%)	23 (6%)	0	100	100
11	i	381/648 (59%)	365 (96%)	16 (4%)	0	100	100
12	J	420/507 (83%)	404 (96%)	16 (4%)	0	100	100
13	K	212/331 (64%)	207 (98%)	5 (2%)	0	100	100
14	L	230/303 (76%)	220 (96%)	9 (4%)	1 (0%)	30	65
15	M	114/178 (64%)	112 (98%)	2 (2%)	0	100	100
15	m	107/178 (60%)	100 (94%)	7 (6%)	0	100	100
16	N	535/770 (70%)	521 (97%)	14 (3%)	0	100	100
17	O	86/167 (52%)	84 (98%)	2 (2%)	0	100	100
18	P	103/143 (72%)	99 (96%)	4 (4%)	0	100	100
All	All	8157/10762 (76%)	7876 (97%)	280 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
14	L	134	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/308 (72%)	222 (100%)	1 (0%)	84	90
1	a	257/308 (83%)	251 (98%)	6 (2%)	44	74
2	B	667/924 (72%)	661 (99%)	6 (1%)	70	85
3	C	429/612 (70%)	421 (98%)	8 (2%)	50	76
4	c	679/1238 (55%)	665 (98%)	14 (2%)	47	75
5	D	528/770 (69%)	522 (99%)	6 (1%)	65	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
6	E	300/741 (40%)	291 (97%)	9 (3%)	36	69
7	F	395/615 (64%)	392 (99%)	3 (1%)	73	86
8	G	172/232 (74%)	169 (98%)	3 (2%)	53	78
9	H	212/472 (45%)	210 (99%)	2 (1%)	70	85
10	I	287/437 (66%)	284 (99%)	3 (1%)	68	84
11	i	316/567 (56%)	308 (98%)	8 (2%)	42	72
12	J	352/449 (78%)	346 (98%)	6 (2%)	53	78
13	K	199/300 (66%)	197 (99%)	2 (1%)	68	84
14	L	83/258 (32%)	74 (89%)	9 (11%)	6	26
15	M	103/160 (64%)	103 (100%)	0	100	100
15	m	92/160 (58%)	89 (97%)	3 (3%)	33	67
16	N	95/659 (14%)	91 (96%)	4 (4%)	26	61
17	O	69/144 (48%)	69 (100%)	0	100	100
18	P	91/129 (70%)	89 (98%)	2 (2%)	45	74
All	All	5549/9483 (58%)	5454 (98%)	95 (2%)	52	78

5 of 95 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
9	H	354	THR
12	J	245	GLN
10	I	323	VAL
11	i	531	VAL
13	K	173	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 152 such sidechains are listed below:

Mol	Chain	Res	Type
9	H	353	GLN
15	M	155	ASN
10	I	193	GLN
11	i	607	GLN
17	O	99	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-37386. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 250




Y Index: 223




Z Index: 184


6.3.2 Raw map



X Index: 250



Y Index: 223




Z Index: 184


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1962 nm^3 ; this corresponds to an approximate mass of 1772 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.333 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.333 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.00	-	-
Author-provided FSC curve	3.03	3.41	3.08
Unmasked-calculated*	3.65	4.20	3.68

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.65 differs from the reported value 3.0 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-37386 and PDB model 8W9Z. Per-residue inclusion information can be found in section [3](#) on page [8](#).


9.1 Map-model overlay [i](#)



X



Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.1 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.1).

9.4 Atom inclusion [i](#)



At the recommended contour level, 100% of all backbone atoms, 99% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9910	 0.4050
A	 0.9810	 0.4340
B	 0.9900	 0.4720
C	 0.9950	 0.3800
D	 0.9950	 0.3680
E	 0.9750	 0.1700
F	 0.9960	 0.4270
G	 0.9940	 0.4700
H	 0.9920	 0.5000
I	 0.9930	 0.5180
J	 0.9940	 0.4570
K	 0.9910	 0.4560
L	 1.0000	 0.4250
M	 0.9900	 0.5070
N	 1.0000	 0.1620
O	 0.9930	 0.4450
P	 0.9930	 0.4410
a	 0.9910	 0.4690
c	 0.9910	 0.4090
i	 0.9950	 0.4700
m	 0.9950	 0.4400

