



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 02:34 AM UTC

PDB ID : 2WAH / pdb_00002wah
Title : Crystal Structure of an IgG1 Fc Glycoform (Man9GlcNAc2)
Authors : Crispin, M.; Bowden, T.A.; Coles, C.H.; Harlos, K.; Aricescu, A.R.; Harvey, D.J.; Stuart, D.I.; Jones, E.Y.
Deposited on : 2009-02-06
Resolution : 2.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	
1	B	209	
2	C	9	
3	D	3	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3565 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called IG GAMMA-1 CHAIN C REGION.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	0	0
			1658	1056	279	317	6			
1	B	209	Total	C	N	O	S	0	0	0
			1669	1063	281	319	6			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	393	ALA	THR	engineered mutation	UNP P01857
B	393	ALA	THR	engineered mutation	UNP P01857

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	9	Total	C	N	O	0	0	0
			105	58	2	45			

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	3	Total	C	N	O	0	0	0
			39	22	2	15			



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	54	Total	O	0	0
			54	54		
4	B	40	Total	O	0	0
			40	40		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: IG GAMMA-1 CHAIN C REGION



- Molecule 1: IG GAMMA-1 CHAIN C REGION



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.39Å 74.98Å 149.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.87 – 2.51 29.87 – 2.51	Depositor EDS
% Data completeness (in resolution range)	97.5 (29.87-2.51) 97.5 (29.87-2.51)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.5.0047	Depositor
R, R_{free}	0.216 , 0.268 (Not available) , 0.307	Depositor DCC
R_{free} test set	988 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	40.1	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3565	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.15 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.4513e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, MAN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/1704	0.75	1/2321 (0.0%)
1	B	0.61	0/1716	0.74	0/2339
All	All	0.64	0/3420	0.74	1/4660 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	391	TYR	N-CA-C	5.36	117.12	109.14

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	389	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1658	0	1626	10	1
1	B	1669	0	1635	14	1
2	C	105	0	88	0	0
3	D	39	0	34	0	0
4	A	54	0	0	0	0
4	B	40	0	0	0	0
All	All	3565	0	3383	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:240:VAL:HG22	1:B:263:VAL:HG22	1.69	0.72
1:B:350:THR:HB	1:B:441:LEU:HD22	1.77	0.65
1:A:355:ARG:HA	1:A:358:LEU:HD23	1.87	0.56
1:B:353:PRO:HD3	1:B:365:LEU:HD12	1.94	0.49
1:B:368:LEU:HD13	1:B:407:TYR:CZ	2.47	0.49
1:A:238:PRO:HG2	1:A:328:LEU:HD13	1.96	0.47
1:B:406:LEU:HD12	1:B:406:LEU:C	2.40	0.47
1:A:242:LEU:HD12	1:A:260:THR:O	2.14	0.47
1:B:418:GLN:HA	1:B:443:LEU:HD22	1.98	0.46
1:B:332:ILE:HD11	1:B:334:LYS:NZ	2.30	0.46
1:B:346:PRO:HB3	1:B:372:PHE:HB3	1.97	0.45
1:B:271:PRO:O	1:B:272:GLU:C	2.59	0.44
1:A:242:LEU:HD11	1:A:259:VAL:CG1	2.48	0.44
1:B:422:VAL:HG22	1:B:442:SER:OG	2.18	0.44
1:A:406:LEU:C	1:A:406:LEU:HD12	2.44	0.43
1:B:367:CYS:HB2	1:B:381:TRP:CZ2	2.54	0.43
1:A:388:GLU:HB3	1:A:410:LEU:HD11	2.00	0.43
1:A:257:PRO:HG2	1:A:308:VAL:O	2.20	0.42
1:A:344:ARG:NH1	1:A:403:SER:HB3	2.35	0.42
1:A:391:TYR:HB2	1:A:409:LYS:O	2.20	0.42
1:A:343:PRO:HA	1:A:373:TYR:O	2.20	0.41
1:B:388:GLU:CD	1:B:416:ARG:HH22	2.28	0.41
1:B:443:LEU:HD12	1:B:444:SER:N	2.36	0.41
1:B:393:ALA:HA	1:B:408:SER:HA	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:389:ASN:OD1	1:B:389:ASN:O[3_555]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/209 (98%)	199 (97%)	6 (3%)	0	100	100
1	B	207/209 (99%)	199 (96%)	7 (3%)	1 (0%)	24	43
All	All	412/418 (99%)	398 (97%)	13 (3%)	1 (0%)	43	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	329	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/194 (100%)	189 (98%)	4 (2%)	47	74
1	B	194/194 (100%)	187 (96%)	7 (4%)	31	58
All	All	387/388 (100%)	376 (97%)	11 (3%)	38	66

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	252	MET
1	A	270	ASP
1	A	384	ASN
1	A	406	LEU
1	B	267	SER
1	B	273	VAL
1	B	299	THR
1	B	360	LYS
1	B	365	LEU
1	B	406	LEU
1	B	444	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	311	GLN
1	A	315	ASN
1	A	325	ASN
1	A	421	ASN
1	B	433	HIS
1	B	435	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2,1	14,14,15	0.41	0	17,19,21	1.28	1 (5%)
2	NAG	C	2	2	14,14,15	0.58	0	17,19,21	0.84	0
2	BMA	C	3	2	11,11,12	0.42	0	15,15,17	0.59	0
2	MAN	C	4	2	11,11,12	0.54	0	15,15,17	1.16	1 (6%)
2	MAN	C	5	2	11,11,12	0.54	0	15,15,17	0.88	0
2	MAN	C	6	2	11,11,12	0.57	0	15,15,17	1.06	1 (6%)
2	MAN	C	7	2	11,11,12	0.51	0	15,15,17	0.69	1 (6%)
2	MAN	C	8	2	11,11,12	0.51	0	15,15,17	1.04	1 (6%)
2	MAN	C	9	2	11,11,12	0.55	0	15,15,17	1.04	1 (6%)
3	NAG	D	1	3,1	14,14,15	0.35	0	17,19,21	1.50	1 (5%)
3	NAG	D	2	3	14,14,15	0.74	0	17,19,21	1.04	0
3	BMA	D	3	3	11,11,12	0.36	0	15,15,17	0.86	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	MAN	C	5	2	-	2/2/19/22	0/1/1/1
2	MAN	C	6	2	-	0/2/19/22	0/1/1/1
2	MAN	C	7	2	-	0/2/19/22	0/1/1/1
2	MAN	C	8	2	-	0/2/19/22	0/1/1/1
2	MAN	C	9	2	-	2/2/19/22	0/1/1/1
3	NAG	D	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
3	BMA	D	3	3	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	C1-O5-C5	4.94	118.81	112.19
2	C	4	MAN	C1-O5-C5	3.37	116.71	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	9	MAN	C1-O5-C5	3.10	116.34	112.19
2	C	6	MAN	C1-O5-C5	2.91	116.09	112.19
2	C	1	NAG	C8-C7-N2	2.90	120.93	116.12
2	C	8	MAN	C1-O5-C5	2.70	115.80	112.19
3	D	3	BMA	C1-O5-C5	2.32	115.30	112.19
2	C	7	MAN	C1-O5-C5	2.02	114.89	112.19

There are no chirality outliers.


All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	5	MAN	O5-C5-C6-O6
3	D	3	BMA	O5-C5-C6-O6
2	C	9	MAN	O5-C5-C6-O6
2	C	5	MAN	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
3	D	1	NAG	C8-C7-N2-C2
3	D	1	NAG	O7-C7-N2-C2
2	C	9	MAN	C4-C5-C6-O6
3	D	3	BMA	C4-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/209 (99%)	0.44	9 (4%) 40 35	16, 22, 27, 31	0
1	B	209/209 (100%)	1.05	43 (20%) 2 2	14, 21, 26, 40	0
All	All	416/418 (99%)	0.74	52 (12%) 8 6	14, 21, 27, 40	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	296	TYR	4.6
1	B	315	ASN	4.2
1	B	251	LEU	3.9
1	B	311	GLN	3.8
1	B	338	LYS	3.7
1	B	272	GLU	3.5
1	B	298	SER	3.5
1	B	273	VAL	3.5
1	B	268	HIS	3.4
1	B	314	LEU	3.3
1	A	253	ILE	3.3
1	B	316	GLY	3.2
1	B	271	PRO	3.1
1	B	254	SER	3.1
1	B	324	SER	3.1
1	A	330	ALA	2.9
1	A	255	ARG	2.9
1	B	313	TRP	2.9
1	B	267	SER	2.8
1	B	317	LYS	2.7
1	B	275	PHE	2.7
1	A	248	LYS	2.7
1	B	276	ASN	2.7
1	B	308	VAL	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	266	VAL	2.6
1	B	336	ILE	2.6
1	B	312	ASP	2.5
1	B	247	PRO	2.5
1	B	300	TYR	2.4
1	B	253	ILE	2.4
1	B	269	GLU	2.4
1	A	385	GLY	2.3
1	B	326	LYS	2.3
1	B	299	THR	2.3
1	B	385	GLY	2.3
1	B	256	THR	2.3
1	A	249	ASP	2.2
1	B	280	ASP	2.2
1	B	277	TRP	2.2
1	B	244	PRO	2.2
1	B	263	VAL	2.2
1	B	274	LYS	2.2
1	A	326	LYS	2.2
1	B	250	THR	2.1
1	B	319	TYR	2.1
1	A	251	LEU	2.1
1	B	309	LEU	2.1
1	B	292	ARG	2.0
1	B	330	ALA	2.0
1	A	254	SER	2.0
1	B	445	PRO	2.0
1	B	257	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MAN	C	9	11/12	0.25	0.17	84,84,85,85	0
3	BMA	D	3	11/12	0.35	0.17	81,82,82,82	0
2	MAN	C	7	11/12	0.36	0.18	80,81,82,82	0
2	MAN	C	4	11/12	0.53	0.18	75,76,77,79	0
2	MAN	C	8	11/12	0.53	0.15	77,79,80,82	0
3	NAG	D	2	14/15	0.58	0.20	72,74,77,79	0
3	NAG	D	1	14/15	0.63	0.16	58,65,66,70	0
2	MAN	C	6	11/12	0.68	0.19	78,79,79,80	0
2	MAN	C	5	11/12	0.75	0.14	76,77,78,78	0
2	NAG	C	2	14/15	0.78	0.13	53,58,61,64	0
2	NAG	C	1	14/15	0.80	0.12	48,53,55,55	0
2	BMA	C	3	11/12	0.81	0.12	67,69,73,74	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.