



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:06 AM UTC

PDB ID : 2WAJ / pdb_00002waj
Title : Crystal structure of human Jnk3 complexed with a 1-aryl-3,4- dihydroisoquinoline inhibitor
Authors : Bax, B.D.; Christopher, J.A.; Jones, E.J.; Mosley, J.E.
Deposited on : 2009-02-08
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3076 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MITOGEN-ACTIVATED PROTEIN KINASE 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	348	Total	C	N	O	S	0	8	0
			2878	1843	492	520	23			

- Molecule 2 is 1-(3-BROMOPHENYL)-7-CHLORO-6-METHOXY-3,4-DIHYDROISOQUINOLINE (CCD ID: SNB) (formula: C₁₆H₁₃BrClNO).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	Br	C	Cl	N	O	
			20	1	16	1	1	1	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	178	Total 178	0	0

- Molecule 1: MITOGEN-ACTIVATED PROTEIN KINASE 10



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.66Å 71.34Å 107.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40 20.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.8 (20.00-2.40) 98.5 (20.00-2.40)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.50 (at 2.41Å)	Xtriage
Refinement program	REFMAC 5.3.0006	Depositor
R, R_{free}	0.183 , 0.264 (Not available) , 0.257	Depositor DCC
R_{free} test set	799 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	30.2	Xtriage
Anisotropy	1.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3076	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SNB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/2940	0.95	9/3974 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	THR	CA-C-N	6.46	133.32	121.70
1	A	216	THR	C-N-CA	6.46	133.32	121.70
1	A	283	CYS	CA-C-N	6.08	127.43	119.84
1	A	283	CYS	C-N-CA	6.08	127.43	119.84
1	A	371	PRO	CA-C-N	5.58	126.13	120.38
1	A	371	PRO	C-N-CA	5.58	126.13	120.38
1	A	189	ASP	N-CA-C	5.38	117.33	110.61
1	A	364	ASP	CA-C-N	5.03	125.05	119.32
1	A	364	ASP	C-N-CA	5.03	125.05	119.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2878	0	2885	50	0
2	A	20	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	178	0	0	2	0
All	All	3076	0	2898	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:216:THR:HB	1:A:217:SER:HB3	1.16	1.11
1:A:216:THR:CB	1:A:217:SER:HB3	1.96	0.95
1:A:120:HIS:HD2	1:A:122:ASN:H	1.16	0.92
1:A:102:GLN:NE2	1:A:390:TRP:HE1	1.70	0.89
1:A:67:LEU:HD22	1:A:81:ALA:HB2	1.66	0.77
1:A:377:ASP:O	1:A:378:LYS:HB2	1.85	0.77
1:A:53:VAL:HG11	1:A:79[A]:CYS:SG	2.25	0.77
1:A:216:THR:N	1:A:217:SER:HG	1.86	0.74
1:A:173:MET:O	1:A:177:ILE:HD12	1.92	0.69
1:A:102:GLN:NE2	1:A:385:HIS:O	2.28	0.66
1:A:97:ARG:NH2	1:A:140:GLN:OE1	2.31	0.63
1:A:102:GLN:HE22	1:A:385:HIS:H	1.46	0.61
1:A:383:ARG:HE	1:A:385:HIS:HE1	1.47	0.60
1:A:115[B]:MET:HE3	1:A:126:LEU:HD11	1.84	0.60
1:A:120:HIS:CD2	1:A:122:ASN:H	2.08	0.60
1:A:380:LEU:O	1:A:381:ASP:HB2	2.01	0.60
1:A:371:PRO:O	1:A:373:PRO:HD3	2.02	0.59
1:A:216:THR:CA	1:A:217:SER:HB3	2.34	0.57
1:A:377:ASP:O	1:A:378:LYS:CB	2.52	0.57
1:A:110:ARG:HD2	1:A:381:ASP:OD1	2.06	0.56
1:A:315:ASP:OD1	1:A:333:ARG:NH2	2.38	0.56
1:A:233:GLU:HB2	1:A:238:MET:HB2	1.89	0.55
1:A:102:GLN:NE2	1:A:390:TRP:NE1	2.50	0.53
1:A:110:ARG:HA	1:A:380:LEU:HD11	1.90	0.52
1:A:170:LEU:HA	1:A:173:MET:HE3	1.92	0.52
1:A:111:GLU:O	1:A:115[B]:MET:HB2	2.11	0.51
1:A:360:ASN:C	1:A:360:ASN:HD22	2.19	0.50
1:A:297:TYR:CE1	1:A:301:ARG:HD3	2.48	0.48
1:A:107:ARG:NH1	1:A:207:ASP:OD2	2.47	0.48
1:A:356:HIS:ND1	1:A:357:PRO:HD2	2.29	0.48
1:A:360:ASN:C	1:A:360:ASN:ND2	2.72	0.48
1:A:212:ARG:HD3	1:A:212:ARG:C	2.38	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:ALA:HA	1:A:247:TRP:CD1	2.49	0.47
1:A:287:MET:HA	1:A:290:LEU:HD12	1.97	0.47
1:A:246:ILE:HD11	1:A:350:VAL:HA	1.97	0.47
1:A:121:LYS:HE2	1:A:367:GLU:HG2	1.96	0.46
1:A:227:ARG:HD3	1:A:268:TYR:CE1	2.50	0.46
1:A:93:LYS:HE2	3:A:2029:HOH:O	2.16	0.46
1:A:371:PRO:HA	1:A:372:PRO:HD3	1.72	0.44
1:A:178:LYS:HD2	3:A:2088:HOH:O	2.17	0.44
1:A:76:GLY:HA3	1:A:93:LYS:HE3	2.00	0.43
1:A:142:VAL:HG22	1:A:398:VAL:HG21	2.01	0.43
1:A:119:ASN:H	1:A:179:HIS:HE1	1.66	0.42
1:A:102:GLN:O	1:A:106:LYS:HB2	2.20	0.42
1:A:307:LEU:HD12	1:A:312:LEU:HD13	2.02	0.42
1:A:373:PRO:HB2	1:A:374:GLN:CB	2.50	0.41
1:A:168:TYR:HD2	1:A:201[B]:CYS:SG	2.43	0.41
1:A:96:SER:O	1:A:97:ARG:C	2.64	0.41
1:A:188:ARG:NH2	1:A:211:ALA:O	2.54	0.41
1:A:114:LEU:HD22	1:A:185:ILE:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	239/364 (66%)	225 (94%)	10 (4%)	4 (2%)	7 10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	378	LYS
1	A	381	ASP
1	A	372	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	284	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/329 (97%)	304 (95%)	16 (5%)	22	38

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	LYS
1	A	95	LEU
1	A	96	SER
1	A	118	VAL
1	A	121	LYS
1	A	137	GLU
1	A	140	GLN
1	A	153	LEU
1	A	212	ARG
1	A	220	MET
1	A	262	LEU
1	A	303	LYS
1	A	312	LEU
1	A	360	ASN
1	A	378	LYS
1	A	384	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	ASN
1	A	89	ASN
1	A	102	GLN
1	A	104	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	119	ASN
1	A	120	HIS
1	A	122	ASN
1	A	155	GLN
1	A	243	ASN
1	A	355	GLN
1	A	360	ASN
1	A	385	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SNB	A	1401	-	21,22,22	0.78	1 (4%)	30,31,31	2.08	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SNB	A	1401	-	-	0/6/16/16	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1401	SNB	C16-CL	2.21	1.78	1.73

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1401	SNB	C6-N-C7	9.50	128.29	116.21
2	A	1401	SNB	C8-C7-N	3.59	121.73	117.69
2	A	1401	SNB	O-C2-C16	2.51	118.99	116.62
2	A	1401	SNB	C5-C6-N	-2.02	107.41	114.35


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	348/364 (95%)	0.19	12 (3%) 48 44	25, 51, 84, 109	8 (2%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	374	GLN	4.5
1	A	373	PRO	4.0
1	A	223	TYR	4.0
1	A	370	ALA	3.7
1	A	117	CYS	3.6
1	A	224	VAL	3.4
1	A	222	PRO	2.8
1	A	221	THR	2.7
1	A	216	THR	2.7
1	A	73	GLY	2.6
1	A	371	PRO	2.2
1	A	220	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SNB	A	1401	20/20	0.99	0.07	35,40,42,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.