



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:51 PM UTC

PDB ID : 5WAT / pdb_00005wat
Title : Corynebacterium glutamicum Full length Homoserine kinase
Authors : Petit, C.; Ronning, D.R.
Deposited on : 2017-06-27
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	7% 81% 17% ..
1	B	315	19% 83% 14% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	403	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4862 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Homoserine kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	8	0
			2340	1472	414	451	3			
1	B	306	Total	C	N	O	S	0	2	0
			2245	1418	392	432	3			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	310	HIS	-	expression tag	UNP P07128
A	311	HIS	-	expression tag	UNP P07128
A	312	HIS	-	expression tag	UNP P07128
A	313	HIS	-	expression tag	UNP P07128
A	314	HIS	-	expression tag	UNP P07128
A	315	HIS	-	expression tag	UNP P07128
B	310	HIS	-	expression tag	UNP P07128
B	311	HIS	-	expression tag	UNP P07128
B	312	HIS	-	expression tag	UNP P07128
B	313	HIS	-	expression tag	UNP P07128
B	314	HIS	-	expression tag	UNP P07128
B	315	HIS	-	expression tag	UNP P07128

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is HEXAETHYLENE GLYCOL (CCD ID: P6G) (formula: C₁₂H₂₆O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			19	12	7		
3	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	150	Total	O	0	0
			150	150		
6	B	49	Total	O	0	0
			49	49		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Homoserine kinase



• Molecule 1: Homoserine kinase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	68.08Å 130.08Å 195.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.34 – 2.14 42.34 – 2.14	Depositor EDS
% Data completeness (in resolution range)	99.5 (42.34-2.14) 95.4 (42.34-2.14)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.75 (at 2.14Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.189 , 0.225 0.195 , 0.229	Depositor DCC
R_{free} test set	2409 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtriage
Anisotropy	0.543	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4862	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P6G, SO4, PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2396	0.86	3/3277 (0.1%)
1	B	0.44	0/2290	0.84	1/3133 (0.0%)
All	All	0.54	0/4686	0.85	4/6410 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	135	GLU	N-CA-C	5.79	118.06	111.11
1	B	259	TYR	N-CA-C	5.46	118.35	108.69
1	A	137	HIS	CA-C-N	5.12	125.15	119.32
1	A	137	HIS	C-N-CA	5.12	125.15	119.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2340	0	2330	39	0
1	B	2245	0	2240	24	0
2	A	15	0	0	2	0
2	B	15	0	0	1	0
3	A	32	0	43	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	10	0	0	1	0
4	B	5	0	0	0	0
5	A	1	0	0	0	0
6	A	150	0	0	4	0
6	B	49	0	0	3	0
All	All	4862	0	4613	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:ARG:HH12	3:A:404:P6G:H112	1.42	0.84
1:A:292:LYS:NZ	4:A:406:SO4:O4	2.14	0.80
1:A:163:GLN:OE1	6:A:501:HOH:O	2.00	0.80
1:A:52:VAL:HG21	1:A:65:SER:HB3	1.65	0.78
1:A:297:GLU:HG2	3:A:404:P6G:H61	1.68	0.75
1:A:9:ARG:HB2	1:A:43:ILE:HD12	1.69	0.73
1:A:240:ARG:NH1	2:A:403:PO4:O3	2.20	0.73
1:B:15:VAL:HG21	1:B:109:ALA:HB2	1.72	0.70
1:A:178:ARG:NH1	3:A:404:P6G:H112	2.08	0.69
1:A:98:ARG:NH2	1:A:185:ASN:O	2.26	0.68
1:A:178:ARG:HH12	3:A:404:P6G:C11	2.05	0.68
1:A:277:PRO:HB3	3:A:404:P6G:H141	1.74	0.67
1:A:189:SER:HB3	1:A:244:LEU:HD21	1.77	0.65
2:B:402:PO4:O4	6:B:501:HOH:O	2.14	0.65
1:A:172:GLU:O	1:A:221[A]:GLN:HG2	1.97	0.64
1:B:191:GLU:HG2	1:B:192:ALA:H	1.61	0.63
1:A:12:THR:HB	1:A:305:GLU:CG	2.29	0.63
1:B:191:GLU:CG	1:B:192:ALA:H	2.12	0.62
1:A:245:PRO:O	1:A:249:GLU:HG3	2.02	0.59
1:A:222:GLN:HG3	1:A:223:ARG:HG2	1.83	0.59
1:B:95:PRO:HG2	1:B:100:LEU:HD12	1.85	0.59
1:A:178:ARG:HH22	3:A:404:P6G:H142	1.70	0.56
1:A:160:GLY:C	1:A:162:SER:HB3	2.31	0.55
1:B:202:THR:HG23	1:B:205:ASP:H	1.71	0.54
1:A:253:ARG:O	1:A:257[A]:ARG:HG3	2.07	0.54
1:A:4:GLU:OE1	1:A:82:GLU:HG3	2.09	0.53
1:A:98:ARG:NH1	1:A:187:HIS:HA	2.24	0.53
1:A:12:THR:HB	1:A:305:GLU:HG2	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:HIS:HD2	6:A:540:HOH:O	1.92	0.52
1:B:182:LEU:HD22	1:B:293:VAL:HG22	1.92	0.52
1:B:46:SER:O	6:B:502:HOH:O	2.19	0.51
1:B:184:PRO:HG2	1:B:268:PRO:HB2	1.93	0.50
1:A:59:GLU:OE2	6:A:503:HOH:O	2.19	0.50
1:B:196:VAL:HB	1:B:239:TYR:HB3	1.94	0.50
1:B:9:ARG:HB3	1:B:43:ILE:HD12	1.93	0.49
1:A:205:ASP:OD2	6:A:504:HOH:O	2.20	0.48
1:B:98:ARG:HA	1:B:268:PRO:HD2	1.96	0.48
1:A:55:GLU:N	1:A:62:LEU:HD11	2.29	0.48
1:B:66:HIS:NE2	1:B:70:LYS:HD2	2.29	0.47
1:B:182:LEU:HB3	1:B:291:ILE:HG21	1.96	0.47
1:A:198:PRO:HG3	1:A:239:TYR:CZ	2.50	0.47
1:A:62:LEU:HD22	1:A:92:ASN:HB2	1.97	0.47
1:A:161:LYS:N	1:A:162:SER:HB3	2.31	0.46
1:B:139:ASP:N	1:B:139:ASP:OD1	2.50	0.45
1:A:163:GLN:HG3	1:A:164:PRO:HD2	1.99	0.45
1:A:51:GLU:OE2	1:A:87:ARG:NH2	2.45	0.45
1:B:162:SER:OG	1:B:163:GLN:N	2.50	0.45
1:B:96:GLN:OE1	6:B:503:HOH:O	2.20	0.45
3:A:404:P6G:H112	3:A:404:P6G:H142	1.70	0.44
1:B:153:TRP:HD1	1:B:156:LEU:HD13	1.83	0.43
1:B:250:TRP:CZ2	1:B:286:ALA:HA	2.54	0.43
1:A:190:THR:OG1	1:A:191:GLU:N	2.51	0.43
1:B:172:GLU:O	1:B:221:GLN:HG2	2.19	0.43
1:B:13:VAL:HG22	1:B:304:VAL:HG22	2.01	0.42
1:A:96:GLN:HG2	1:A:102:SER:OG	2.19	0.42
1:A:125[A]:GLU:H	1:A:125[A]:GLU:CD	2.28	0.42
1:A:251:VAL:O	1:A:255:ARG:HG3	2.19	0.42
1:B:158:ILE:HA	1:B:159:ASP:HA	1.88	0.41
1:A:191:GLU:OE1	1:A:191:GLU:HA	2.19	0.41
1:B:191:GLU:CG	1:B:192:ALA:N	2.83	0.41
1:B:28:THR:HG23	1:B:152:SER:O	2.20	0.41
1:A:216:MET:HE1	1:A:273:LEU:HD12	2.02	0.41
1:A:178:ARG:NH1	3:A:404:P6G:C11	2.75	0.41
1:A:266:ALA:HB2	2:A:403:PO4:O4	2.21	0.41
1:B:98:ARG:O	1:B:269:THR:OG1	2.30	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/315 (100%)	303 (96%)	12 (4%)	1 (0%)	36	33
1	B	304/315 (96%)	282 (93%)	21 (7%)	1 (0%)	36	33
All	All	620/630 (98%)	585 (94%)	33 (5%)	2 (0%)	36	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	160	GLY
1	A	160	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/251 (98%)	243 (98%)	4 (2%)	55	61
1	B	233/251 (93%)	225 (97%)	8 (3%)	32	32
All	All	480/502 (96%)	468 (98%)	12 (2%)	40	44

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	GLN
1	A	193	VAL
1	A	281	LYS
1	A	289	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	3	ILE
1	B	15	VAL
1	B	163	GLN
1	B	173	VAL
1	B	183	VAL
1	B	193	VAL
1	B	197	LEU
1	B	217	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	187	HIS
1	A	307	ASN
1	B	92	ASN
1	B	174	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	P6G	A	404	-	18,18,18	0.68	0	17,17,17	0.55	0
4	SO4	B	404	-	4,4,4	0.23	0	6,6,6	0.12	0
2	PO4	B	403	-	4,4,4	0.89	0	6,6,6	0.80	0
4	SO4	A	406	-	4,4,4	0.24	0	6,6,6	0.07	0
2	PO4	B	402	-	4,4,4	0.99	0	6,6,6	0.69	0
4	SO4	A	405	-	4,4,4	0.21	0	6,6,6	0.18	0
2	PO4	A	403	-	4,4,4	1.08	0	6,6,6	0.58	0
2	PO4	A	402	-	4,4,4	1.08	0	6,6,6	0.41	0
2	PO4	B	401	-	4,4,4	1.02	0	6,6,6	0.54	0
3	P6G	A	408	-	12,12,18	0.52	0	11,11,17	0.41	0
2	PO4	A	401	-	4,4,4	1.18	0	6,6,6	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	P6G	A	404	-	-	8/16/16/16	-
3	P6G	A	408	-	-	7/10/10/16	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	404	P6G	O7-C8-C9-O10
3	A	408	P6G	O7-C8-C9-O10
3	A	408	P6G	O13-C14-C15-O16
3	A	404	P6G	O13-C14-C15-O16
3	A	408	P6G	O16-C17-C18-O19
3	A	404	P6G	O4-C5-C6-O7
3	A	404	P6G	O1-C2-C3-O4
3	A	404	P6G	C11-C12-O13-C14
3	A	408	P6G	C14-C15-O16-C17
3	A	408	P6G	C11-C12-O13-C14
3	A	404	P6G	C5-C6-O7-C8

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	408	P6G	C15-C14-O13-C12
3	A	404	P6G	C15-C14-O13-C12
3	A	408	P6G	C12-C11-O10-C9
3	A	404	P6G	C6-C5-O4-C3

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	404	P6G	8	0
4	A	406	SO4	1	0
2	B	402	PO4	1	0
2	A	403	PO4	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	310/315 (98%)	0.22	23 (7%) 20 24	14, 34, 76, 106	8 (2%)
1	B	306/315 (97%)	1.31	59 (19%) 3 4	20, 58, 96, 122	2 (0%)
All	All	616/630 (97%)	0.76	82 (13%) 7 8	14, 48, 93, 122	10 (1%)

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	154	THR	8.3
1	A	188	ALA	6.4
1	B	193	VAL	6.2
1	B	158	ILE	6.0
1	B	192	ALA	5.6
1	B	196	VAL	5.2
1	B	309	PRO	5.1
1	A	192	ALA	5.0
1	B	156	LEU	4.7
1	B	258	GLY	4.6
1	A	158	ILE	4.4
1	B	243	VAL	4.2
1	A	187	HIS	4.2
1	B	188	ALA	4.2
1	B	135	GLU	4.1
1	B	250	TRP	4.0
1	B	290	GLY	4.0
1	B	185	ASN	3.9
1	B	155	ASN	3.8
1	B	137	HIS	3.8
1	A	190	THR	3.8
1	A	157	SER	3.7
1	B	190	THR	3.6
1	B	246	ILE	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	135	GLU	3.4
1	B	171	LEU	3.4
1	B	136	GLY	3.2
1	B	138	PRO	3.2
1	A	310	HIS	3.2
1	A	161	LYS	3.1
1	A	185	ASN	3.1
1	B	201	VAL	3.1
1	B	278	ILE	3.1
1	B	291	ILE	3.1
1	B	157	SER	3.0
1	B	202	THR	3.0
1	A	156	LEU	3.0
1	B	172	GLU	2.9
1	B	189	SER	2.9
1	B	160	GLY	2.9
1	A	193	VAL	2.8
1	B	247	THR	2.8
1	B	164	PRO	2.8
1	B	184	PRO	2.8
1	A	194	ARG	2.8
1	B	194	ARG	2.8
1	B	259	TYR	2.7
1	B	159	ASP	2.7
1	B	234	ARG	2.7
1	B	284	GLU	2.6
1	B	281	LYS	2.6
1	A	243	VAL	2.5
1	B	239	TYR	2.5
1	B	256	ASN	2.5
1	A	58	GLY	2.5
1	B	251	VAL	2.4
1	B	204	ILE	2.3
1	B	242	GLU	2.3
1	B	285	ASP	2.3
1	B	195	ARG	2.3
1	A	186	PHE	2.3
1	B	286	ALA	2.3
1	B	228	TRP	2.3
1	A	246	ILE	2.2
1	B	257	ARG	2.2
1	B	307	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	197	LEU	2.2
1	A	62	LEU	2.2
1	B	245	PRO	2.1
1	B	2	ALA	2.1
1	B	226	LEU	2.1
1	B	177	ILE	2.1
1	B	308	GLN	2.1
1	B	253	ARG	2.1
1	A	191	GLU	2.1
1	B	127	ILE	2.1
1	B	133	ALA	2.1
1	B	260	ALA	2.1
1	A	137	HIS	2.0
1	A	54	GLY	2.0
1	A	159	ASP	2.0
1	A	56	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	B	403	5/5	0.58	0.19	49,59,65,71	5
2	PO4	A	402	5/5	0.75	0.24	40,47,55,59	5
4	SO4	B	404	5/5	0.83	0.10	71,78,95,98	0
2	PO4	B	402	5/5	0.84	0.14	52,56,59,64	5
5	MG	A	407	1/1	0.84	0.15	58,58,58,58	0
2	PO4	A	403	5/5	0.86	0.21	43,45,56,57	5
3	P6G	A	408	13/19	0.86	0.16	56,64,72,77	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	B	401	5/5	0.88	0.11	58,60,72,74	0
4	SO4	A	406	5/5	0.90	0.09	54,69,70,75	5
4	SO4	A	405	5/5	0.93	0.10	42,50,57,63	5
3	P6G	A	404	19/19	0.93	0.10	29,44,59,63	0
2	PO4	A	401	5/5	0.96	0.07	35,38,47,48	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around P6G A 408:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around P6G A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.