



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:02 AM UTC

PDB ID : 1WBH / pdb_00001wbh
Title : Crystal structure of the E45N mutant from KDPG aldolase from Escherichia coli
Authors : Fullerton, S.W.B.; Merkel, A.B.; Naismith, J.H.
Deposited on : 2004-11-01
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	4% 90% 7%
1	B	214	5% 89% 11%
1	C	214	7% 86% 11%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5440 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called KHG/KDPG ALDOLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	213	Total	C	N	O	S	26	0	0
			1565	1005	261	292	7			
1	B	214	Total	C	N	O	S	5	0	0
			1570	1008	262	293	7			
1	C	213	Total	C	N	O	S	36	0	0
			1565	1005	261	292	7			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	45	ASN	GLU	engineered mutation	UNP P10177
B	45	ASN	GLU	engineered mutation	UNP P10177
C	45	ASN	GLU	engineered mutation	UNP P10177

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	232	Total	O	0	0
			232	232		
3	B	246	Total	O	0	0
			246	246		
3	C	242	Total	O	0	0
			242	242		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: KHG/KDPG ALDOLASE



- Molecule 1: KHG/KDPG ALDOLASE



- Molecule 1: KHG/KDPG ALDOLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.49Å 84.24Å 132.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.16 – 1.55 40.16 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.0 (40.16-1.55) 98.0 (40.16-1.55)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.2.0007	Depositor
R, R_{free}	0.168 , 0.217 0.172 , 0.219	Depositor DCC
R_{free} test set	4409 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	14.3	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5440	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	3/1593 (0.2%)	1.00	2/2167 (0.1%)
1	B	0.93	2/1598 (0.1%)	1.01	3/2174 (0.1%)
1	C	0.95	3/1593 (0.2%)	1.06	4/2167 (0.2%)
All	All	0.95	8/4784 (0.2%)	1.03	9/6508 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	2	LYS	CA-C	13.15	1.70	1.52
1	A	89	GLN	CG-CD	10.56	1.78	1.52
1	C	212	LYS	CG-CD	-10.11	1.22	1.52
1	A	89	GLN	CD-OE1	8.53	1.39	1.23
1	A	89	GLN	CB-CG	6.73	1.72	1.52
1	B	151	GLY	N-CA	6.67	1.53	1.44
1	B	133	LYS	CG-CD	6.38	1.71	1.52
1	C	151	GLY	N-CA	5.94	1.52	1.44

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	212	LYS	CB-CG-CD	14.02	143.54	111.30
1	C	212	LYS	CG-CD-CE	10.06	134.44	111.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	89	GLN	CB-CG-CD	9.53	128.80	112.60
1	C	151	GLY	CA-C-N	9.09	131.20	119.84
1	C	151	GLY	C-N-CA	9.09	131.20	119.84
1	B	133	LYS	CB-CG-CD	-7.50	94.05	111.30
1	B	133	LYS	CG-CD-CE	-5.96	97.60	111.30
1	A	82	GLU	CB-CG-CD	-5.66	102.98	112.60
1	B	145	ALA	N-CA-C	-5.39	105.31	111.14

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	2	LYS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1565	0	1632	16	0
1	B	1570	0	1637	21	0
1	C	1565	0	1632	22	0
2	A	5	0	0	0	0
2	B	10	0	0	0	0
2	C	5	0	0	0	0
3	A	232	0	0	7	0
3	B	246	0	0	12	0
3	C	242	0	0	10	0
All	All	5440	0	4901	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:GLN:CD	1:A:89:GLN:CG	1.78	1.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:102:LYS:HD3	3:C:2151:HOH:O	1.33	1.28
1:C:5:LYS:HE2	3:C:2010:HOH:O	1.05	1.23
1:B:106:GLU:HG2	3:B:2152:HOH:O	1.43	1.18
1:C:209:GLU:HB3	3:C:2233:HOH:O	1.50	1.11
1:A:198:ASP:HB3	3:A:2208:HOH:O	1.63	0.98
1:C:145:ALA:O	1:C:149:ILE:N	2.03	0.92
1:A:66:GLU:HG2	3:A:2103:HOH:O	1.68	0.90
1:C:62:LYS:HG3	3:C:2098:HOH:O	1.77	0.85
1:C:5:LYS:CE	3:C:2010:HOH:O	1.84	0.77
1:A:205:ARG:HG3	3:A:2221:HOH:O	1.86	0.75
1:C:42:ARG:HD2	1:C:68:ILE:HG12	1.68	0.75
1:A:66:GLU:HG2	3:A:2047:HOH:O	1.90	0.72
1:C:146:LEU:O	1:C:150:ALA:HB3	1.91	0.70
1:B:130:LYS:HE3	3:B:2019:HOH:O	1.92	0.69
1:A:202:LYS:HG2	1:A:205:ARG:HH22	1.57	0.68
1:C:102:LYS:HE3	3:C:2170:HOH:O	1.94	0.68
1:C:209:GLU:HG2	3:C:2226:HOH:O	1.94	0.66
1:B:0:ALA:HB1	3:B:2075:HOH:O	1.98	0.62
1:C:42:ARG:HD3	3:C:2109:HOH:O	2.00	0.60
1:A:66:GLU:CG	3:A:2103:HOH:O	2.38	0.59
1:B:1:MET:CE	1:B:105:THR:HG22	2.36	0.55
1:B:186:LEU:HB2	3:B:2217:HOH:O	2.05	0.55
1:B:78:GLN:HG2	3:B:2130:HOH:O	2.07	0.53
1:B:0:ALA:CB	3:B:2075:HOH:O	2.54	0.53
1:C:150:ALA:HB1	3:C:2191:HOH:O	2.08	0.52
1:B:149:ILE:HD11	3:B:2165:HOH:O	2.08	0.52
1:B:94:PRO:O	1:C:118:VAL:HG21	2.10	0.51
1:B:1:MET:HE2	1:B:105:THR:HG22	1.91	0.51
1:C:42:ARG:HD2	1:C:68:ILE:CG1	2.40	0.51
1:C:42:ARG:CD	3:C:2109:HOH:O	2.57	0.51
1:A:19:PRO:HD2	1:A:43:VAL:O	2.12	0.49
1:A:114:GLY:HA2	1:A:133:LYS:HB3	1.95	0.49
1:B:193:GLU:HG3	3:B:2220:HOH:O	2.13	0.48
1:A:136:PRO:HG2	1:A:139:ALA:HB3	1.94	0.48
1:C:19:PRO:HD2	1:C:43:VAL:O	2.12	0.48
1:B:209:GLU:HG2	3:B:2229:HOH:O	2.13	0.48
1:A:118:VAL:HG21	1:C:94:PRO:O	2.15	0.47
1:B:170:ARG:NH1	1:B:212:LYS:O	2.48	0.47
1:C:130:LYS:HG3	1:C:131:GLU:OE2	2.15	0.46
1:B:193:GLU:CG	3:B:2220:HOH:O	2.63	0.46
1:A:205:ARG:CZ	3:A:2212:HOH:O	2.64	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:MET:HE3	1:B:1:MET:HB3	1.77	0.46
1:A:89:GLN:CG	1:A:89:GLN:NE2	2.67	0.44
1:B:19:PRO:HD2	1:B:43:VAL:O	2.18	0.44
1:A:9:GLU:OE2	3:A:2021:HOH:O	2.21	0.43
1:B:121:LEU:HD11	1:B:156:VAL:HG11	1.99	0.43
1:C:32:MET:HB2	1:C:192:LEU:HD11	2.01	0.43
1:C:190:ASP:HA	1:C:193:GLU:CD	2.43	0.43
1:C:181:ILE:HD12	1:C:181:ILE:C	2.44	0.43
1:B:187:VAL:HB	3:B:2037:HOH:O	2.20	0.42
1:A:95:GLY:HA2	1:A:114:GLY:O	2.20	0.42
1:A:129:LEU:HD12	1:A:129:LEU:N	2.35	0.41
1:B:1:MET:HE1	1:B:105:THR:HG22	2.03	0.41
1:B:1:MET:HE1	1:B:105:THR:CG2	2.51	0.41
1:C:165:SER:HB2	1:C:166:PRO:CD	2.51	0.41
1:B:102:LYS:HD2	3:B:2176:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/214 (99%)	207 (98%)	4 (2%)	0	100	100
1	B	212/214 (99%)	209 (99%)	3 (1%)	0	100	100
1	C	211/214 (99%)	206 (98%)	4 (2%)	1 (0%)	24	8
All	All	634/642 (99%)	622 (98%)	11 (2%)	1 (0%)	43	24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	151	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/163 (100%)	156 (96%)	7 (4%)	26	4
1	B	163/163 (100%)	162 (99%)	1 (1%)	78	64
1	C	163/163 (100%)	158 (97%)	5 (3%)	35	8
All	All	489/489 (100%)	476 (97%)	13 (3%)	39	12

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	2	LYS
1	A	9	GLU
1	A	66	GLU
1	A	169	TYR
1	A	192	LEU
1	A	198	ASP
1	B	169	TYR
1	C	2	LYS
1	C	42	ARG
1	C	66	GLU
1	C	169	TYR
1	C	192	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	HIS
1	A	78	GLN
1	B	89	GLN
1	B	147	GLN
1	C	28	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	C	1214	-	4,4,4	0.95	0	6,6,6	1.23	1 (16%)
2	PO4	B	1215	-	4,4,4	1.30	0	6,6,6	0.49	0
2	PO4	B	1214	-	4,4,4	1.02	0	6,6,6	0.62	0
2	PO4	A	1214	-	4,4,4	1.18	0	6,6,6	0.78	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	C	1214	PO4	O4-P-O2	2.31	115.08	107.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	212/214 (99%)	0.02	9 (4%)	40	48	3, 16, 28, 40	3 (1%)
1	B	214/214 (100%)	0.22	11 (5%)	33	40	10, 18, 30, 38	2 (0%)
1	C	211/214 (98%)	0.27	15 (7%)	22	25	7, 18, 30, 39	4 (1%)
All	All	637/642 (99%)	0.17	35 (5%)	30	37	3, 17, 30, 40	9 (1%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	150	ALA	6.6
1	C	149	ILE	5.5
1	C	151	GLY	5.5
1	B	151	GLY	4.8
1	B	149	ILE	4.8
1	B	0	ALA	3.8
1	C	148	ALA	3.8
1	A	205	ARG	3.7
1	B	213	LEU	3.5
1	C	146	LEU	3.3
1	C	192	LEU	3.2
1	C	42	ARG	3.2
1	B	150	ALA	3.1
1	A	155	GLN	2.8
1	B	155	GLN	2.7
1	B	106	GLU	2.7
1	B	5	LYS	2.7
1	A	194	ALA	2.5
1	C	24	LYS	2.5
1	B	193	GLU	2.4
1	A	193	GLU	2.4
1	C	62	LYS	2.3
1	A	139	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	194	ALA	2.3
1	C	190	ASP	2.3
1	C	212	LYS	2.3
1	A	192	LEU	2.3
1	A	150	ALA	2.2
1	A	3	ASN	2.2
1	C	63	GLU	2.2
1	B	28	HIS	2.2
1	B	205	ARG	2.1
1	A	149	ILE	2.1
1	C	193	GLU	2.0
1	C	139	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	B	1214	5/5	0.90	0.11	28,28,34,37	0
2	PO4	C	1214	5/5	0.92	0.12	23,26,29,29	0
2	PO4	A	1214	5/5	0.94	0.10	20,24,28,30	0
2	PO4	B	1215	5/5	0.99	0.05	14,17,18,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.