



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 12:42 PM UTC

PDB ID : 2WBY / pdb_00002wby
Title : Crystal structure of human insulin-degrading enzyme in complex with insulin
Authors : Manolopoulou, M.; Guo, Q.; Malito, E.; Schilling, A.B.; Tang, W.J.
Deposited on : 2009-03-06
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	990	10% 72%21%...
1	B	990	69%23%...
2	C	20	10% 75%25%
2	E	20	5% 60%30%10%
3	D	19	26% 63%21%11%5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	19	 A horizontal bar chart showing the quality of chain F. The bar is divided into four segments: red (37%), green (58%), yellow (37%), and orange (5%). The segments are labeled with their respective percentages: 37%, 58%, 37%, and 5%.

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ZN	A	3012	-	-	X	-
4	ZN	B	3012	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 16926 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INSULIN-DEGRADING ENZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	958	Total	C	N	O	S	0	0	1
			7813	5032	1313	1445	23			
1	B	955	Total	C	N	O	S	0	0	1
			7787	5018	1307	1440	22			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	110	LEU	CYS	engineered mutation	UNP P14735
A	111	GLN	GLU	engineered mutation	UNP P14735
A	171	SER	CYS	engineered mutation	UNP P14735
A	178	ALA	CYS	engineered mutation	UNP P14735
A	257	VAL	CYS	engineered mutation	UNP P14735
A	414	LEU	CYS	engineered mutation	UNP P14735
A	573	ASN	CYS	engineered mutation	UNP P14735
A	590	SER	CYS	engineered mutation	UNP P14735
A	789	SER	CYS	engineered mutation	UNP P14735
A	812	ALA	CYS	engineered mutation	UNP P14735
A	819	ALA	CYS	engineered mutation	UNP P14735
A	904	SER	CYS	engineered mutation	UNP P14735
A	966	ASN	CYS	engineered mutation	UNP P14735
A	974	ALA	CYS	engineered mutation	UNP P14735
B	110	LEU	CYS	engineered mutation	UNP P14735
B	111	GLN	GLU	engineered mutation	UNP P14735
B	171	SER	CYS	engineered mutation	UNP P14735
B	178	ALA	CYS	engineered mutation	UNP P14735
B	257	VAL	CYS	engineered mutation	UNP P14735
B	414	LEU	CYS	engineered mutation	UNP P14735
B	573	ASN	CYS	engineered mutation	UNP P14735
B	590	SER	CYS	engineered mutation	UNP P14735
B	789	SER	CYS	engineered mutation	UNP P14735
B	812	ALA	CYS	engineered mutation	UNP P14735
B	819	ALA	CYS	engineered mutation	UNP P14735

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	904	SER	CYS	engineered mutation	UNP P14735
B	966	ASN	CYS	engineered mutation	UNP P14735
B	974	ALA	CYS	engineered mutation	UNP P14735

- Molecule 2 is a protein called INSULIN A CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	20	Total	C	N	O	S	0	0	0
			155	95	23	33	4			
2	E	20	Total	C	N	O	S	0	0	0
			155	95	23	33	4			

- Molecule 3 is a protein called INSULIN B CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	18	Total	C	N	O	S	0	0	0
			143	94	24	24	1			
3	F	19	Total	C	N	O	S	0	0	0
			149	97	25	25	2			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		
4	B	1	Total	Zn	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	408	Total	O	0	0
			408	408		
5	B	295	Total	O	0	0
			295	295		
5	C	6	Total	O	0	0
			6	6		
5	D	4	Total	O	0	0
			4	4		
5	E	5	Total	O	0	0
			5	5		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	F	4	Total	O	0	0
			4	4		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: INSULIN-DEGRADING ENZYME





• Molecule 1: INSULIN-DEGRADING ENZYME







● Molecule 2: INSULIN A CHAIN



● Molecule 2: INSULIN A CHAIN



● Molecule 3: INSULIN B CHAIN



● Molecule 3: INSULIN B CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	262.32Å 262.32Å 90.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	32.08 – 2.60 32.08 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (32.08-2.60) 99.8 (32.08-2.60)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.39 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.164 , 0.218 0.168 , 0.220	Depositor DCC
R_{free} test set	5438 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16926	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.35	22/8007 (0.3%)	1.30	40/10833 (0.4%)
1	B	1.34	24/7982 (0.3%)	1.29	45/10801 (0.4%)
2	C	1.53	0/156	1.52	0/209
2	E	1.48	1/156 (0.6%)	1.34	0/209
3	D	1.74	1/146 (0.7%)	1.50	0/198
3	F	1.67	3/152 (2.0%)	1.48	1/206 (0.5%)
All	All	1.36	51/16599 (0.3%)	1.30	86/22456 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	189	GLU	CD-OE2	10.00	1.44	1.25
1	B	764	VAL	CA-CB	9.84	1.65	1.53
1	A	189	GLU	CD-OE1	9.27	1.43	1.25
1	B	716	ILE	CA-CB	8.22	1.58	1.54
1	B	741	ILE	CA-CB	7.47	1.63	1.54

The worst 5 of 86 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	52	ASN	N-CA-C	18.75	133.90	111.02
1	A	189	GLU	CG-CD-OE2	-12.84	88.88	118.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	636	ASP	N-CA-C	12.59	127.87	111.75
1	A	189	GLU	CG-CD-OE1	11.78	145.50	118.40
1	B	52	ASN	CB-CA-C	-10.74	93.84	110.92

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	979	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7813	0	7744	156	0
1	B	7787	0	7715	172	0
2	C	155	0	144	5	0
2	E	155	0	143	8	0
3	D	143	0	142	14	0
3	F	149	0	146	12	0
4	A	1	0	0	2	0
4	B	1	0	0	3	0
5	A	408	0	0	24	0
5	B	295	0	0	14	0
5	C	6	0	0	0	0
5	D	4	0	0	1	0
5	E	5	0	0	0	0
5	F	4	0	0	0	0
All	All	16926	0	16034	339	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 339 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:491:ARG:HG3	1:A:491:ARG:HH11	1.02	1.17
4:B:3012:ZN:ZN	3:F:1:PHE:N	1.11	1.14
4:A:3012:ZN:ZN	3:D:1:PHE:N	1.17	1.06
1:B:189:GLU:OE2	3:F:1:PHE:N	1.90	1.03
1:B:119:LYS:HE3	1:B:171:SER:HB2	1.44	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	954/990 (96%)	918 (96%)	36 (4%)	0	100	100
1	B	951/990 (96%)	921 (97%)	29 (3%)	1 (0%)	48	70
2	C	18/20 (90%)	16 (89%)	2 (11%)	0	100	100
2	E	18/20 (90%)	17 (94%)	1 (6%)	0	100	100
3	D	16/19 (84%)	12 (75%)	4 (25%)	0	100	100
3	F	17/19 (90%)	17 (100%)	0	0	100	100
All	All	1974/2058 (96%)	1901 (96%)	72 (4%)	1 (0%)	48	70

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1010	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	848/879 (96%)	758 (89%)	90 (11%)	6	14
1	B	845/879 (96%)	753 (89%)	92 (11%)	6	13
2	C	19/19 (100%)	17 (90%)	2 (10%)	6	14
2	E	19/19 (100%)	16 (84%)	3 (16%)	2	4
3	D	16/17 (94%)	14 (88%)	2 (12%)	4	9
3	F	17/17 (100%)	15 (88%)	2 (12%)	5	10
All	All	1764/1830 (96%)	1573 (89%)	191 (11%)	6	13

5 of 191 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	287	GLU
1	B	657	LYS
1	B	417	LEU
1	B	556	MET
1	B	728	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	783	ASN
1	B	805	ASN
1	B	93	HIS
1	B	783	ASN
1	B	922	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	958/990 (96%)	-0.86	7 (0%) 84 82	13, 24, 40, 71	0
1	B	955/990 (96%)	-0.73	10 (1%) 79 76	15, 28, 44, 78	0
2	C	20/20 (100%)	0.59	2 (10%) 12 9	22, 52, 60, 61	0
2	E	20/20 (100%)	0.68	1 (5%) 34 28	21, 46, 63, 67	0
3	D	18/19 (94%)	1.43	5 (27%) 1 1	26, 52, 56, 57	0
3	F	19/19 (100%)	1.47	7 (36%) 1 0	32, 49, 63, 64	0
All	All	1990/2058 (96%)	-0.73	32 (1%) 70 66	13, 26, 47, 78	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	965	SER	5.7
1	B	52	ASN	5.6
1	B	964	ASP	4.7
1	A	964	ASP	4.6
1	B	50	ILE	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ZN	A	3012	1/1	1.00	0.30	2,2,2,2	0
4	ZN	B	3012	1/1	1.00	0.30	2,2,2,2	0

6.5 Other polymers [i](#)

There are no such residues in this entry.