



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 10:47 PM UTC

PDB ID : 4WCX / pdb_00004wcx
Title : Crystal structure of HydG: A maturase of the [FeFe]-hydrogenase
Authors : Dinis, P.C.; Harmer, J.E.; Driesener, R.C.; Roach, P.L.
Deposited on : 2014-09-05
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	
1	C	480	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	H2S	A	506	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 15198 atoms, of which 7116 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Biotin and thiamin synthesis associated.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	454	Total	C	H	N	O	S	0	2	0
			7214	2305	3583	623	683	20			
1	C	450	Total	C	H	N	O	S	0	0	0
			7059	2261	3502	607	669	20			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP D3T7F1
A	-12	GLY	-	expression tag	UNP D3T7F1
A	-11	SER	-	expression tag	UNP D3T7F1
A	-10	SER	-	expression tag	UNP D3T7F1
A	-9	HIS	-	expression tag	UNP D3T7F1
A	-8	HIS	-	expression tag	UNP D3T7F1
A	-7	HIS	-	expression tag	UNP D3T7F1
A	-6	HIS	-	expression tag	UNP D3T7F1
A	-5	HIS	-	expression tag	UNP D3T7F1
A	-4	HIS	-	expression tag	UNP D3T7F1
A	-3	SER	-	expression tag	UNP D3T7F1
A	-2	GLN	-	expression tag	UNP D3T7F1
A	-1	ASP	-	expression tag	UNP D3T7F1
A	0	PRO	-	expression tag	UNP D3T7F1
C	-13	MET	-	initiating methionine	UNP D3T7F1
C	-12	GLY	-	expression tag	UNP D3T7F1
C	-11	SER	-	expression tag	UNP D3T7F1
C	-10	SER	-	expression tag	UNP D3T7F1
C	-9	HIS	-	expression tag	UNP D3T7F1
C	-8	HIS	-	expression tag	UNP D3T7F1
C	-7	HIS	-	expression tag	UNP D3T7F1
C	-6	HIS	-	expression tag	UNP D3T7F1
C	-5	HIS	-	expression tag	UNP D3T7F1
C	-4	HIS	-	expression tag	UNP D3T7F1
C	-3	SER	-	expression tag	UNP D3T7F1

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLN	-	expression tag	UNP D3T7F1
C	-1	ASP	-	expression tag	UNP D3T7F1
C	0	PRO	-	expression tag	UNP D3T7F1

- Molecule 2 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	C	1	Total	Fe	S	0	0
			8	4	4		
2	C	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 3 is METHIONINE (CCD ID: MET) (formula: $\text{C}_5\text{H}_{11}\text{NO}_2\text{S}$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	S	
			17	5	8	1	2	1	

- Molecule 4 is FE (III) ION (CCD ID: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Fe		
			1	1	0	0

- Molecule 5 is ALANINE (CCD ID: ALA) (formula: C₃H₇NO₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	0	0
			10	3	4	1	2		

- Molecule 6 is HYDROSULFURIC ACID (CCD ID: H2S) (formula: H₂S).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	S	0	0
			1	1		
6	C	1	Total	S	0	0
			1	1		

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Na	0	0
			1	1		
7	C	2	Total	Na	0	0
			2	2		

- Molecule 8 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: C₁₅H₂₂N₆O₅S).





Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
8	C	1	Total	C	H	N	O	S	0	0
			46	15	19	6	5	1		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	373	Total	O	0	0
			373	373		
9	C	441	Total	O	0	0
			441	441		

- Molecule 1: Biotin and thiamin synthesis associated



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.13Å 56.19Å 84.92Å 89.59° 83.62° 66.84°	Depositor
Resolution (Å)	84.33 – 1.59 84.33 – 1.59	Depositor EDS
% Data completeness (in resolution range)	96.2 (84.33-1.59) 96.2 (84.33-1.59)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 1.59Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.180 , 0.213 0.185 , 0.215	Depositor DCC
R_{free} test set	1995 reflections (1.62%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.416	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	15198	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAM, SF4, NA, H2S, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/3710	0.85	1/5008 (0.0%)
1	C	0.79	2/3625 (0.1%)	0.91	0/4897
All	All	0.76	2/7335 (0.0%)	0.88	1/9905 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	419	ILE	CA-CB	5.63	1.61	1.54
1	C	301	MET	SD-CE	-5.39	1.66	1.79

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	308	MET	CG-SD-CE	-5.08	89.72	100.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	387	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3631	3583	3576	31	1
1	C	3557	3502	3501	35	1
2	A	16	0	0	0	0
2	C	16	0	0	0	0
3	A	9	8	8	4	0
4	A	1	0	0	0	0
5	A	6	4	4	2	0
6	A	1	0	0	2	0
6	C	1	0	0	0	0
7	A	1	0	0	0	0
7	C	2	0	0	0	0
8	C	27	19	21	0	0
9	A	373	0	0	6	0
9	C	441	0	0	7	0
All	All	8082	7116	7110	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:505:ALA:HB3	6:A:506:H2S:S	1.57	1.42
5:A:505:ALA:CB	6:A:506:H2S:S	2.44	1.05
1:C:335:CYS:SG	9:C:1003:HOH:O	2.38	0.82
1:A:392:ARG:NH2	1:C:258:GLU:OE2	2.11	0.81
1:A:198:ARG:NH1	9:A:756:HOH:O	2.06	0.78
1:A:109:LYS:NZ	9:A:601:HOH:O	2.19	0.75
1:A:388:ARG:NH2	1:A:466:PHE:OXT	2.29	0.65
1:C:126:GLY:O	1:C:128:LYS:NZ	2.32	0.62
1:A:99:TYR:CD2	3:A:503:MET:HE1	2.36	0.61
1:A:385:ARG:NH1	9:A:880:HOH:O	2.34	0.61
1:C:272:LEU:HD23	1:C:286:ILE:HD11	1.84	0.60
1:C:287:VAL:CG1	1:C:291:ASP:HB2	2.33	0.58
1:A:70:LYS:NZ	1:A:327:ILE:O	2.35	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:254:ASN:ND2	9:C:993:HOH:O	2.39	0.56
1:A:112:MET:CE	1:A:148:ASP:HB3	2.39	0.53
1:C:20:LYS:NZ	9:C:809:HOH:O	2.40	0.53
1:A:100:ARG:NH1	9:A:603:HOH:O	2.41	0.53
1:C:305:TYR:HB3	1:C:394:MET:HE1	1.91	0.53
1:A:448[B]:GLU:HG3	1:A:452:ARG:NH1	2.25	0.51
1:C:323:ILE:HA	1:C:327:ILE:HB	1.92	0.51
1:C:122:LEU:HD23	1:C:125:MET:CE	2.42	0.50
1:C:397:ALA:HA	1:C:402:ILE:HD13	1.92	0.50
1:A:99:TYR:CE2	3:A:503:MET:HE1	2.46	0.50
1:C:287:VAL:HG13	1:C:291:ASP:HB2	1.94	0.50
1:A:283:PHE:O	1:A:286:ILE:HG12	2.12	0.50
1:A:33:ILE:CG2	1:A:65:VAL:HG21	2.43	0.48
1:C:280:ILE:HG22	1:C:283:PHE:CE1	2.48	0.48
1:A:112:MET:HE3	1:A:148:ASP:HB3	1.97	0.47
1:C:81:PHE:CZ	1:C:331:SER:HB2	2.49	0.47
1:C:401:GLN:NE2	9:C:605:HOH:O	2.45	0.47
1:C:424:GLU:OE1	9:C:763:HOH:O	2.20	0.47
1:A:33:ILE:HG22	1:A:65:VAL:HG21	1.95	0.47
1:A:20:LYS:HE2	1:A:45:GLU:HG3	1.98	0.46
1:A:133:GLU:HB3	3:A:503:MET:HG3	1.97	0.45
1:A:448[B]:GLU:HG3	1:A:452:ARG:HH12	1.81	0.45
1:A:112:MET:HE1	1:A:148:ASP:C	2.42	0.45
1:C:122:LEU:HD13	1:C:130:LEU:HD22	1.99	0.45
1:A:354:GLN:CD	3:A:503:MET:HE3	2.42	0.44
1:C:280:ILE:HA	1:C:283:PHE:CD1	2.53	0.44
1:C:300:ARG:NH2	1:C:308:MET:HG3	2.33	0.44
1:A:359:ASP:OD1	1:A:361:ARG:HD3	2.18	0.44
1:C:81:PHE:CZ	1:C:331:SER:CB	3.01	0.44
1:C:407:LEU:HB3	1:C:408:PRO:HD3	2.00	0.43
1:A:198:ARG:HB2	1:A:199:PRO:HD3	2.01	0.43
1:A:6:ALA:HB1	1:A:217:LEU:HD23	2.01	0.43
1:A:118:GLU:HA	1:A:121:ILE:HD12	2.01	0.43
1:C:112:MET:O	1:C:116:ARG:HG3	2.18	0.43
1:A:20:LYS:HE2	1:A:45:GLU:OE2	2.19	0.42
1:A:20:LYS:HE3	9:A:837:HOH:O	2.19	0.42
1:C:122:LEU:HD21	1:C:335:CYS:HB2	2.01	0.42
1:C:280:ILE:HA	1:C:283:PHE:HD1	1.82	0.42
1:A:20:LYS:HE2	1:A:45:GLU:CG	2.49	0.42
1:C:158:LEU:HD23	1:C:159:LYS:N	2.35	0.42
1:C:335:CYS:HB3	9:C:1000:HOH:O	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:413:THR:HG22	1:C:465:TYR:CE1	2.55	0.42
1:C:415:LYS:NZ	9:C:833:HOH:O	2.52	0.42
1:C:394:MET:O	1:C:398:LYS:HG3	2.20	0.42
1:A:20:LYS:CE	9:A:837:HOH:O	2.67	0.41
1:A:323:ILE:HA	1:A:327:ILE:HB	2.01	0.41
1:C:394:MET:CA	1:C:394:MET:HE3	2.50	0.41
1:C:342:HIS:CE1	1:C:344:GLU:HB2	2.55	0.41
1:A:112:MET:CE	1:A:148:ASP:CB	2.99	0.41
1:C:6:ALA:CB	1:C:217:LEU:HD23	2.51	0.41
1:A:164:ARG:HG2	1:A:385:ARG:HH22	1.86	0.40
1:C:330:ILE:HG13	1:C:331:SER:H	1.85	0.40
1:C:402:ILE:HD12	1:C:402:ILE:HA	1.91	0.40
1:C:409:ASN:O	1:C:413:THR:HG23	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26:SER:OG	1:C:114:GLU:OE2[1_654]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	452/480 (94%)	443 (98%)	8 (2%)	1 (0%)	43	24
1	C	446/480 (93%)	431 (97%)	15 (3%)	0	100	100
All	All	898/960 (94%)	874 (97%)	23 (3%)	1 (0%)	48	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	326	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/422 (92%)	381 (98%)	7 (2%)	51	28
1	C	377/422 (89%)	375 (100%)	2 (0%)	81	70
All	All	765/844 (91%)	756 (99%)	9 (1%)	63	43

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	225	GLU
1	A	262	VAL
1	A	272	LEU
1	A	325	ILE
1	A	342	HIS
1	A	367	LEU
1	A	431	GLU
1	C	262	VAL
1	C	398	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	177	ASN
1	A	277	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic and 2 are modelled with single atom - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	ALA	A	505	4	5,5,5	1.39	2 (40%)	6,6,6	1.16	1 (16%)
2	SF4	C	502	1,6	0,12,12	-	-	-		
3	MET	A	503	2	7,8,8	1.07	1 (14%)	5,9,9	1.53	1 (20%)
2	SF4	A	502	1,6	0,12,12	-	-	-		
2	SF4	C	501	1,8	0,12,12	-	-	-		
2	SF4	A	501	1,3	0,12,12	-	-	-		
8	SAM	C	503	2	27,29,29	1.10	2 (7%)	34,42,42	2.42	11 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ALA	A	505	4	-	1/4/4/4	-
2	SF4	C	502	1,6	-	-	0/6/5/5
3	MET	A	503	2	-	5/8/8/8	-
2	SF4	A	502	1,6	-	-	0/6/5/5
2	SF4	C	501	1,8	-	-	0/6/5/5
2	SF4	A	501	1,3	-	-	0/6/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	SAM	C	503	2	-	3/17/33/33	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	C	503	SAM	C2-N1	2.52	1.38	1.33
8	C	503	SAM	OXT-C	-2.44	1.22	1.30
3	A	503	MET	OXT-C	-2.26	1.23	1.30
5	A	505	ALA	CA-C	-2.16	1.51	1.54
5	A	505	ALA	OXT-C	-2.07	1.24	1.30

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	C	503	SAM	N3-C2-N1	-8.08	116.34	128.58
8	C	503	SAM	C2-N3-C4	4.44	122.67	111.83
8	C	503	SAM	N9-C8-N7	-4.26	107.90	113.94
8	C	503	SAM	C4-N9-C8	3.95	109.89	105.74
8	C	503	SAM	C5-C4-N3	-3.88	121.38	126.72
3	A	503	MET	OXT-C-O	-3.20	116.83	124.08
8	C	503	SAM	C5-N7-C8	3.12	108.35	103.45
8	C	503	SAM	N3-C4-N9	2.85	132.01	127.17
8	C	503	SAM	C4-C5-N7	-2.81	107.37	110.58
8	C	503	SAM	OXT-C-O	-2.63	118.12	124.08
8	C	503	SAM	O4'-C1'-N9	2.55	112.99	108.09
8	C	503	SAM	C2-N1-C6	2.47	122.79	118.73
5	A	505	ALA	OXT-C-O	-2.20	119.10	124.08

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	503	MET	O-C-CA-N
8	C	503	SAM	N-CA-CB-CG
8	C	503	SAM	C-CA-CB-CG
3	A	503	MET	OXT-C-CA-N
3	A	503	MET	CB-CG-SD-CE
3	A	503	MET	OXT-C-CA-CB
3	A	503	MET	O-C-CA-CB
5	A	505	ALA	OXT-C-CA-CB
8	C	503	SAM	CB-CG-SD-C5'

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	505	ALA	2	0
3	A	503	MET	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	454/480 (94%)	0.46	20 (4%) 39 40	16, 34, 55, 73	1 (0%)
1	C	450/480 (93%)	0.26	21 (4%) 36 38	17, 29, 55, 75	0
All	All	904/960 (94%)	0.36	41 (4%) 38 40	16, 31, 55, 75	1 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	333	GLY	5.3
1	C	158	LEU	5.3
1	C	332	ALA	5.0
1	A	139	VAL	3.9
1	C	335	CYS	3.8
1	C	393	PHE	3.3
1	C	160	ASN	3.1
1	C	334	SER	3.0
1	A	153	ILE	2.9
1	C	280	ILE	2.9
1	C	395	SER	2.9
1	C	465	TYR	2.9
1	C	396	PHE	2.9
1	A	429	ILE	2.8
1	A	276	LEU	2.8
1	C	390	GLY	2.8
1	A	71	GLU	2.7
1	A	149	VAL	2.7
1	A	89	TYR	2.6
1	C	159	LYS	2.6
1	C	384	TYR	2.5
1	C	207	GLN	2.4
1	A	278	ILE	2.4
1	A	141	CYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	101	HIS	2.3
1	C	282	LYS	2.3
1	A	280	ILE	2.2
1	A	111	THR	2.2
1	A	352	LYS	2.2
1	C	402	ILE	2.1
1	C	336	THR	2.1
1	A	148	ASP	2.1
1	C	399	SER	2.1
1	A	355	PHE	2.1
1	C	154	TYR	2.1
1	A	344	GLU	2.1
1	C	400	GLY	2.1
1	A	86	VAL	2.0
1	A	207	GLN	2.0
1	A	384	TYR	2.0
1	A	341	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	NA	A	507	1/1	0.55	0.27	65,65,65,65	0
7	NA	C	506	1/1	0.84	0.16	53,53,53,53	0
5	ALA	A	505	6/6	0.87	0.11	23,27,33,34	10
3	MET	A	503	9/9	0.90	0.13	23,44,64,64	0
2	SF4	C	502	8/8	0.94	0.08	24,25,30,31	0
6	H2S	C	504	1/1	0.94	0.08	25,25,25,25	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SF4	A	501	8/8	0.95	0.07	26,27,30,31	0
8	SAM	C	503	27/27	0.95	0.06	16,21,35,43	4
6	H2S	A	506	1/1	0.97	0.05	25,25,25,25	0
7	NA	C	505	1/1	0.98	0.11	39,39,39,39	0
2	SF4	C	501	8/8	0.99	0.06	17,18,18,19	0
2	SF4	A	502	8/8	0.99	0.03	19,20,21,22	0
4	FE	A	504	1/1	1.00	0.04	24,24,24,24	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.