



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 12:11 PM UTC

PDB ID : 1WDK / pdb_00001wdk
Title : fatty acid beta-oxidation multienzyme complex from *Pseudomonas fragi*, form I (native2)
Authors : Ishikawa, M.; Tsuchiya, D.; Oyama, T.; Tsunaka, Y.; Morikawa, K.
Deposited on : 2004-05-17
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	715	% 72% 24% ..
1	B	715	2% 72% 25% ..
2	C	390	74% 24% .
2	D	390	% 72% 26% .

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 17701 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fatty oxidation complex alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	708	Total	C	N	O	S	0	0	0
			5324	3400	894	1003	27			
1	B	711	Total	C	N	O	S	0	0	0
			5333	3404	898	1004	27			

- Molecule 2 is a protein called 3-ketoacyl-CoA thiolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	390	Total	C	N	O	S	0	0	0
			2893	1801	515	548	29			
2	D	390	Total	C	N	O	S	0	0	0
			2889	1798	514	548	29			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 44	C 21	N 7	O 14	P 2	0	0
4	B	1	Total 44	C 21	N 7	O 14	P 2	0	0

- Molecule 5 is 3,6,9,12,15-PENTAOXATRICOSAN-1-OL (CCD ID: N8E) (formula: $C_{18}H_{38}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			24	18	6		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			24	18	6		
5	B	1	Total	C	O	0	0
			24	18	6		
5	B	1	Total	C	O	0	0
			24	18	6		

- Molecule 6 is MERCURY (II) ION (CCD ID: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	1	Total	Hg	0	0
			1	1		
6	D	1	Total	Hg	0	0
			1	1		


- Molecule 7 is ACETYL COENZYME *A (CCD ID: ACO) (formula: C₂₃H₃₈N₇O₁₇P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
7	C	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		
7	D	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	286	Total 286	O 286	0	0
8	B	309	Total 309	O 309	0	0
8	C	205	Total 205	O 205	0	0
8	D	173	Total 173	O 173	0	0




• Molecule 2: 3-ketoacyl-CoA thiolase

Chain C: 74% 24%



• Molecule 2: 3-ketoacyl-CoA thiolase

Chain D: 72% 26%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	180.71Å 94.58Å 160.00Å 90.00° 111.45° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 20.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.50) 95.1 (20.00-2.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.48 (at 2.51Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.203 , 0.244 0.202 , 0.244	Depositor DCC
R_{free} test set	5875 reflections (7.09%)	wwPDB-VP
Wilson B-factor (Å ²)	28.5	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	17701	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAD, ACO, N8E, HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.41	0/5410	0.92	16/7310 (0.2%)
1	B	0.42	0/5419	0.92	20/7325 (0.3%)
2	C	0.49	2/2941 (0.1%)	0.97	6/3967 (0.2%)
2	D	0.48	1/2937 (0.0%)	0.94	5/3963 (0.1%)
All	All	0.44	3/16707 (0.0%)	0.93	47/22565 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	255	MET	SD-CE	-7.59	1.60	1.79
2	C	151	MET	SD-CE	-6.16	1.64	1.79
2	D	255	MET	SD-CE	-5.12	1.66	1.79

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	249	TYR	CA-C-N	8.33	127.63	118.97
1	A	249	TYR	C-N-CA	8.33	127.63	118.97
2	C	93	ARG	N-CA-C	-8.19	100.74	110.41
2	C	352	SER	N-CA-C	8.08	120.80	111.11
1	B	249	TYR	CA-C-N	8.00	127.94	119.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5324	0	5391	165	0
1	B	5333	0	5385	159	0
2	C	2893	0	2903	112	0
2	D	2889	0	2892	118	0
3	A	1	0	0	0	0
4	A	44	0	26	2	0
4	B	44	0	26	2	0
5	A	48	0	76	8	0
5	B	48	0	76	8	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	C	51	0	34	3	0
7	D	51	0	34	4	0
8	A	286	0	0	8	0
8	B	309	0	0	13	0
8	C	205	0	0	4	0
8	D	173	0	0	9	0
All	All	17701	0	16843	540	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 540 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:525:ASP:HB2	1:B:536:GLY:HA3	1.23	1.17
1:A:525:ASP:HB2	1:A:536:GLY:HA3	1.21	1.13
1:A:695:ALA:HA	1:A:698:HIS:HD2	1.19	1.04
1:A:695:ALA:HA	1:A:698:HIS:CD2	1.93	1.02
1:B:595:ASP:HA	1:B:600:GLN:HA	1.41	1.00

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	704/715 (98%)	668 (95%)	35 (5%)	1 (0%)	48	68
1	B	707/715 (99%)	672 (95%)	34 (5%)	1 (0%)	48	68
2	C	388/390 (100%)	370 (95%)	17 (4%)	1 (0%)	36	55
2	D	388/390 (100%)	367 (95%)	20 (5%)	1 (0%)	36	55
All	All	2187/2210 (99%)	2077 (95%)	106 (5%)	4 (0%)	43	63

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	324	ILE
1	B	324	ILE
2	C	349	PHE
2	D	349	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	547/562 (97%)	522 (95%)	25 (5%)	24	48
1	B	545/562 (97%)	519 (95%)	26 (5%)	23	46
2	C	307/307 (100%)	297 (97%)	10 (3%)	33	61
2	D	306/307 (100%)	297 (97%)	9 (3%)	37	65
All	All	1705/1738 (98%)	1635 (96%)	70 (4%)	27	53

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	318	GLN
2	C	351	CYS
2	D	289	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	703	LEU
1	A	686	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 40 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	162	HIS
2	D	107	GLN
2	C	169	GLN
2	D	4	ASN
2	D	169	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 3 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	N8E	A	1002	-	23,23,23	0.46	0	22,22,22	0.44	0
5	N8E	B	2003	-	23,23,23	0.49	0	22,22,22	0.41	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	N8E	A	1003	-	23,23,23	0.46	0	22,22,22	0.44	0
7	ACO	C	3001	6	51,53,53	0.75	0	73,79,79	0.75	1 (1%)
4	NAD	A	1001	-	46,48,48	1.56	6 (13%)	64,73,73	1.34	4 (6%)
7	ACO	D	4001	6	51,53,53	0.69	0	73,79,79	0.67	1 (1%)
4	NAD	B	2001	-	46,48,48	1.59	6 (13%)	64,73,73	1.34	4 (6%)
5	N8E	B	2002	-	23,23,23	0.46	0	22,22,22	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	N8E	A	1002	-	-	15/21/21/21	-
5	N8E	B	2003	-	-	13/21/21/21	-
5	N8E	A	1003	-	-	15/21/21/21	-
7	ACO	C	3001	6	-	8/51/67/67	0/3/3/3
4	NAD	A	1001	-	-	12/30/62/62	0/5/5/5
7	ACO	D	4001	6	-	12/51/67/67	0/3/3/3
4	NAD	B	2001	-	-	11/30/62/62	0/5/5/5
5	N8E	B	2002	-	-	13/21/21/21	-

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1001	NAD	C3N-C7N	5.91	1.59	1.50
4	B	2001	NAD	C3N-C7N	5.58	1.58	1.50
4	B	2001	NAD	C2N-N1N	4.79	1.40	1.35
4	A	1001	NAD	C2N-N1N	4.26	1.39	1.35
4	A	1001	NAD	C4N-C3N	3.80	1.45	1.39

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1001	NAD	C5N-C4N-C3N	-6.77	113.70	120.36
4	B	2001	NAD	C5N-C4N-C3N	-6.62	113.86	120.36
4	A	1001	NAD	C6N-C5N-C4N	5.58	127.49	119.45
4	B	2001	NAD	C6N-C5N-C4N	5.53	127.42	119.45
7	C	3001	ACO	C2P-S1P-C	4.58	123.44	101.42

There are no chirality outliers.

5 of 99 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1001	NAD	O4D-C1D-N1N-C2N
4	A	1001	NAD	C2N-C3N-C7N-O7N
4	A	1001	NAD	C2N-C3N-C7N-N7N
4	B	2001	NAD	O4D-C1D-N1N-C2N
4	B	2001	NAD	C2N-C3N-C7N-O7N




There are no ring outliers.


8 monomers are involved in 27 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1002	N8E	2	0
5	B	2003	N8E	7	0
5	A	1003	N8E	6	0
7	C	3001	ACO	3	0
4	A	1001	NAD	2	0
7	D	4001	ACO	4	0
4	B	2001	NAD	2	0
5	B	2002	N8E	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	708/715 (99%)	-0.08	8 (1%) 78 75	14, 29, 51, 72	0
1	B	711/715 (99%)	-0.07	15 (2%) 63 59	13, 29, 52, 71	0
2	C	390/390 (100%)	-0.44	1 (0%) 90 87	11, 21, 35, 52	0
2	D	390/390 (100%)	-0.43	3 (0%) 82 80	10, 21, 35, 52	0
All	All	2199/2210 (99%)	-0.20	27 (1%) 76 73	10, 26, 48, 72	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	2	SER	4.8
1	B	594	ALA	4.3
1	B	567	ASP	3.9
1	B	294	PHE	3.5
1	B	620	ARG	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	N8E	A	1003	24/24	0.71	0.22	62,83,88,89	0
5	N8E	B	2003	24/24	0.74	0.19	29,61,81,81	0
7	ACO	C	3001	51/51	0.77	0.20	69,102,115,116	0
5	N8E	B	2002	24/24	0.79	0.16	42,55,59,60	0
7	ACO	D	4001	51/51	0.80	0.20	72,90,97,98	0
5	N8E	A	1002	24/24	0.81	0.18	50,67,70,71	0
4	NAD	B	2001	44/44	0.90	0.11	41,51,73,73	0
4	NAD	A	1001	44/44	0.92	0.10	46,50,60,61	0
3	ZN	A	716	1/1	0.97	0.06	43,43,43,43	0
6	HG	C	392	1/1	0.98	0.08	87,87,87,87	0
6	HG	D	1	1/1	0.99	0.05	75,75,75,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around N8E B 2003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ACO C 3001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around N8E B 2002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ACO D 4001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around N8E A 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAD B 2001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NAD A 1001:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.