



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:27 AM UTC

PDB ID : 1WDR / pdb_00001wdr
Title : The role of an inner loop in the catalytic mechanism of soybean beta-amylase
Authors : Kang, Y.N.; Adachi, M.; Utsumi, S.; Mikami, B.
Deposited on : 2004-05-17
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1232 (1.36-1.36)
Ramachandran outliers	187476	1220 (1.36-1.36)
Sidechain outliers	187428	1220 (1.36-1.36)
RSRZ outliers	180081	1214 (1.36-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	495	87% 11% .
2	B	4	50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	B	3[A]	X	-	-	-
2	GLC	B	3[B]	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5007 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-amylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	493	4012	2573	671	750	18	0	20	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	76	LEU	PHE	SEE REMARK 999	UNP P10538
A	202	GLY	ARG	SEE REMARK 999	UNP P10538
A	342	SER	THR	engineered mutation	UNP P10538
A	399	ARG	LYS	SEE REMARK 999	UNP P10538

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	4	47	24	23	0	3	0

- Molecule 3 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	1
			12	6	6		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	906	Total	O	0	0
			906	906		

- Molecule 1: Beta-amylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	85.13Å 85.13Å 143.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 1.35 10.00 – 1.35	Depositor EDS
% Data completeness (in resolution range)	94.3 (10.00-1.35) 94.7 (10.00-1.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.71 (at 1.35Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.114 , 0.172 0.125 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	10.0	Xtriage
Anisotropy	0.536	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 70.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5007	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, CME, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	2/4187 (0.0%)	1.46	57/5684 (1.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	199	GLU	N-CA	-6.36	1.38	1.46
1	A	334	ARG	CD-NE	-6.03	1.37	1.46

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	239	ASN	OD1-CG-ND2	-12.63	109.97	122.60
1	A	87	GLN	OE1-CD-NE2	12.26	134.86	122.60
1	A	198	TRP	CA-C-N	10.27	139.58	122.33
1	A	198	TRP	C-N-CA	10.27	139.58	122.33
1	A	216	ASP	CA-CB-CG	9.62	122.22	112.60
1	A	11	TYR	CA-CB-CG	9.49	130.98	113.90
1	A	279	ASP	CA-CB-CG	9.39	121.99	112.60
1	A	347	ARG	CD-NE-CZ	8.88	136.83	124.40
1	A	239	ASN	CB-CG-OD1	8.28	137.36	120.80
1	A	282[A]	ASN	OD1-CG-ND2	8.16	130.76	122.60
1	A	282[B]	ASN	OD1-CG-ND2	8.16	130.76	122.60
1	A	334	ARG	CD-NE-CZ	8.13	135.79	124.40
1	A	340	ASN	CA-CB-CG	7.92	120.52	112.60
1	A	351	GLN	OE1-CD-NE2	7.90	130.50	122.60
1	A	410	LYS	N-CA-CB	7.65	121.37	110.12
1	A	87	GLN	CG-CD-NE2	-7.57	105.05	116.40
1	A	151	ILE	O-C-N	-7.29	114.80	121.87
1	A	493	VAL	O-C-N	7.16	129.10	121.87
1	A	268	ASN	CA-CB-CG	6.95	119.55	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	464	SER	CA-CB-OG	-6.92	97.27	111.10
1	A	458	ILE	N-CA-CB	6.58	120.50	111.41
1	A	300	HIS	ND1-CE1-NE2	-6.52	101.88	108.40
1	A	247	PHE	O-C-N	-6.52	114.72	122.15
1	A	417	THR	CA-CB-OG1	-6.38	100.04	109.60
1	A	347	ARG	NE-CZ-NH2	6.37	124.94	119.20
1	A	5	SER	O-C-N	-6.25	114.58	122.27
1	A	273	HIS	CA-CB-CG	6.11	119.91	113.80
1	A	151	ILE	CA-C-N	6.05	128.39	120.28
1	A	151	ILE	C-N-CA	6.05	128.39	120.28
1	A	347	ARG	NH1-CZ-NH2	-6.02	111.48	119.30
1	A	3	SER	CA-C-N	6.00	132.99	121.54
1	A	3	SER	C-N-CA	6.00	132.99	121.54
1	A	97	GLY	O-C-N	-5.93	116.34	121.97
1	A	282[A]	ASN	CA-CB-CG	-5.91	106.69	112.60
1	A	282[B]	ASN	CA-CB-CG	-5.91	106.69	112.60
1	A	30	GLU	CA-C-N	-5.83	116.43	122.85
1	A	30	GLU	C-N-CA	-5.83	116.43	122.85
1	A	334	ARG	NE-CZ-NH1	5.83	127.33	121.50
1	A	336	HIS	CA-CB-CG	-5.81	107.99	113.80
1	A	176	ASP	CA-CB-CG	5.68	118.28	112.60
1	A	254	TYR	CA-CB-CG	5.68	124.12	113.90
1	A	118	ASN	OD1-CG-ND2	-5.66	116.94	122.60
1	A	247	PHE	CA-C-N	5.61	131.19	122.26
1	A	247	PHE	C-N-CA	5.61	131.19	122.26
1	A	167	ASP	CA-CB-CG	5.48	118.08	112.60
1	A	198	TRP	O-C-N	-5.41	116.15	122.81
1	A	266	TYR	CA-CB-CG	5.37	123.57	113.90
1	A	419	LEU	CA-C-N	5.35	131.33	121.70
1	A	419	LEU	C-N-CA	5.35	131.33	121.70
1	A	493	VAL	CA-C-O	-5.32	114.88	120.57
1	A	239	ASN	N-CA-C	5.31	119.45	112.92
1	A	429	SER	O-C-N	5.12	127.62	122.09
1	A	423	ASP	CA-CB-CG	5.10	117.70	112.60
1	A	376	ARG	CD-NE-CZ	5.06	131.49	124.40
1	A	300	HIS	ND1-CG-CD2	-5.04	101.06	106.10
1	A	20	LEU	CA-C-O	5.02	126.71	121.19
1	A	97	GLY	CA-C-O	5.01	123.60	119.04

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4012	0	3907	30	0
2	B	47	0	26	3	0
3	A	12	0	4	4	0
4	A	30	0	0	0	0
5	A	906	0	0	17	0
All	All	5007	0	3937	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:2[A]:GLC:O4	2:B:3[A]:GLC:H1	0.83	1.00
3:A:499[B]:GLC:C5	3:A:499[B]:GLC:O5	2.11	0.98
3:A:499[B]:GLC:C5	3:A:499[B]:GLC:C6	2.42	0.98
3:A:499[B]:GLC:C5	3:A:499[B]:GLC:C1	2.48	0.90
2:B:2[A]:GLC:HO4	2:B:3[A]:GLC:H1	1.35	0.90
2:B:2[A]:GLC:C4	2:B:3[A]:GLC:H1	2.08	0.84
1:A:458:ILE:HD12	5:A:1097:HOH:O	1.82	0.80
1:A:468:ILE:HD11	5:A:1301:HOH:O	1.84	0.77
1:A:30:GLU:HB2	5:A:1257:HOH:O	1.86	0.74
1:A:424:ASP:O	1:A:427[B]:GLN:HG3	1.92	0.69
1:A:116:GLU:HG2	5:A:876:HOH:O	1.93	0.68
1:A:3:SER:O	1:A:7:MET:HG2	1.96	0.66
1:A:376:ARG:HD2	5:A:982:HOH:O	1.96	0.65
1:A:471:GLU:HG3	5:A:1276:HOH:O	1.96	0.65
1:A:288:CME:HZ2	5:A:1230:HOH:O	1.96	0.64
1:A:351:GLN:HG3	1:A:357[A]:SER:OG	1.96	0.64
1:A:123:TYR:OH	1:A:137[B]:VAL:HG23	2.00	0.61
1:A:493:VAL:O	1:A:494:ASP:HB2	1.99	0.61
1:A:25[B]:VAL:HG13	5:A:1042:HOH:O	2.00	0.61
1:A:456:HIS:HD2	5:A:566:HOH:O	1.86	0.58
1:A:467:LYS:HE2	5:A:792:HOH:O	2.05	0.56
1:A:364[B]:GLN:HG2	1:A:485:TRP:CD2	2.42	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:401:GLN:HG2	5:A:1223:HOH:O	2.06	0.55
3:A:499[B]:GLC:O5	3:A:499[B]:GLC:C4	2.55	0.54
1:A:25[A]:VAL:HG22	5:A:1042:HOH:O	2.07	0.54
1:A:351:GLN:HG3	1:A:357[A]:SER:HG	1.74	0.53
1:A:282[A]:ASN:ND2	5:A:1231:HOH:O	2.46	0.47
1:A:7:MET:SD	1:A:403[B]:VAL:HG11	2.55	0.46
1:A:214[A]:LYS:NZ	5:A:1192:HOH:O	2.49	0.46
1:A:401:GLN:HG2	5:A:1167:HOH:O	2.18	0.43
1:A:403[A]:VAL:HG22	1:A:490:ASP:OD2	2.18	0.42
1:A:126[B]:ARG:NH1	5:A:1090:HOH:O	2.52	0.42
1:A:400:PRO:HB3	1:A:485:TRP:CE2	2.56	0.41
1:A:451:PRO:HA	1:A:454:TYR:CE2	2.57	0.40
1:A:7:MET:HE3	1:A:443:ALA:O	2.21	0.40
1:A:4:ASP:HB2	5:A:1342:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	510/495 (103%)	499 (98%)	9 (2%)	2 (0%)	30	11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	494	ASP
1	A	4	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	442/423 (104%)	441 (100%)	1 (0%)	87	74

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	471	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	87	GLN
1	A	118	ASN
1	A	164	ASN
1	A	196	GLN
1	A	207	GLN
1	A	239	ASN
1	A	268	ASN
1	A	276	GLN
1	A	340	ASN
1	A	393	GLN
1	A	401	GLN
1	A	456	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	CME	A	288	1	8,9,10	0.85	0	6,9,11	2.10	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	288	1	-	1/5/8/10	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	288	CME	CZ-CE-SD	-3.65	101.19	113.39
1	A	288	CME	OH-CZ-CE	3.17	123.18	110.82

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	288	CME	SD-CE-CZ-OH

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	288	CME	1	0

5.5 Carbohydrates [i](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	B	1[A]	2	12,12,12	0.79	0	17,17,17	1.86	6 (35%)
2	GLC	B	2[A]	2	11,11,12	1.26	1 (9%)	15,15,17	2.39	5 (33%)
2	GLC	B	3[A]	-	12,12,12	0.91	0	17,17,17	1.72	4 (23%)
2	GLC	B	3[B]	-	12,12,12	0.94	0	17,17,17	1.77	4 (23%)
2	GLC	B	4	2	11,11,12	1.11	0	15,15,17	1.49	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1[A]	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2[A]	2	-	1/2/19/22	0/1/1/1
2	GLC	B	3[A]	-	1/1/5/5	0/2/22/22	0/1/1/1
2	GLC	B	3[B]	-	1/1/5/5	0/2/22/22	0/1/1/1
2	GLC	B	4	2	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2[A]	GLC	C4-C5	3.15	1.59	1.53

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2[A]	GLC	O5-C5-C6	-4.79	98.35	107.66
2	B	1[A]	GLC	C4-C3-C2	-4.51	102.92	110.83
2	B	2[A]	GLC	O5-C5-C4	-4.33	100.29	110.83
2	B	2[A]	GLC	C6-C5-C4	-4.23	102.64	113.02
2	B	3[B]	GLC	O1-C1-O5	-3.45	100.17	110.41
2	B	2[A]	GLC	C3-C4-C5	-3.31	104.22	110.23
2	B	3[A]	GLC	O1-C1-O5	-3.09	101.25	110.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3[A]	GLC	O2-C2-C1	-3.07	102.15	109.25
2	B	3[B]	GLC	O2-C2-C1	-3.07	102.15	109.25
2	B	4	GLC	C2-C3-C4	-2.91	105.74	110.86
2	B	4	GLC	C1-O5-C5	2.77	115.91	112.19
2	B	1[A]	GLC	O1-C1-C2	2.71	116.86	108.98
2	B	1[A]	GLC	O3-C3-C2	-2.60	104.25	110.38
2	B	3[A]	GLC	O5-C5-C4	-2.60	105.02	109.70
2	B	3[B]	GLC	O5-C5-C4	-2.60	105.02	109.70
2	B	1[A]	GLC	O5-C1-C2	2.57	114.81	110.30
2	B	1[A]	GLC	O5-C5-C4	-2.37	105.42	109.70
2	B	3[A]	GLC	O4-C4-C5	-2.19	103.92	109.32
2	B	3[B]	GLC	O4-C4-C5	-2.19	103.92	109.32
2	B	4	GLC	O4-C4-C3	-2.15	105.31	110.38
2	B	2[A]	GLC	O2-C2-C3	-2.15	105.71	110.15
2	B	4	GLC	O5-C1-C2	-2.12	105.72	110.79
2	B	1[A]	GLC	C1-C2-C3	-2.10	106.06	110.36

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	3[A]	GLC	C1
2	B	3[B]	GLC	C1

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2[A]	GLC	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2[A]	GLC	3	0
2	B	3[A]	GLC	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	2003	-	4,4,4	0.42	0	6,6,6	0.70	0
4	SO4	A	2001	-	4,4,4	0.60	0	6,6,6	0.23	0
4	SO4	A	2000	-	4,4,4	0.48	0	6,6,6	0.61	0
4	SO4	A	2005	-	4,4,4	0.50	0	6,6,6	0.64	0
4	SO4	A	2004	-	4,4,4	0.42	0	6,6,6	0.34	0
4	SO4	A	2002	-	4,4,4	0.42	0	6,6,6	0.53	0
3	GLC	A	499[B]	-	12,12,12	11.15	2 (16%)	17,17,17	6.43	9 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	A	499[B]	-	-	1/2/22/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	499[B]	GLC	O5-C5	27.33	2.11	1.44
3	A	499[B]	GLC	C6-C5	27.15	2.42	1.51

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	499[B]	GLC	C1-O5-C5	-13.40	87.73	113.65
3	A	499[B]	GLC	O5-C5-C6	-13.29	73.51	106.44
3	A	499[B]	GLC	O5-C5-C4	-11.99	88.10	109.70
3	A	499[B]	GLC	C3-C4-C5	8.32	125.32	110.23
3	A	499[B]	GLC	O6-C6-C5	-7.28	86.53	111.33
3	A	499[B]	GLC	O5-C1-C2	6.11	121.05	110.30
3	A	499[B]	GLC	C6-C5-C4	-4.24	102.61	113.02
3	A	499[B]	GLC	O1-C1-O5	3.25	120.07	110.41
3	A	499[B]	GLC	O3-C3-C2	-2.18	105.24	110.38

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	499[B]	GLC	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	499[B]	GLC	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	492/495 (99%)	-0.67	5 (1%) 79 88	6, 11, 25, 96	20 (4%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	SER	6.0
1	A	495	GLY	5.2
1	A	6	ASN	2.7
1	A	494	ASP	2.4
1	A	4	ASP	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	288	10/11	0.97	0.07	11,15,44,49	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	B	2[A]	11/12	0.90	0.11	13,19,22,24	11

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	B	1[A]	12/12	0.95	0.06	16,20,22,24	12
2	GLC	B	3[A]	12/12	0.98	0.04	12,14,17,19	1
2	GLC	B	3[B]	12/12	0.98	0.04	12,14,17,20	1
2	GLC	B	4	11/12	0.99	0.02	10,11,12,13	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	A	2002	5/5	0.83	0.17	78,81,96,122	0
3	GLC	A	499[B]	12/12	0.87	0.09	16,21,24,25	12
4	SO4	A	2000	5/5	0.89	0.11	52,55,67,69	0
4	SO4	A	2001	5/5	0.91	0.09	37,39,57,63	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	A	2005	5/5	0.97	0.08	26,31,36,42	0
4	SO4	A	2003	5/5	0.98	0.07	26,27,42,46	0
4	SO4	A	2004	5/5	0.99	0.06	14,16,17,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.