



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:56 PM UTC

PDB ID : 1WDU / pdb_00001wdu
Title : Endonuclease domain of TRAS1, a telomere-specific non-LTR retrotransposon
Authors : Maita, N.; Anzai, T.; Aoyagi, H.; Mizuno, H.; Fujiwara, H.
Deposited on : 2004-05-17
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	
1	B	245	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3504 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRAS1 ORF2p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1707	1082	294	326	5			
1	B	219	Total	C	N	O	S	0	0	0
			1735	1097	305	328	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	167	SER	CYS	engineered mutation	UNP Q93139
B	167	SER	CYS	engineered mutation	UNP Q93139

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Cl 1	0	0
3	B	1	Total 1	Cl 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	20	Total 20	O 20	0	0
4	B	35	Total 35	O 35	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRAS1 ORF2p



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	64.64Å 64.64Å 117.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.40 20.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-2.40) 99.5 (20.00-2.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.96 (at 2.41Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.225 , 0.253 0.224 , 0.251	Depositor DCC
R_{free} test set	1681 reflections (7.88%)	wwPDB-VP
Wilson B-factor (Å ²)	37.3	Xtriage
Anisotropy	0.469	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.005 for -h,-k,l 0.033 for h,-h-k,-l 0.022 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3504	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1738	0.92	5/2353 (0.2%)
1	B	0.42	0/1766	0.91	4/2388 (0.2%)
All	All	0.42	0/3504	0.91	9/4741 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	54	ILE	N-CA-C	6.86	118.94	108.71
1	A	220	LEU	N-CA-C	-6.17	101.28	110.23
1	A	196	THR	N-CA-C	-5.79	106.21	113.28
1	A	38	ALA	N-CA-C	-5.38	105.33	111.14
1	B	100	TYR	CA-C-N	5.36	125.03	119.56
1	B	100	TYR	C-N-CA	5.36	125.03	119.56
1	A	54	ILE	N-CA-C	5.36	117.27	108.97
1	B	246	LEU	N-CA-C	-5.20	108.69	114.62
1	A	245	ARG	CB-CA-C	-5.12	109.00	116.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1707	0	1679	68	0
1	B	1735	0	1722	63	0
2	A	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	20	0	0	1	0
4	B	35	0	0	1	0
All	All	3504	0	3401	131	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (131) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:ARG:HH11	1:A:72:ARG:HG3	1.18	1.06
1:A:117:ALA:HB1	1:A:246:LEU:HD21	1.41	0.97
1:B:102:GLN:NE2	1:B:102:GLN:H	1.66	0.93
1:B:201:ARG:HA	1:B:201:ARG:HE	1.31	0.93
1:A:35:LEU:H	1:A:35:LEU:HD23	1.34	0.91
1:B:79:GLN:HB3	1:B:83:THR:HB	1.57	0.87
1:B:31:GLN:HE21	1:B:235:SER:HB2	1.42	0.84
1:A:72:ARG:HG3	1:A:72:ARG:NH1	1.95	0.81
1:B:31:GLN:NE2	1:B:235:SER:HB2	1.98	0.79
1:A:220:LEU:O	1:A:221:ASP:HB2	1.84	0.76
1:B:102:GLN:H	1:B:102:GLN:HE21	1.33	0.75
1:A:153:ILE:HD11	1:A:219:MET:HE2	1.69	0.75
1:B:76:SER:HB2	1:B:105:THR:O	1.86	0.74
1:A:229:ASP:HB2	1:A:232:LEU:HB2	1.69	0.73
1:A:152:LEU:HB2	1:A:216:THR:HG23	1.72	0.70
1:A:128:GLU:HG3	1:A:131:LYS:HG3	1.73	0.69
1:B:102:GLN:HE21	1:B:102:GLN:N	1.91	0.69
1:A:104:THR:HG22	1:A:110:VAL:H	1.60	0.67
1:B:232:LEU:HD23	1:B:239:GLY:HA3	1.76	0.67
1:A:229:ASP:CB	1:A:232:LEU:HB2	2.24	0.67
1:B:229:ASP:O	1:B:230:GLU:HB2	1.93	0.66
1:B:201:ARG:HA	1:B:201:ARG:NE	2.08	0.64
1:B:145:ARG:HH21	1:B:146:LYS:HE2	1.63	0.63
1:A:117:ALA:HB1	1:A:246:LEU:CD2	2.25	0.62
1:B:159:ASN:H	1:B:210:VAL:HG13	1.65	0.61
1:B:79:GLN:CB	1:B:83:THR:HB	2.29	0.61
1:A:232:LEU:HB3	1:A:239:GLY:HA3	1.81	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:217:GLU:HA	1:B:220:LEU:HD23	1.82	0.60
1:B:152:LEU:HB2	1:B:216:THR:HG23	1.82	0.60
1:A:189:LEU:HD12	1:A:213:THR:HG22	1.84	0.59
1:A:21:PRO:N	1:A:22:PRO:CD	2.65	0.59
1:A:33:LYS:NZ	1:A:235:SER:HA	2.16	0.59
1:A:34:LYS:HE2	1:A:58:TYR:CE2	2.37	0.59
1:A:227:ARG:HG3	1:A:241:VAL:HG22	1.84	0.58
1:B:35:LEU:O	1:B:39:GLU:HG3	2.03	0.58
1:B:151:ARG:HG2	1:B:151:ARG:HH11	1.67	0.58
1:B:226:TRP:HA	1:B:241:VAL:O	2.03	0.57
1:A:220:LEU:O	1:A:221:ASP:CB	2.52	0.57
1:A:198:ASP:OD1	1:A:207:GLN:HG2	2.04	0.57
1:A:151:ARG:HH12	1:A:222:LEU:HD21	1.69	0.56
1:A:76:SER:HB2	1:A:105:THR:O	2.05	0.56
1:A:235:SER:C	1:A:237:HIS:H	2.13	0.56
1:A:35:LEU:H	1:A:35:LEU:CD2	2.11	0.56
1:B:128:GLU:HB2	1:B:131:LYS:HB2	1.86	0.56
1:B:30:LEU:HD21	1:B:40:LEU:HD22	1.88	0.56
1:A:79:GLN:HG2	1:A:83:THR:O	2.05	0.55
1:B:230:GLU:HG2	1:B:238:ASN:CG	2.31	0.55
1:B:178:MET:SD	1:B:188:ILE:HD11	2.48	0.54
1:B:145:ARG:NH2	1:B:146:LYS:HE2	2.22	0.54
1:A:121:THR:HG21	1:A:147:MET:HG2	1.89	0.54
1:B:51:ILE:HG22	1:B:52:ALA:N	2.24	0.53
1:B:217:GLU:O	1:B:220:LEU:HD23	2.08	0.53
1:B:107:ASN:ND2	1:B:136:TYR:HE2	2.07	0.52
1:B:120:ILE:HD12	1:B:246:LEU:HD11	1.90	0.52
1:A:33:LYS:HD2	1:A:235:SER:HB3	1.91	0.52
1:A:123:VAL:HG13	1:A:154:PHE:HA	1.91	0.52
1:A:57:PRO:HD2	1:A:84:VAL:O	2.09	0.52
1:A:178:MET:SD	1:A:188:ILE:HD11	2.49	0.52
1:B:229:ASP:O	1:B:230:GLU:CB	2.57	0.52
1:B:151:ARG:HG2	1:B:151:ARG:NH1	2.25	0.52
1:B:166:GLY:HA3	1:B:209:ARG:HD2	1.93	0.51
1:A:54:ILE:HG22	1:A:55:GLN:N	2.26	0.50
1:B:121:THR:HB	1:B:152:LEU:HD23	1.93	0.50
1:B:159:ASN:N	1:B:210:VAL:HG13	2.26	0.50
1:A:33:LYS:HD2	1:A:235:SER:CB	2.41	0.50
1:A:68:PHE:HB3	1:A:71:VAL:HB	1.94	0.50
1:B:52:ALA:HB3	1:B:89:ALA:HB3	1.93	0.49
1:B:209:ARG:NH2	4:B:264:HOH:O	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:ALA:O	1:B:51:ILE:HD13	2.13	0.48
1:A:145:ARG:HG3	1:A:146:LYS:N	2.28	0.48
1:A:229:ASP:O	1:A:231:ASP:N	2.47	0.48
1:A:51:ILE:HG22	1:A:52:ALA:N	2.28	0.48
1:A:30:LEU:HD22	1:A:54:ILE:CG2	2.44	0.48
1:A:232:LEU:HD23	1:A:239:GLY:HA3	1.96	0.48
1:A:190:ASN:O	1:A:191:GLU:HG3	2.14	0.48
1:A:22:PRO:HB3	1:A:245:ARG:NH1	2.29	0.47
1:B:133:ILE:O	1:B:137:LEU:HG	2.15	0.47
1:B:114:ARG:HB3	1:B:119:GLU:HG3	1.97	0.47
1:B:69:ARG:C	1:B:71:VAL:H	2.21	0.47
1:A:33:LYS:HZ1	1:A:235:SER:HA	1.77	0.46
1:A:164:TRP:CD1	1:A:178:MET:HE2	2.51	0.46
1:A:123:VAL:HG11	1:A:154:PHE:CD2	2.51	0.45
1:A:229:ASP:C	1:A:231:ASP:N	2.74	0.45
1:B:109:VAL:O	1:B:123:VAL:HA	2.16	0.45
1:A:47:ARG:NH2	1:A:229:ASP:OD2	2.50	0.45
1:A:164:TRP:HB3	1:A:188:ILE:CD1	2.47	0.45
1:A:229:ASP:O	1:A:230:GLU:C	2.60	0.45
1:B:54:ILE:HG21	1:B:57:PRO:HB3	1.99	0.45
1:B:230:GLU:HG2	1:B:238:ASN:ND2	2.32	0.45
1:A:100:TYR:O	1:A:104:THR:HG23	2.17	0.45
1:B:210:VAL:HG13	1:B:210:VAL:O	2.17	0.44
1:A:223:ILE:HA	1:A:243:ASN:O	2.16	0.44
1:B:101:PRO:HD2	1:B:102:GLN:HE22	1.83	0.44
1:B:230:GLU:HA	1:B:238:ASN:OD1	2.18	0.44
1:A:43:GLU:OE2	1:A:232:LEU:HD12	2.17	0.43
1:A:246:LEU:O	1:A:247:GLN:C	2.60	0.43
1:B:120:ILE:HD13	1:B:219:MET:HE1	2.00	0.43
1:A:30:LEU:HD22	1:A:54:ILE:HG23	2.00	0.43
1:A:52:ALA:HB2	1:A:91:PHE:HE1	1.84	0.43
1:A:113:ILE:HD13	1:A:122:LEU:HD11	2.01	0.43
1:A:231:ASP:C	1:A:233:VAL:H	2.26	0.43
1:B:234:SER:O	1:B:235:SER:C	2.61	0.43
1:B:79:GLN:NE2	1:B:79:GLN:HA	2.33	0.42
1:A:143:VAL:O	1:A:147:MET:HB2	2.19	0.42
1:B:51:ILE:CG2	1:B:52:ALA:N	2.82	0.42
1:B:115:THR:C	1:B:117:ALA:H	2.27	0.42
1:A:195:PRO:O	1:A:238:ASN:ND2	2.51	0.42
1:A:104:THR:HG21	4:A:258:HOH:O	2.20	0.42
1:B:197:PHE:O	1:B:207:GLN:HA	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:124:SER:HA	1:B:155:GLY:O	2.19	0.42
1:A:33:LYS:O	1:A:37:THR:HG23	2.19	0.41
1:A:143:VAL:O	1:A:147:MET:CB	2.68	0.41
1:A:231:ASP:C	1:A:233:VAL:N	2.75	0.41
1:A:158:ALA:O	1:A:159:ASN:C	2.63	0.41
1:B:120:ILE:CD1	1:B:246:LEU:HD11	2.49	0.41
1:B:217:GLU:HA	1:B:220:LEU:CD2	2.48	0.41
1:A:177:LEU:O	1:A:181:LEU:HD13	2.20	0.41
1:A:235:SER:C	1:A:237:HIS:N	2.78	0.41
1:B:26:LEU:O	1:B:52:ALA:HA	2.20	0.41
1:A:127:PHE:CE2	1:A:136:TYR:HB3	2.55	0.41
1:B:127:PHE:HZ	1:B:140:ILE:HD11	1.85	0.41
1:B:76:SER:HB3	1:B:109:VAL:HG22	2.01	0.41
1:B:201:ARG:NE	1:B:201:ARG:CA	2.79	0.41
1:A:197:PHE:HA	1:A:236:ASP:O	2.21	0.41
1:B:102:GLN:NE2	1:B:102:GLN:N	2.44	0.41
1:B:202:GLY:C	1:B:204:LYS:H	2.29	0.41
1:B:159:ASN:HA	1:B:210:VAL:CG1	2.51	0.41
1:A:184:LEU:HB2	1:A:186:LEU:HG	2.03	0.40
1:B:128:GLU:HG2	1:B:136:TYR:OH	2.20	0.40
1:B:201:ARG:HE	1:B:201:ARG:CA	2.15	0.40
1:A:24:ARG:HG2	1:A:243:ASN:OD1	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/245 (86%)	196 (93%)	11 (5%)	4 (2%)	6	8
1	B	215/245 (88%)	198 (92%)	13 (6%)	4 (2%)	6	8
All	All	426/490 (87%)	394 (92%)	24 (6%)	8 (2%)	6	8

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	81	ASP
1	A	230	GLU
1	B	235	SER
1	B	247	GLN
1	A	221	ASP
1	A	247	GLN
1	B	230	GLU
1	A	233	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/204 (88%)	167 (93%)	13 (7%)	13	23
1	B	183/204 (90%)	174 (95%)	9 (5%)	22	39
All	All	363/408 (89%)	341 (94%)	22 (6%)	17	30

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	LEU
1	A	35	LEU
1	A	37	THR
1	A	40	LEU
1	A	72	ARG
1	A	83	THR
1	A	98	ILE
1	A	111	VAL
1	A	118	TRP
1	A	123	VAL
1	A	212	VAL
1	A	229	ASP
1	A	241	VAL
1	B	31	GLN
1	B	35	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	98	ILE
1	B	102	GLN
1	B	108	ILE
1	B	123	VAL
1	B	130	ASP
1	B	201	ARG
1	B	210	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	79	GLN
1	A	106	ASN
1	A	159	ASN
1	A	176	GLN
1	A	207	GLN
1	B	27	GLN
1	B	31	GLN
1	B	79	GLN
1	B	102	GLN
1	B	107	ASN
1	B	159	ASN
1	B	176	GLN
1	B	187	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	249	-	4,4,4	1.55	1 (25%)	6,6,6	1.39	1 (16%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	249	PO4	P-O1	2.94	1.57	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	249	PO4	O3-P-O2	2.48	115.63	107.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/245 (88%)	0.38	20 (9%) 14 12	23, 41, 73, 90	0
1	B	219/245 (89%)	0.37	19 (8%) 16 13	19, 40, 74, 84	0
All	All	436/490 (88%)	0.38	39 (8%) 15 12	19, 40, 74, 90	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	234	SER	4.6
1	A	230	GLU	4.5
1	B	82	GLY	4.2
1	A	235	SER	4.0
1	B	59	VAL	3.7
1	B	247	GLN	3.7
1	A	35	LEU	3.6
1	A	231	ASP	3.4
1	B	235	SER	3.4
1	A	229	ASP	3.3
1	B	106	ASN	3.2
1	A	232	LEU	3.2
1	B	69	ARG	3.1
1	A	59	VAL	3.1
1	B	79	GLN	3.1
1	A	233	VAL	3.0
1	B	202	GLY	2.9
1	A	246	LEU	2.7
1	A	248	LYS	2.7
1	A	80	GLY	2.6
1	A	228	VAL	2.6
1	B	203	GLY	2.5
1	B	234	SER	2.5
1	B	83	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	117	ALA	2.5
1	B	200	ILE	2.4
1	B	201	ARG	2.4
1	A	205	ARG	2.4
1	A	21	PRO	2.3
1	B	22	PRO	2.3
1	A	217	GLU	2.2
1	A	67	GLY	2.2
1	B	80	GLY	2.2
1	B	78	ALA	2.2
1	A	37	THR	2.1
1	B	68	PHE	2.1
1	A	83	THR	2.1
1	B	81	ASP	2.1
1	A	43	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	A	249	5/5	0.88	0.11	75,75,75,76	0
3	CL	A	250	1/1	0.97	0.04	38,38,38,38	0
3	CL	B	249	1/1	1.00	0.06	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.