



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 12:01 PM UTC

PDB ID : 1WEK / pdb_00001wek
Title : Crystal structure of the conserved hypothetical protein TT1465 from *Thermus thermophilus* HB8
Authors : Kukimoto-Niino, M.; Murayama, K.; Kato-Murayama, M.; Terada, T.; Shirouzu, M.; Kuramitsu, S.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2004-05-25
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	217	 64% 29% • •
1	B	217	 62% 29% 5% 5%
1	C	217	 63% 28% • 5%
1	D	217	 62% 29% 5% •
1	E	217	 64% 26% 5% 5%
1	F	217	 65% 26% • 6%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 10248 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hypothetical protein TT1465.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	208	Total	C	N	O	Se	0	0	0
			1651	1076	281	293	1			
1	B	207	Total	C	N	O	Se	0	0	0
			1644	1071	280	292	1			
1	C	206	Total	C	N	O	Se	0	0	0
			1632	1066	277	288	1			
1	D	208	Total	C	N	O	Se	0	0	0
			1646	1074	279	292	1			
1	E	206	Total	C	N	O	Se	0	0	0
			1629	1062	276	290	1			
1	F	205	Total	C	N	O	Se	0	0	0
			1624	1060	276	287	1			

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0
2	E	1	Total O P 5 4 1	0	0
2	E	1	Total O P 5 4 1	0	0
2	F	1	Total O P 5 4 1	0	0
2	F	1	Total O P 5 4 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	75	Total O 75 75	0	0
3	B	56	Total O 56 56	0	0
3	C	48	Total O 48 48	0	0
3	D	51	Total O 51 51	0	0
3	E	65	Total O 65 65	0	0
3	F	67	Total O 67 67	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: hypothetical protein TT1465




- Molecule 1: hypothetical protein TT1465




- Molecule 1: hypothetical protein TT1465




• Molecule 1: hypothetical protein TT1465



• Molecule 1: hypothetical protein TT1465



• Molecule 1: hypothetical protein TT1465



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.65Å 83.80Å 265.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.94 – 2.20	Depositor
% Data completeness (in resolution range)	93.9 (14.94-2.20)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.198 , 0.250	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	10248	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/1695	1.01	11/2298 (0.5%)
1	B	0.48	0/1687	0.97	5/2287 (0.2%)
1	C	0.46	0/1674	0.94	3/2266 (0.1%)
1	D	0.45	0/1688	0.94	4/2286 (0.2%)
1	E	0.46	0/1671	1.01	4/2264 (0.2%)
1	F	0.49	0/1666	0.94	7/2255 (0.3%)
All	All	0.47	0/10081	0.97	34/13656 (0.2%)

There are no bond length outliers.

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	133	VAL	N-CA-C	-7.79	104.94	113.43
1	A	106	GLN	N-CA-C	-7.64	97.86	110.17
1	A	120	TYR	N-CA-C	6.86	120.11	109.07
1	D	190	GLY	CA-C-N	6.68	127.22	119.47
1	D	190	GLY	C-N-CA	6.68	127.22	119.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1651	0	1627	62	0
1	B	1644	0	1619	56	0
1	C	1632	0	1619	53	0
1	D	1646	0	1629	60	0
1	E	1629	0	1605	50	0
1	F	1624	0	1608	53	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	15	0	0	1	0
2	D	15	0	0	1	0
2	E	10	0	0	0	0
2	F	10	0	0	0	0
3	A	75	0	0	3	0
3	B	56	0	0	3	0
3	C	48	0	0	3	0
3	D	51	0	0	1	0
3	E	65	0	0	1	0
3	F	67	0	0	4	0
All	All	10248	0	9707	300	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 300 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:200:THR:HG22	1:C:202:GLU:H	1.24	1.03
1:D:81:MSE:HE2	1:D:81:MSE:HA	1.52	0.88
1:F:202:GLU:HB2	1:F:205:GLU:HG3	1.55	0.87
1:A:107:LYS:H	1:A:107:LYS:HD2	1.37	0.86
1:C:159:LYS:NZ	1:D:102:LEU:HD23	1.95	0.81

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/217 (95%)	197 (96%)	8 (4%)	1 (0%)	24	27
1	B	205/217 (94%)	194 (95%)	7 (3%)	4 (2%)	6	4
1	C	202/217 (93%)	196 (97%)	5 (2%)	1 (0%)	24	27
1	D	204/217 (94%)	194 (95%)	9 (4%)	1 (0%)	24	27
1	E	202/217 (93%)	196 (97%)	4 (2%)	2 (1%)	12	11
1	F	201/217 (93%)	194 (96%)	5 (2%)	2 (1%)	12	11
All	All	1220/1302 (94%)	1171 (96%)	38 (3%)	11 (1%)	14	14

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	105	GLU
1	C	169	ASP
1	E	169	ASP
1	E	214	ALA
1	F	169	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/175 (97%)	161 (95%)	8 (5%)	23	31
1	B	168/175 (96%)	156 (93%)	12 (7%)	13	16
1	C	167/175 (95%)	155 (93%)	12 (7%)	13	15
1	D	168/175 (96%)	157 (94%)	11 (6%)	15	18
1	E	166/175 (95%)	152 (92%)	14 (8%)	10	11
1	F	166/175 (95%)	156 (94%)	10 (6%)	17	21
All	All	1004/1050 (96%)	937 (93%)	67 (7%)	15	17

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	13	GLU
1	F	116	LEU
1	F	196	LEU
1	C	124	ARG
1	C	116	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	99	ASN
1	F	208	GLN
1	D	195	GLN
1	F	9	GLN
1	D	114	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	A	501	-	4,4,4	1.66	0	6,6,6	0.46	0
2	PO4	C	511	-	4,4,4	1.75	1 (25%)	6,6,6	0.47	0
2	PO4	F	502	-	4,4,4	1.67	1 (25%)	6,6,6	0.47	0
2	PO4	C	505	-	4,4,4	1.76	2 (50%)	6,6,6	0.50	0
2	PO4	E	504	-	4,4,4	1.44	0	6,6,6	0.46	0
2	PO4	D	506	-	4,4,4	1.61	0	6,6,6	0.49	0
2	PO4	F	510	-	4,4,4	1.64	0	6,6,6	0.46	0
2	PO4	D	512	-	4,4,4	1.61	0	6,6,6	0.42	0
2	PO4	D	508	-	4,4,4	1.72	1 (25%)	6,6,6	0.49	0
2	PO4	B	503	-	4,4,4	1.58	0	6,6,6	0.47	0
2	PO4	C	507	-	4,4,4	1.68	0	6,6,6	0.48	0
2	PO4	E	509	-	4,4,4	1.60	0	6,6,6	0.44	0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	508	PO4	P-O4	-2.11	1.48	1.54
2	C	505	PO4	P-O2	-2.04	1.48	1.54
2	C	505	PO4	P-O3	-2.02	1.48	1.54
2	C	511	PO4	P-O4	-2.01	1.48	1.54
2	F	502	PO4	P-O4	-2.00	1.48	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	512	PO4	1	0
2	C	507	PO4	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.