



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 29, 2026 – 12:20 AM UTC

PDB ID : 5WE2 / pdb_00005we2
Title : Structural Basis for Telomere Length Regulation by the Shelterin Bridge
Authors : Kim, J.-K.; Liu, J.; Hu, X.; Sankaran, B.; Qiao, F.
Deposited on : 2017-07-06
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	6% 73% 18% 8%
1	C	249	10% 69% 20% 8%
2	B	33	9% 79% 15%
2	D	33	3% 61% 27% 12%
3	F	30	10% 13% 80%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4450 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protection of telomeres protein poz1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	0	1	0
			1934	1233	334	350	17			
1	C	229	Total	C	N	O	S	0	0	0
			1918	1218	332	353	15			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP O13852
C	1	SER	-	expression tag	UNP O13852

- Molecule 2 is a protein called Protection of telomeres protein tpz1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	32	Total	C	N	O	S	0	0	0
			270	172	48	47	3			
2	D	29	Total	C	N	O	S	0	0	0
			246	158	44	41	3			

- Molecule 3 is a protein called DNA-binding protein rap1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	6	Total	C	N	O	0	0	0
			46	28	6	12			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	D	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	14	Total 14	O 14	0	0
5	B	6	Total 6	O 6	0	0
5	C	13	Total 13	O 13	0	0
5	D	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protection of telomeres protein poz1



- Molecule 1: Protection of telomeres protein poz1




- Molecule 2: Protection of telomeres protein tpz1



- Molecule 2: Protection of telomeres protein tpz1





● Molecule 3: DNA-binding protein rap1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.36Å 95.39Å 109.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.67 – 2.50 45.67 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (45.67-2.50) 94.0 (45.67-2.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.209 , 0.263 0.221 , 0.266	Depositor DCC
R_{free} test set	2000 reflections (6.50%)	wwPDB-VP
Wilson B-factor (Å ²)	51.2	Xtriage
Anisotropy	0.521	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4450	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	1/1973 (0.1%)	0.93	3/2645 (0.1%)
1	C	0.57	0/1956	0.87	1/2624 (0.0%)
2	B	0.67	0/276	0.70	0/366
2	D	0.53	0/252	0.80	0/335
3	F	0.82	0/45	1.55	1/59 (1.7%)
All	All	0.62	1/4502 (0.0%)	0.89	5/6029 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	77	LEU	CA-C	5.09	1.57	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	77	LEU	C-N-CD	-8.22	91.29	125.00
1	A	77	LEU	CA-C-N	7.61	129.12	120.98
1	A	77	LEU	C-N-CA	7.61	129.12	120.98
3	F	478	LEU	N-CA-C	7.26	119.89	110.53
1	A	75	GLN	N-CA-C	6.16	117.66	111.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1934	0	1927	37	0
1	C	1918	0	1886	54	0
2	B	270	0	261	4	0
2	D	246	0	234	6	0
3	F	46	0	40	3	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	A	14	0	0	1	0
5	B	6	0	0	0	0
5	C	13	0	0	1	0
5	D	1	0	0	0	0
All	All	4450	0	4348	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:182:MET:HA	1:C:185:LEU:HB2	1.57	0.85
1:A:6:ARG:HG3	1:C:16:THR:HG22	1.58	0.85
1:C:137:ALA:HB2	1:C:226:GLN:HE22	1.48	0.78
1:C:79:ILE:CD1	1:C:81:GLN:HB3	2.17	0.75
1:A:236:LYS:NZ	1:A:238:ARG:HH21	1.83	0.75
1:A:236:LYS:HZ3	1:A:238:ARG:HH21	1.35	0.73
1:A:14:LEU:HD13	1:A:56:MET:HE1	1.71	0.71
3:F:476:GLU:O	3:F:478:LEU:HD12	1.89	0.71
2:B:477:GLU:HG3	2:B:478:ALA:H	1.56	0.70
1:C:52:MET:HE2	2:D:490:SER:HB3	1.72	0.69
1:A:54:ILE:HD11	1:A:106:ILE:HA	1.75	0.67
1:C:177:HIS:HD2	1:C:181:GLU:HB3	1.60	0.67
1:C:79:ILE:HD12	1:C:79:ILE:C	2.20	0.66
1:C:79:ILE:HD11	1:C:81:GLN:HB3	1.78	0.65
1:C:76:ASN:OD1	1:C:76:ASN:N	2.29	0.63
1:C:159:GLN:NE2	5:C:301:HOH:O	2.30	0.63
2:D:479:CYS:HB3	2:D:482:CYS:HB2	1.83	0.61
1:C:150:ARG:HH12	1:C:207:MET:HB2	1.67	0.60
1:A:137:ALA:HB2	1:A:226:GLN:HE22	1.66	0.60
1:C:238:ARG:O	1:C:241:VAL:HG22	2.01	0.60
1:C:241:VAL:HG23	1:C:242:LYS:H	1.66	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:144:ALA:HB1	1:C:216:LEU:HD13	1.86	0.58
1:A:74:PHE:CD1	1:A:77:LEU:HB2	2.39	0.57
1:C:109:GLU:OE1	1:C:233:TYR:OH	2.17	0.56
1:A:236:LYS:HZ3	1:A:238:ARG:NH2	2.04	0.56
1:A:165:ILE:HD12	1:A:165:ILE:H	1.72	0.55
1:C:79:ILE:HD12	1:C:79:ILE:O	2.07	0.55
1:C:114:MET:HE1	1:C:230:PHE:CD2	2.41	0.55
1:A:6:ARG:HA	1:C:16:THR:HG21	1.89	0.55
1:C:114:MET:O	1:C:130:LYS:NZ	2.38	0.55
1:C:22:ASN:O	1:C:224:LYS:NZ	2.41	0.54
1:C:76:ASN:O	1:C:78:PRO:HD3	2.08	0.53
2:D:486:LEU:O	2:D:488:HIS:ND1	2.38	0.53
1:C:135:ASN:HA	1:C:138:LEU:HD12	1.91	0.53
1:A:114:MET:HE1	1:A:230:PHE:CD2	2.43	0.53
1:A:27:GLN:HB3	1:A:29:GLU:OE2	2.08	0.53
1:A:236:LYS:HZ2	1:A:236:LYS:HB3	1.73	0.53
1:C:187:ARG:HH11	1:C:187:ARG:HG3	1.73	0.53
1:C:15:GLU:HG3	1:C:19:ILE:HD12	1.90	0.52
1:C:130:LYS:HB2	1:C:230:PHE:CZ	2.45	0.52
1:A:22:ASN:O	1:A:224:LYS:NZ	2.43	0.52
1:A:171:LEU:HD22	1:A:182:MET:SD	2.51	0.50
1:C:150:ARG:NH1	1:C:207:MET:HB2	2.26	0.50
1:C:77:LEU:HD23	1:C:78:PRO:N	2.27	0.50
1:C:39:ARG:HA	2:D:495:LEU:HD22	1.93	0.49
1:C:137:ALA:HB2	1:C:226:GLN:NE2	2.23	0.49
1:C:26:MET:CE	2:D:498:TRP:HE1	2.26	0.49
1:A:72:SER:HB3	1:A:211:LYS:HD3	1.95	0.49
2:B:495:LEU:O	2:B:499:LYS:HG3	2.13	0.48
1:A:29:GLU:H	1:A:29:GLU:CD	2.20	0.48
1:C:150:ARG:HH12	1:C:207:MET:CB	2.26	0.48
1:C:187:ARG:HG3	1:C:187:ARG:NH1	2.29	0.48
1:A:19:ILE:HG22	1:A:236:LYS:HD3	1.96	0.48
1:A:14:LEU:HD13	1:A:56:MET:CE	2.42	0.48
1:C:31:SER:O	1:C:35:ASN:ND2	2.42	0.48
1:C:241:VAL:HG23	1:C:242:LYS:N	2.29	0.47
1:A:208:PRO:HD2	1:A:211:LYS:HG2	1.95	0.47
3:F:476:GLU:HG2	3:F:477:ASP:H	1.79	0.47
1:C:79:ILE:CD1	1:C:79:ILE:C	2.86	0.47
1:C:137:ALA:CB	1:C:226:GLN:HE22	2.24	0.47
1:C:181:GLU:O	1:C:185:LEU:N	2.46	0.46
1:A:114:MET:HE1	1:A:230:PHE:HB2	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:PHE:CD2	1:A:102[A]:ARG:HD2	2.50	0.46
1:A:151:TRP:CZ3	1:A:204:LYS:HE2	2.52	0.45
1:C:177:HIS:HD2	1:C:181:GLU:CB	2.26	0.45
1:A:11:LEU:HD23	1:A:11:LEU:HA	1.83	0.44
1:C:139:LYS:NZ	3:F:479:GLU:HB3	2.32	0.44
1:C:159:GLN:O	1:C:196:ARG:NH2	2.51	0.44
1:A:54:ILE:HG22	5:A:314:HOH:O	2.17	0.44
1:A:14:LEU:HD23	1:A:14:LEU:HA	1.78	0.44
1:A:16:THR:HG21	1:C:6:ARG:HA	2.00	0.43
1:A:183:GLU:O	1:A:187:ARG:HG3	2.18	0.43
1:C:14:LEU:HD13	1:C:56:MET:HE3	2.01	0.43
1:C:160:GLU:HA	1:C:161:PRO:HA	1.79	0.43
2:D:481:MET:HE3	2:D:481:MET:HB2	1.79	0.43
1:A:74:PHE:HE1	1:A:78:PRO:O	2.00	0.43
1:C:140:LEU:HB3	1:C:219:TYR:HB2	2.00	0.43
2:B:477:GLU:HG3	2:B:478:ALA:N	2.29	0.43
1:A:238:ARG:H	1:A:238:ARG:HG2	1.59	0.43
1:A:102[B]:ARG:O	1:A:106:ILE:HG13	2.18	0.42
1:C:228:GLU:HA	1:C:233:TYR:O	2.19	0.42
1:C:150:ARG:NH1	1:C:207:MET:CB	2.83	0.42
1:C:177:HIS:HB3	1:C:181:GLU:HB2	2.02	0.42
1:C:199:TYR:CE1	1:C:203:HIS:CD2	3.08	0.42
1:A:159:GLN:O	1:A:196:ARG:NH2	2.53	0.41
1:C:14:LEU:HD23	1:C:14:LEU:HA	1.91	0.41
1:C:79:ILE:HD13	1:C:81:GLN:HB3	1.96	0.41
1:C:171:LEU:HD23	1:C:171:LEU:HA	1.94	0.41
1:A:28:ARG:HD3	1:A:28:ARG:HA	1.64	0.41
1:A:141:TYR:HB2	1:A:219:TYR:CE1	2.56	0.40
2:B:506:ASN:HB2	2:B:507:LYS:H	1.76	0.40
1:A:87:GLN:H	1:A:87:GLN:CD	2.29	0.40
1:C:163:ASP:OD2	1:C:166:ASN:HB2	2.21	0.40
1:A:25:ASP:OD1	1:A:26:MET:N	2.54	0.40
1:A:199:TYR:CE2	1:A:203:HIS:CE1	3.10	0.40
1:C:14:LEU:HD13	1:C:56:MET:CE	2.52	0.40
1:C:26:MET:HE2	1:C:34:VAL:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/249 (90%)	217 (97%)	6 (3%)	1 (0%)	30	49
1	C	222/249 (89%)	212 (96%)	9 (4%)	1 (0%)	24	43
2	B	30/33 (91%)	27 (90%)	3 (10%)	0	100	100
2	D	27/33 (82%)	26 (96%)	1 (4%)	0	100	100
3	F	4/30 (13%)	4 (100%)	0	0	100	100
All	All	507/594 (85%)	486 (96%)	19 (4%)	2 (0%)	30	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	78	PRO
1	A	22	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/237 (92%)	215 (98%)	4 (2%)	51	77
1	C	215/237 (91%)	204 (95%)	11 (5%)	21	43
2	B	29/30 (97%)	28 (97%)	1 (3%)	32	60
2	D	26/30 (87%)	25 (96%)	1 (4%)	29	56
3	F	5/29 (17%)	4 (80%)	1 (20%)	1	2
All	All	494/563 (88%)	476 (96%)	18 (4%)	30	58

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	LYS
1	A	142	SER
1	A	180	LYS
1	A	190	LYS
2	B	506	ASN
1	C	26	MET
1	C	28	ARG
1	C	44	SER
1	C	76	ASN
1	C	130	LYS
1	C	139	LYS
1	C	190	LYS
1	C	205	LYS
1	C	215	VAL
1	C	238	ARG
1	C	242	LYS
2	D	480	GLU
3	F	480	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	GLN
1	A	12	ASN
1	C	8	GLN
1	C	155	HIS
1	C	177	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	C	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	70:ASP	C	71:PRO	N	3.02

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	229/249 (91%)	0.43	15 (6%) 24 21	21, 57, 89, 99	2 (0%)
1	C	229/249 (91%)	0.79	26 (11%) 10 8	42, 68, 105, 137	5 (2%)
2	B	32/33 (96%)	0.28	3 (9%) 14 12	42, 49, 97, 109	0
2	D	29/33 (87%)	0.57	1 (3%) 48 43	44, 59, 91, 102	0
3	F	6/30 (20%)	2.55	3 (50%) 0 0	109, 119, 126, 129	0
All	All	525/594 (88%)	0.61	48 (9%) 15 13	21, 62, 100, 137	7 (1%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	74	PHE	5.0
3	F	480	ILE	4.8
1	C	77	LEU	4.7
1	C	71	PRO	4.7
1	C	182	MET	4.6
1	C	78	PRO	4.4
1	A	127	LEU	4.4
1	C	241	VAL	4.4
1	A	76	ASN	4.3
1	A	79	ILE	4.1
1	C	74	PHE	3.8
1	C	243	MET	3.6
1	C	79	ILE	3.6
1	C	81	GLN	3.6
1	C	82	THR	3.3
1	C	76	ASN	3.0
1	A	180	LYS	3.0
1	C	130	LYS	2.9
1	C	83	GLN	2.9
1	A	182	MET	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	476	SER	2.8
3	F	478	LEU	2.8
1	C	207	MET	2.8
1	C	21	GLU	2.7
1	C	171	LEU	2.7
1	C	135	ASN	2.5
3	F	476	GLU	2.5
1	C	177	HIS	2.5
1	C	180	LYS	2.4
1	A	77	LEU	2.4
1	C	240	ILE	2.4
2	B	478	ALA	2.4
1	A	139	LYS	2.4
1	C	187	ARG	2.4
1	A	93	TYR	2.3
1	C	186	GLN	2.3
1	A	75	GLN	2.2
1	C	242	LYS	2.2
1	C	202	GLN	2.2
1	A	115	ARG	2.2
1	A	187	ARG	2.2
2	D	484	LEU	2.1
1	A	86	SER	2.1
1	C	80	THR	2.1
1	A	128	LEU	2.0
2	B	507	LYS	2.0
1	C	45	LYS	2.0
1	A	78	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ZN	D	601	1/1	0.98	0.04	87,87,87,87	0
4	ZN	B	601	1/1	1.00	0.01	52,52,52,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.