



wwPDB EM Validation Summary Report ⓘ

Mar 29, 2026 – 12:56 AM UTC

PDB ID : 7WEL / pdb_00007wel
EMDB ID : EMD-32451
Title : Human Nav1.8 with A-803467, class II
Authors : Yan, N.; Pan, X.J.; Huang, X.S.; Huang, G.X.
Deposited on : 2021-12-23
Resolution : 3.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	15020 (2.70 - 3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1956	 38% 17% 44%
2	B	2	 50% 100%
2	C	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	95T	A	2004	-	-	X	-
6	PCW	A	2008	-	-	X	-
6	PCW	A	2012	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 9815 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sodium channel protein type 10 subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1102	Total	C	N	O	S	1	0
			8882	5901	1417	1502	62		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	894	PHE	SER	conflict	UNP Q9Y5Y9

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
2	B	2	Total	C	N	O	0	0
			28	16	2	10		
2	C	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				AltConf
3	A	1	Total	C	N	O	0
			14	8	1	5	
3	A	1	Total	C	N	O	0
			14	8	1	5	
3	A	1	Total	C	N	O	0
			14	8	1	5	

- Molecule 4 is 5-(4-chlorophenyl)- {N}-(3,5-dimethoxyphenyl)furan-2-carboxamide (CCD ID: 95T) (formula: $C_{19}H_{16}ClNO_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
4	A	1	Total	C	Cl	N	O	0
			25	19	1	1	4	

- Molecule 5 is CHOLESTEROL (CCD ID: CLR) (formula: $C_{27}H_{46}O$).




Mol	Chain	Residues	Atoms			AltConf
5	A	1	Total	C	O	0
			28	27	1	
5	A	1	Total	C	O	0
			28	27	1	
5	A	1	Total	C	O	0
			28	27	1	
5	A	1	Total	C	O	0
			28	27	1	
5	A	1	Total	C	O	0
			28	27	1	

- Molecule 6 is 1,2-DIOLEOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PCW) (formula: $C_{44}H_{85}NO_8P$).




Mol	Chain	Residues	Atoms					AltConf
6	A	1	Total	C	N	O	P	0
			54	44	1	8	1	
6	A	1	Total	C	N	O	P	0
			47	37	1	8	1	
6	A	1	Total	C	O	P		0
			32	23	8	1		
6	A	1	Total	C	N	O	P	0
			54	44	1	8	1	
6	A	1	Total	C	O	P		0
			41	32	8	1		
6	A	1	Total	C	N	O	P	0
			40	30	1	8	1	
6	A	1	Total	C	N	O	P	0
			38	28	1	8	1	

- Molecule 7 is 1-O-OCTADECYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: LPE) (formula: $C_{26}H_{57}NO_6P$).



Mol	Chain	Residues	Atoms					AltConf
7	A	1	Total	C	N	O	P	0
			22	14	1	6	1	
7	A	1	Total	C	N	O	P	0
			22	14	1	6	1	
7	A	1	Total	C	N	O	P	0
			22	14	1	6	1	
7	A	1	Total	C	N	O	P	0
			22	14	1	6	1	
7	A	1	Total	C	O	P		0
			16	9	6	1		
7	A	1	Total	C	N	O	P	0
			23	15	1	6	1	
7	A	1	Total	C	N	O	P	0
			34	26	1	6	1	
7	A	1	Total	C	N	O	P	0
			22	14	1	6	1	
7	A	1	Total	C	N	O	P	0
			22	14	1	6	1	
7	A	1	Total	C	N	O	P	0
			21	13	1	6	1	
7	A	1	Total	C	N	O	P	0
			23	15	1	6	1	
7	A	1	Total	C	N	O	P	0
			21	13	1	6	1	

- Molecule 8 is O-[(R)-{[(2R)-2,3-bis(octadecanoyloxy)propyl]oxy}(hydroxy)phosphoryl]-L-serine (CCD ID: P5S) (formula: C₄₂H₈₂NO₁₀P).



Mol	Chain	Residues	Atoms					AltConf
8	A	1	Total	C	N	O	P	0
			49	37	1	10	1	
8	A	1	Total	C	N	O	P	0
			45	33	1	10	1	





- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:

100%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	141848	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.119	Depositor
Minimum map value	-0.046	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.0171	Depositor
Map size (Å)	259.8, 259.8, 259.8	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0825, 1.0825, 1.0825	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P5S, 95T, PCW, CLR, LPE, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	1/9106 (0.0%)	0.68	17/12349 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	385	SER	CA-C	-5.32	1.48	1.52

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1687	GLY	N-CA-C	9.67	124.33	112.73
1	A	155	LYS	N-CA-C	-8.93	104.44	114.62
1	A	1691	ASP	N-CA-C	-7.96	103.68	113.15
1	A	1589	ALA	N-CA-C	7.61	119.65	111.36
1	A	810	GLY	N-CA-C	6.69	120.26	112.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8882	0	9067	384	0
2	B	28	0	25	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	28	0	25	1	0
3	A	42	0	39	2	0
4	A	25	0	0	10	0
5	A	140	0	230	46	0
6	A	306	0	420	97	0
7	A	270	0	362	39	0
8	A	94	0	123	11	0
All	All	9815	0	10291	471	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 471 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:214:LEU:CD2	1:A:215:ARG:HH21	1.04	1.57
1:A:214:LEU:CD2	1:A:215:ARG:NH2	1.85	1.38
1:A:1171:PHE:CD2	7:A:2022:LPE:H2N2	1.72	1.23
1:A:214:LEU:HD21	1:A:215:ARG:HH21	1.12	1.15
1:A:1286:LEU:HD21	5:A:2006:CLR:H212	1.21	1.14

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1093/1956 (56%)	1043 (95%)	44 (4%)	6 (0%)	24 59

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	817	ARG
1	A	211	ILE
1	A	1632	VAL
1	A	294	LYS
1	A	1629	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	973/1727 (56%)	969 (100%)	4 (0%)	84 86

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	140	ILE
1	A	244	LYS
1	A	673	LEU
1	A	1370	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1362	GLN
1	A	1500	GLN
1	A	390	ASN
1	A	687	HIS
1	A	833	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	0.55	0	17,19,21	1.79	4 (23%)
2	NAG	B	2	2	14,14,15	0.37	0	17,19,21	0.98	1 (5%)
2	NAG	C	1	1,2	14,14,15	0.29	0	17,19,21	0.56	0
2	NAG	C	2	2	14,14,15	0.29	0	17,19,21	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	3/6/23/26	0/1/1/1
2	NAG	B	2	2	-	3/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C2-N2-C7	5.53	130.31	122.90
2	B	2	NAG	C4-C3-C2	-3.44	105.98	111.02
2	B	1	NAG	C4-C3-C2	-3.10	106.47	111.02
2	B	1	NAG	C1-C2-N2	2.23	113.95	110.43
2	B	1	NAG	O5-C1-C2	2.20	114.70	111.29

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C1-C2-N2-C7
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2


There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	1	0
2	C	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

30 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	CLR	A	2028	-	31,31,31	0.94	2 (6%)	48,48,48	1.42	7 (14%)
6	PCW	A	2027	-	37,37,53	1.12	2 (5%)	43,45,61	1.12	2 (4%)
7	LPE	A	2025	-	22,22,33	0.58	0	26,28,39	0.60	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	LPE	A	2014	-	21,21,33	0.64	0	25,27,39	0.72	0
7	LPE	A	2016	-	22,22,33	0.62	0	26,28,39	0.71	0
5	CLR	A	2021	-	31,31,31	0.87	1 (3%)	48,48,48	1.47	10 (20%)
3	NAG	A	2001	1	14,14,15	0.30	0	17,19,21	0.56	0
7	LPE	A	2022	-	21,21,33	0.68	0	25,27,39	0.80	1 (4%)
7	LPE	A	2023	-	21,21,33	0.58	0	25,27,39	0.71	0
3	NAG	A	2003	1	14,14,15	0.36	0	17,19,21	1.18	1 (5%)
5	CLR	A	2006	-	31,31,31	0.72	0	48,48,48	1.21	5 (10%)
7	LPE	A	2026	-	20,20,33	0.62	0	24,26,39	0.61	1 (4%)
6	PCW	A	2011	-	31,31,53	1.18	2 (6%)	34,36,61	1.32	3 (8%)
6	PCW	A	2019	-	39,39,53	1.09	2 (5%)	45,47,61	1.09	3 (6%)
7	LPE	A	2010	-	21,21,33	0.58	0	25,27,39	0.61	0
5	CLR	A	2018	-	31,31,31	0.89	1 (3%)	48,48,48	1.69	10 (20%)
4	95T	A	2004	-	27,27,27	2.52	10 (37%)	37,37,37	4.17	16 (43%)
7	LPE	A	2015	-	15,15,33	0.58	0	16,18,39	0.88	1 (6%)
6	PCW	A	2007	-	53,53,53	0.92	2 (3%)	59,61,61	1.11	4 (6%)
7	LPE	A	2013	-	21,21,33	0.58	0	25,27,39	0.61	0
5	CLR	A	2005	-	31,31,31	0.73	0	48,48,48	1.18	5 (10%)
6	PCW	A	2008	-	46,46,53	1.02	2 (4%)	52,54,61	1.05	3 (5%)
6	PCW	A	2012	-	53,53,53	0.94	2 (3%)	59,61,61	1.00	4 (6%)
7	LPE	A	2024	-	20,20,33	0.60	0	24,26,39	0.62	0
7	LPE	A	2009	-	21,21,33	0.58	0	25,27,39	0.54	0
7	LPE	A	2020	-	33,33,33	0.47	0	37,39,39	0.73	0
8	P5S	A	2030	-	43,44,53	0.95	3 (6%)	45,51,60	1.49	8 (17%)
8	P5S	A	2029	-	47,48,53	1.00	2 (4%)	49,55,60	1.37	4 (8%)
6	PCW	A	2017	-	40,40,53	1.02	2 (5%)	43,45,61	1.28	4 (9%)
3	NAG	A	2002	1	14,14,15	0.49	0	17,19,21	1.52	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CLR	A	2028	-	-	6/10/68/68	0/4/4/4
6	PCW	A	2027	-	-	17/41/41/57	-
7	LPE	A	2025	-	-	3/23/23/34	-
7	LPE	A	2014	-	-	7/22/22/34	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	LPE	A	2016	-	-	8/23/23/34	-
5	CLR	A	2021	-	-	4/10/68/68	0/4/4/4
3	NAG	A	2001	1	-	2/6/23/26	0/1/1/1
7	LPE	A	2022	-	-	10/22/22/34	-
7	LPE	A	2023	-	-	4/22/22/34	-
3	NAG	A	2003	1	-	2/6/23/26	0/1/1/1
5	CLR	A	2006	-	-	5/10/68/68	0/4/4/4
7	LPE	A	2026	-	-	6/21/21/34	-
6	PCW	A	2011	-	-	7/33/33/57	-
6	PCW	A	2019	-	-	11/43/43/57	-
7	LPE	A	2010	-	-	3/22/22/34	-
5	CLR	A	2018	-	-	7/10/68/68	0/4/4/4
4	95T	A	2004	-	-	6/16/16/16	0/3/3/3
7	LPE	A	2015	-	-	7/14/14/34	-
6	PCW	A	2007	-	-	24/57/57/57	-
7	LPE	A	2013	-	-	6/22/22/34	-
5	CLR	A	2005	-	-	0/10/68/68	0/4/4/4
6	PCW	A	2008	-	-	13/50/50/57	-
6	PCW	A	2012	-	-	29/57/57/57	-
7	LPE	A	2024	-	-	5/21/21/34	-
7	LPE	A	2009	-	-	13/22/22/34	-
7	LPE	A	2020	-	-	3/34/34/34	-
8	P5S	A	2030	-	-	12/50/50/59	-
8	P5S	A	2029	-	-	7/54/54/59	-
6	PCW	A	2017	-	-	14/42/42/57	-
3	NAG	A	2002	1	-	1/6/23/26	0/1/1/1

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	2004	95T	O02-C09	-7.98	1.25	1.38
4	A	2004	95T	O02-C07	-5.53	1.28	1.38
8	A	2029	P5S	O37-C38	4.47	1.46	1.34
8	A	2029	P5S	O19-C17	4.41	1.46	1.33
6	A	2008	PCW	O3-C11	4.35	1.46	1.33

The worst 5 of 95 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2004	95T	C09-C13-N06	10.78	139.84	115.47
4	A	2004	95T	O02-C07-C10	10.19	128.80	116.73
4	A	2004	95T	O02-C09-C13	9.14	132.35	117.52
4	A	2004	95T	C08-N06-C13	8.23	142.06	127.45
4	A	2004	95T	O05-C13-N06	-8.06	104.76	123.89

There are no chirality outliers.

5 of 242 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	2018	CLR	C13-C17-C20-C21
5	A	2018	CLR	C16-C17-C20-C22
6	A	2007	PCW	O4P-C4-C5-N
6	A	2007	PCW	C4-O4P-P-O2P
6	A	2008	PCW	C1-O3P-P-O1P

There are no ring outliers.

27 monomers are involved in 199 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	2028	CLR	7	0
6	A	2027	PCW	6	0
7	A	2025	LPE	7	0
7	A	2014	LPE	4	0
7	A	2016	LPE	1	0
5	A	2021	CLR	2	0
3	A	2001	NAG	2	0
7	A	2022	LPE	14	0
7	A	2023	LPE	5	0
5	A	2006	CLR	10	0
7	A	2026	LPE	3	0
6	A	2011	PCW	16	0
6	A	2019	PCW	11	0
7	A	2010	LPE	1	0
5	A	2018	CLR	9	0
4	A	2004	95T	10	0
7	A	2015	LPE	1	0
6	A	2007	PCW	14	0
7	A	2013	LPE	4	0
5	A	2005	CLR	18	0
6	A	2008	PCW	29	0
6	A	2012	PCW	32	0

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	2009	LPE	1	0
7	A	2020	LPE	1	0
8	A	2030	P5S	7	0
8	A	2029	P5S	4	0
6	A	2017	PCW	8	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand PCW A 2027


























































Ligand P5S A 2030	
 Bond lengths	 Bond angles
 Torsions	 Rings
Ligand P5S A 2029	
 Bond lengths	 Bond angles
 Torsions	 Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-32451. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 120



Y Index: 120




Z Index: 120


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 134



Y Index: 130




Z Index: 131


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0171. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 65 nm³; this corresponds to an approximate mass of 58 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.312 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-32451 and PDB model 7WEL. Per-residue inclusion information can be found in [section 3](#) on [page 10](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0171 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0171).

9.4 Atom inclusion ⓘ



At the recommended contour level, 84% of all backbone atoms, 73% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0171) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7300	0.4870
A	0.7310	0.4880
B	0.3930	0.3020
C	0.7140	0.4100

