



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:15 AM UTC

PDB ID : 1WFC / pdb_00001wfc
Title : STRUCTURE OF APO, UNPHOSPHORYLATED, P38 MITOGEN ACTIVATED PROTEIN KINASE P38 (P38 MAP KINASE) THE MAMMALIAN HOMOLOGUE OF THE YEAST HOG1 PROTEIN
Authors : Wilson, K.P.; Fitzgibbon, M.J.; Caron, P.R.; Griffith, J.P.; Chen, W.; McCaffrey, P.G.; Chambers, S.P.; Su, M.S.-S.
Deposited on : 1996-09-13
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	366	 71% 20% .. 7%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MITOGEN-ACTIVATED PROTEIN KINASE P38.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	340	Total	C	N	O	S	31	0	0
			2746	1760	465	508	13			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	186	Total	O	0	0
			186	186		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: MITOGEN-ACTIVATED PROTEIN KINASE P38

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	45.19Å 86.25Å 123.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.30	Depositor
% Data completeness (in resolution range)	90.0 (6.00-2.30)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.213 , 0.297	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2932	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.12	17/2808 (0.6%)	1.67	42/3811 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	337	TRP	CD2-CE2	6.93	1.53	1.41
1	A	18	TRP	CD2-CE2	6.72	1.52	1.41
1	A	64	HIS	CD2-NE2	-6.67	1.30	1.37
1	A	187	TRP	CD2-CE2	6.62	1.52	1.41
1	A	228	HIS	CD2-NE2	-6.52	1.30	1.37
1	A	142	HIS	CD2-NE2	-6.34	1.30	1.37
1	A	207	TRP	CD2-CE2	6.34	1.52	1.41
1	A	197	TRP	CD2-CE2	6.27	1.52	1.41
1	A	80	HIS	CD2-NE2	-6.26	1.30	1.37
1	A	107	HIS	CD2-NE2	-6.22	1.31	1.37
1	A	305	HIS	CD2-NE2	-6.06	1.31	1.37
1	A	91	THR	C-N	-6.03	1.28	1.34
1	A	126	HIS	CD2-NE2	-6.00	1.31	1.37
1	A	148	HIS	CD2-NE2	-5.96	1.31	1.37
1	A	312	HIS	CD2-NE2	-5.88	1.31	1.37
1	A	349	VAL	C-N	-5.43	1.28	1.33
1	A	77	HIS	CD2-NE2	-5.12	1.32	1.37

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	337	TRP	CG-CD2-CE3	10.67	144.57	133.90
1	A	18	TRP	CG-CD2-CE3	10.39	144.29	133.90
1	A	207	TRP	CG-CD2-CE3	10.04	143.94	133.90
1	A	197	TRP	CG-CD2-CE3	9.86	143.76	133.90
1	A	187	TRP	CG-CD2-CE3	9.63	143.53	133.90
1	A	18	TRP	CE2-CD2-CG	-8.86	96.57	107.20
1	A	337	TRP	CE2-CD2-CG	-8.82	96.61	107.20
1	A	207	TRP	CE2-CD2-CG	-8.74	96.71	107.20
1	A	187	TRP	CE2-CD2-CG	-8.57	96.91	107.20
1	A	169	PHE	CA-CB-CG	-8.55	105.25	113.80
1	A	197	TRP	CE2-CD2-CG	-8.54	96.95	107.20
1	A	13	LEU	N-CA-C	-8.17	100.28	111.56
1	A	301	GLN	OE1-CD-NE2	-8.15	114.45	122.60
1	A	149	ARG	NE-CZ-NH2	7.51	125.96	119.20
1	A	33	GLY	N-CA-C	-6.62	104.83	111.85
1	A	168	ASP	CA-CB-CG	-6.51	106.09	112.60
1	A	349	VAL	CA-C-N	6.50	124.40	119.66
1	A	349	VAL	C-N-CA	6.50	124.40	119.66
1	A	331	ASP	CA-CB-CG	-6.47	106.13	112.60
1	A	148	HIS	CB-CG-CD2	-6.24	123.09	131.20
1	A	312	HIS	CB-CG-CD2	-5.81	123.65	131.20
1	A	26	ASN	CA-CB-CG	-5.71	106.89	112.60
1	A	310	GLN	OE1-CD-NE2	-5.52	117.08	122.60
1	A	121	LYS	CG-CD-CE	5.45	123.84	111.30
1	A	155	ASN	OD1-CG-ND2	-5.45	117.15	122.60
1	A	183	VAL	N-CA-C	5.41	115.90	108.12
1	A	80	HIS	CB-CG-CD2	-5.38	124.21	131.20
1	A	199	HIS	CA-CB-CG	-5.38	108.42	113.80
1	A	187	TRP	CB-CG-CD2	-5.36	119.29	126.80
1	A	91	THR	O-C-N	-5.33	116.63	122.06
1	A	156	LEU	N-CA-C	-5.29	99.67	108.34
1	A	149	ARG	NE-CZ-NH1	-5.28	116.22	121.50
1	A	148	HIS	CB-CG-ND1	5.27	130.61	122.70
1	A	228	HIS	CB-CG-CD2	-5.26	124.37	131.20
1	A	196	ASN	CA-CB-CG	-5.19	107.41	112.60
1	A	305	HIS	CB-CG-CD2	-5.17	124.48	131.20
1	A	250	ILE	N-CA-C	-5.14	100.37	108.23
1	A	107	HIS	CB-CG-CD2	-5.12	124.54	131.20
1	A	147	ILE	O-C-N	-5.12	117.73	123.26
1	A	202	GLN	N-CA-C	5.08	118.57	112.38
1	A	321	ASP	CA-C-N	5.04	125.04	119.90
1	A	321	ASP	C-N-CA	5.04	125.04	119.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	169	PHE	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2746	0	2733	27	0
2	A	186	0	0	0	0
All	All	2932	0	2733	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:LEU:HD22	1:A:241:THR:HA	1.72	0.69
1:A:157:ALA:HB2	1:A:167:LEU:HD11	1.75	0.68
1:A:109:MET:HE2	1:A:157:ALA:HB1	1.77	0.67
1:A:32:SER:HA	1:A:37:SER:HA	1.84	0.59
1:A:236:LEU:HD13	1:A:241:THR:HG22	1.83	0.59
1:A:283:ASP:O	1:A:287:LYS:HG2	2.08	0.53
1:A:109:MET:HE3	1:A:110:GLY:H	1.75	0.52
1:A:298:THR:H	1:A:301:GLN:NE2	2.08	0.50
1:A:179:MET:HE3	1:A:186:ARG:HG3	1.94	0.48
1:A:13:LEU:O	1:A:16:THR:N	2.47	0.47
1:A:168:ASP:O	1:A:169:PHE:HB2	2.15	0.46
1:A:111:ALA:HB1	1:A:115:ASN:HB2	1.99	0.44
1:A:283:ASP:OD2	1:A:287:LYS:HE2	2.17	0.44
1:A:7:THR:O	1:A:21:PRO:HA	2.17	0.44
1:A:282:VAL:O	1:A:286:GLU:HG3	2.18	0.43
1:A:122:LEU:HD12	1:A:216:LEU:HD22	1.99	0.43
1:A:193:ILE:HG21	1:A:193:ILE:HD13	1.82	0.43
1:A:42:PHE:HD1	1:A:49:ARG:HD2	1.84	0.42
1:A:55:LEU:HD12	1:A:102:VAL:HB	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:ALA:O	1:A:164:LEU:HA	2.20	0.41
1:A:159:ASN:OD1	1:A:159:ASN:C	2.63	0.41
1:A:213:MET:SD	1:A:288:MET:HE1	2.61	0.41
1:A:332:LEU:HB2	1:A:337:TRP:CE2	2.56	0.41
1:A:10:ARG:NH2	1:A:17:ILE:HD13	2.36	0.41
1:A:88:ASP:HA	1:A:348:PHE:CE2	2.55	0.41
1:A:332:LEU:HB3	1:A:336:GLU:HB2	2.03	0.41
1:A:13:LEU:C	1:A:16:THR:N	2.79	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/366 (91%)	315 (94%)	17 (5%)	2 (1%)	21	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	306	ALA
1	A	168	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	302/325 (93%)	296 (98%)	6 (2%)	48 67

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	LYS
1	A	128	GLN
1	A	162	CYS
1	A	175	THR
1	A	204	VAL
1	A	265	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	126	HIS
1	A	228	HIS
1	A	301	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.