



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:55 AM UTC

PDB ID : 8WFP / pdb_00008wfp
Title : Crystal structure of polo-like kinase(PLK1)PBD in complex with DD-1
Authors : Park, J.; La, Y.K.; Bang, J.K.; Lee, S.J.
Deposited on : 2023-09-20
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	81% 17% .
1	B	226	79% 17% ..
2	C	7	14% 71% 14%
2	D	7	86% 14%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3961 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase PLK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	0	0	0
			1816	1150	316	340	10			
1	B	226	Total	C	N	O	S	0	0	0
			1821	1153	317	341	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	371	ALA	-	expression tag	UNP P53350
A	372	ALA	-	expression tag	UNP P53350
B	371	ALA	-	expression tag	UNP P53350
B	372	ALA	-	expression tag	UNP P53350

- Molecule 2 is a protein called DD-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	7	Total	C	N	O	P	0	0	1
			46	26	8	11	1			
2	D	7	Total	C	N	O	P	0	0	1
			46	26	8	11	1			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	115	Total	O	0	0
			115	115		
3	B	103	Total	O	0	0
			103	103		
3	C	6	Total	O	0	0
			6	6		
3	D	8	Total	O	0	0
			8	8		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase PLK1




- Molecule 1: Serine/threonine-protein kinase PLK1



- Molecule 2: DD-1



- Molecule 2: DD-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	77.83Å 35.37Å 95.55Å 90.00° 89.98° 90.00°	Depositor
Resolution (Å)	35.37 – 1.99 35.37 – 1.99	Depositor EDS
% Data completeness (in resolution range)	96.6 (35.37-1.99) 96.7 (35.37-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 1.98Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.198 , 0.229 0.198 , 0.227	Depositor DCC
R_{free} test set	1987 reflections (5.44%)	wwPDB-VP
Wilson B-factor (Å ²)	33.8	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.466 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3961	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1854	0.91	9/2508 (0.4%)
1	B	0.71	8/1859 (0.4%)	1.20	21/2515 (0.8%)
2	C	4.52	6/33 (18.2%)	2.07	2/45 (4.4%)
2	D	4.57	7/33 (21.2%)	2.03	2/45 (4.4%)
All	All	0.86	21/3779 (0.6%)	1.09	34/5113 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	LEU	C-N	13.27	1.49	1.33
2	C	3	LEU	C-N	13.16	1.49	1.33
2	D	2	PRO	C-N	9.58	1.48	1.33
2	C	2	PRO	N-CD	-9.10	1.35	1.47
2	D	2	PRO	N-CD	-9.04	1.35	1.47
2	C	2	PRO	C-N	9.04	1.47	1.33
2	C	4	HIS	C-N	8.91	1.45	1.33
2	D	4	HIS	C-N	8.05	1.44	1.33
1	B	488	GLU	CD-OE2	7.33	1.39	1.25
1	B	503	ASP	C-N	-6.24	1.24	1.33
1	B	504	GLU	CG-CD	6.23	1.67	1.52
1	B	503	ASP	CA-C	-5.91	1.46	1.53
1	B	474	LYS	CG-CD	-5.83	1.34	1.52
2	D	4	HIS	CB-CG	5.64	1.58	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	PRO	N-CA	5.57	1.55	1.47
1	B	505	LEU	CB-CG	5.50	1.64	1.53
2	D	1	ACE	C-N	5.40	1.48	1.43
2	C	4	HIS	CB-CG	5.30	1.57	1.50
1	B	388	LYS	C-O	-5.19	1.21	1.23
2	C	2	PRO	N-CA	5.07	1.54	1.47
1	B	399	GLU	CG-CD	-5.04	1.39	1.52

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	504	GLU	OE1-CD-OE2	-20.28	74.24	122.90
1	B	505	LEU	N-CA-C	-16.68	75.26	110.80
1	A	505	LEU	CB-CG-CD2	16.40	159.91	110.70
1	B	504	GLU	CG-CD-OE2	-15.81	82.03	118.40
1	B	504	GLU	CG-CD-OE1	15.47	153.98	118.40
1	B	504	GLU	N-CA-C	-12.88	93.71	111.87
1	B	505	LEU	CA-C-N	11.04	142.63	121.54
1	B	505	LEU	C-N-CA	11.04	142.63	121.54
1	B	474	LYS	CD-CE-NZ	-9.70	80.87	111.90
1	A	505	LEU	CA-CB-CG	9.65	150.06	116.30
1	A	505	LEU	CD1-CG-CD2	-9.44	90.03	110.80
1	A	505	LEU	CB-CG-CD1	-9.29	82.84	110.70
1	A	504	GLU	CA-C-N	9.21	136.69	122.24
1	A	504	GLU	C-N-CA	9.21	136.69	122.24
1	B	504	GLU	CA-CB-CG	8.72	131.55	114.10
1	B	505	LEU	N-CA-CB	8.59	125.00	110.49
1	A	505	LEU	N-CA-CB	-7.42	101.04	110.90
1	B	502	GLY	CA-C-N	7.06	132.67	122.08
1	B	502	GLY	C-N-CA	7.06	132.67	122.08
1	B	504	GLU	CB-CA-C	6.61	121.72	111.28
1	A	505	LEU	CB-CA-C	6.39	119.75	109.02
1	B	398	GLU	CA-C-N	-6.26	111.79	122.56
1	B	398	GLU	C-N-CA	-6.26	111.79	122.56
1	B	417	TYR	N-CA-CB	-6.03	101.38	110.91
1	B	505	LEU	CB-CG-CD2	5.91	128.42	110.70
1	B	488	GLU	CB-CA-C	-5.90	97.81	109.67
1	A	505	LEU	N-CA-C	-5.81	106.82	114.31
2	C	3	LEU	CA-C-N	-5.39	112.43	121.75
2	C	3	LEU	C-N-CA	-5.39	112.43	121.75
2	D	3	LEU	CA-C-N	-5.38	112.44	121.75
2	D	3	LEU	C-N-CA	-5.38	112.44	121.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	399	GLU	CA-C-N	-5.32	112.76	122.38
1	B	399	GLU	C-N-CA	-5.32	112.76	122.38
1	B	505	LEU	O-C-N	5.26	129.59	122.59

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	417	TYR	Sidechain
1	B	503	ASP	Peptide
1	B	504	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1816	0	1795	36	1
1	B	1821	0	1800	37	1
2	C	46	0	39	4	0
2	D	46	0	39	3	0
3	A	115	0	0	8	0
3	B	103	0	0	2	0
3	C	6	0	0	1	0
3	D	8	0	0	0	0
All	All	3961	0	3673	76	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:503:ASP:C	1:A:505:LEU:H	1.71	0.95
1:A:443:ILE:HD11	1:A:510:TYR:HB3	1.51	0.92
1:A:452:GLN:HE21	1:A:454:ILE:HD11	1.32	0.92

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:592:SER:OG	3:A:601:HOH:O	1.90	0.88
1:A:405:CYS:SG	3:A:679:HOH:O	2.31	0.86
1:A:434:VAL:HG23	1:A:479:LEU:HD13	1.58	0.86
1:A:460:GLU:OE1	1:A:507:ARG:NH2	2.08	0.86
1:A:456:ARG:NH2	3:A:603:HOH:O	2.10	0.84
1:B:434:VAL:HG23	1:B:479:LEU:HD13	1.59	0.83
1:A:488:GLU:HB3	1:A:489:HIS:HD2	1.48	0.77
1:A:513:THR:OG1	3:A:602:HOH:O	2.04	0.74
1:A:507:ARG:NH1	3:A:605:HOH:O	2.20	0.73
1:A:503:ASP:O	1:A:505:LEU:N	2.22	0.71
1:B:452:GLN:HE21	1:B:454:ILE:HD11	1.57	0.69
1:A:488:GLU:HB3	1:A:489:HIS:CD2	2.28	0.69
1:B:408:ILE:HD13	1:B:501:GLU:O	1.93	0.68
1:A:503:ASP:O	1:A:506:ALA:N	2.25	0.67
1:B:459:THR:OG1	3:B:601:HOH:O	2.13	0.66
1:B:503:ASP:OD1	1:B:506:ALA:HB2	1.95	0.65
1:B:416:ASP:OD1	1:B:418:SER:OG	2.13	0.62
1:A:452:GLN:NE2	1:A:454:ILE:HD11	2.08	0.62
1:A:384:VAL:HA	1:A:568:GLU:CD	2.24	0.62
1:B:419:ASP:OD1	1:B:420:LYS:N	2.33	0.61
1:B:592:SER:OG	3:B:602:HOH:O	2.16	0.61
1:A:463:LEU:HD12	1:A:469:PRO:HG3	1.86	0.58
1:A:414:TRP:CD1	2:C:5:SER:HB3	2.39	0.57
1:B:556:LYS:N	1:B:556:LYS:HD3	2.18	0.57
1:B:491:LEU:H	2:D:7:NH2:N	2.01	0.57
1:A:503:ASP:C	1:A:505:LEU:N	2.50	0.57
1:A:443:ILE:HD11	1:A:510:TYR:CB	2.29	0.57
3:A:629:HOH:O	1:B:589:LYS:HE3	2.04	0.57
1:A:507:ARG:NH1	3:A:604:HOH:O	2.19	0.57
2:C:4:HIS:HD2	2:C:5:SER:O	1.89	0.56
1:A:412:SER:HB2	1:A:492:LYS:HG3	1.89	0.55
1:B:501:GLU:OE1	1:B:503:ASP:HB2	2.06	0.55
2:D:4:HIS:HD2	2:D:5:SER:O	1.90	0.55
1:B:452:GLN:NE2	1:B:454:ILE:HD11	2.22	0.53
1:B:414:TRP:CD1	2:D:5:SER:HB3	2.44	0.53
1:A:541:LEU:HD11	1:A:579:ARG:HB3	1.91	0.52
1:B:412:SER:HB2	1:B:492:LYS:HG3	1.91	0.52
1:B:400:ALA:HA	1:B:566:LEU:HB3	1.92	0.51
1:B:488:GLU:HB3	1:B:489:HIS:CD2	2.46	0.51
1:A:491:LEU:H	2:C:7:NH2:N	2.08	0.50
1:B:556:LYS:H	1:B:556:LYS:HE2	1.76	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:416:ASP:OD1	1:A:418:SER:OG	2.18	0.50
1:B:541:LEU:HD11	1:B:579:ARG:HB3	1.94	0.49
1:A:469:PRO:HD2	1:A:473:MET:SD	2.53	0.49
1:A:537:ASP:OD2	1:A:579:ARG:NH2	2.47	0.48
1:B:474:LYS:HD2	1:B:474:LYS:H	1.79	0.48
1:A:487:SER:HA	3:A:646:HOH:O	2.13	0.48
1:B:428:CYS:O	1:B:492:LYS:NZ	2.37	0.47
1:B:502:GLY:C	1:B:504:GLU:HB2	2.40	0.47
1:A:503:ASP:OD2	1:A:506:ALA:HB2	2.15	0.46
1:A:419:ASP:OD1	1:A:420:LYS:N	2.49	0.46
1:B:556:LYS:N	1:B:556:LYS:CD	2.79	0.45
1:B:563:ARG:CB	1:B:566:LEU:HD23	2.46	0.45
1:A:408:ILE:CD1	1:A:502:GLY:HA3	2.46	0.45
2:C:7:NH2:N	3:C:101:HOH:O	2.50	0.44
1:B:563:ARG:HB3	1:B:566:LEU:HD23	1.98	0.44
1:A:500:ARG:HG2	1:A:501:GLU:H	1.83	0.44
1:B:415:VAL:HG22	1:B:417:TYR:CD1	2.54	0.43
1:B:534:PHE:CE2	1:B:579:ARG:HG2	2.54	0.43
1:B:505:LEU:O	1:B:505:LEU:HD12	2.19	0.42
1:A:489:HIS:CD2	1:A:489:HIS:N	2.86	0.42
1:B:392:ARG:HH11	1:B:392:ARG:HD3	1.74	0.42
1:A:408:ILE:HD13	1:A:502:GLY:HA3	2.00	0.42
1:A:534:PHE:CE2	1:A:579:ARG:HG2	2.55	0.42
1:B:434:VAL:CG2	1:B:479:LEU:HD13	2.41	0.42
1:B:508:LEU:HD23	1:B:508:LEU:HA	1.81	0.42
1:B:482:PHE:O	1:B:486:MET:HG3	2.20	0.41
1:B:567:LEU:HD22	1:B:572:CYS:HB3	2.02	0.41
1:A:463:LEU:CD1	1:A:469:PRO:HG3	2.49	0.41
1:B:432:VAL:HG23	1:B:479:LEU:HD11	2.02	0.41
1:B:481:TYR:CD1	1:B:481:TYR:C	2.99	0.41
1:B:377:MET:HE3	1:B:377:MET:HB2	1.70	0.40
1:B:501:GLU:O	1:B:502:GLY:C	2.62	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:568:GLU:OE1	1:B:474:LYS:NZ[2_555]	1.90	0.30

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/226 (99%)	220 (99%)	2 (1%)	1 (0%)	30	27
1	B	224/226 (99%)	216 (96%)	7 (3%)	1 (0%)	30	27
2	C	4/7 (57%)	3 (75%)	1 (25%)	0	100	100
2	D	4/7 (57%)	4 (100%)	0	0	100	100
All	All	455/466 (98%)	443 (97%)	10 (2%)	2 (0%)	30	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	504	GLU
1	B	505	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/202 (99%)	197 (98%)	3 (2%)	57	64
1	B	200/202 (99%)	199 (100%)	1 (0%)	81	87
2	C	4/4 (100%)	4 (100%)	0	100	100
2	D	4/4 (100%)	4 (100%)	0	100	100
All	All	408/412 (99%)	404 (99%)	4 (1%)	68	75

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	418	SER
1	A	498	THR
1	A	505	LEU
1	B	501	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	426	GLN
1	A	452	GLN
1	A	489	HIS
1	B	426	GLN
1	B	452	GLN
2	C	4	HIS
2	D	4	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TPO	C	6	2	8,10,11	1.62	1 (12%)	10,14,16	1.13	1 (10%)
2	TPO	D	6	2	8,10,11	1.46	1 (12%)	10,14,16	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	C	6	2	-	1/9/11/13	-
2	TPO	D	6	2	-	1/9/11/13	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	6	TPO	P-OG1	4.15	1.66	1.59
2	D	6	TPO	P-OG1	3.65	1.65	1.59

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	6	TPO	P-OG1-CB	-2.38	116.87	123.33

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	6	TPO	O-C-CA-CB
2	D	6	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	225/226 (99%)	-1.17	0 100 100	25, 37, 56, 88	0
1	B	226/226 (100%)	-1.16	0 100 100	26, 36, 56, 94	0
2	C	4/7 (57%)	-1.23	0 100 100	27, 30, 37, 38	0
2	D	4/7 (57%)	-1.34	0 100 100	26, 30, 36, 39	0
All	All	459/466 (98%)	-1.17	0 100 100	25, 36, 56, 94	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TPO	C	6	11/12	0.99	0.04	26,29,32,34	0
2	TPO	D	6	11/12	1.00	0.03	27,30,32,33	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.