



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 05:28 AM UTC

PDB ID : 6WGB / pdb_00006wgb
Title : Crystal structure of the fab portion of dupilumab
Authors : Druzina, Z.; Atwell, S.; Pustilnik, A.; Antonysamy, S.; Ho, C.; Lieu, R.; Hendle, J.; Benach, J.; Wang, J.
Deposited on : 2020-04-05
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	234	
1	C	234	
2	B	219	
2	D	219	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6179 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dupilumab Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	208	Total	C	N	O	S	0	0	0
			1519	953	256	303	7			
1	C	198	Total	C	N	O	S	0	0	0
			1388	873	234	275	6			

- Molecule 2 is a protein called Dupilumab Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	215	Total	C	N	O	S	0	0	0
			1627	1024	268	329	6			
2	D	208	Total	C	N	O	S	0	0	0
			1528	955	254	313	6			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	38	Total	O	0	0
			38	38		
3	B	57	Total	O	0	0
			57	57		
3	C	8	Total	O	0	0
			8	8		
3	D	14	Total	O	0	0
			14	14		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dupilumab Fab heavy chain

Chain A: 



- Molecule 1: Dupilumab Fab heavy chain

Chain C: 




- Molecule 2: Dupilumab Fab light chain

Chain B: 



- Molecule 2: Dupilumab Fab light chain

Chain D: 





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.75Å 66.29Å 135.48Å 90.00° 91.64° 90.00°	Depositor
Resolution (Å)	30.00 – 1.99 30.00 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.6 (30.00-1.99) 99.6 (30.00-1.99)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.243 , 0.299 0.244 , 0.300	Depositor DCC
R_{free} test set	3098 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 19.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.125 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6179	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	1/1550 (0.1%)	1.28	5/2110 (0.2%)
1	C	0.67	1/1416 (0.1%)	1.29	6/1935 (0.3%)
2	B	0.69	0/1663	1.25	6/2263 (0.3%)
2	D	0.64	0/1559	1.24	8/2124 (0.4%)
All	All	0.68	2/6188 (0.0%)	1.26	25/8432 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	C	0	2
2	B	0	2
2	D	0	3
All	All	0	10

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	176	HIS	CE1-NE2	6.36	1.39	1.32
1	A	176	HIS	CE1-NE2	5.17	1.37	1.32

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	205	THR	CA-CB-OG1	-8.71	96.53	109.60
1	A	195	THR	CA-CB-OG1	-8.18	97.33	109.60
2	B	17	GLU	CB-CA-C	7.21	119.93	109.11
1	A	205	THR	CA-CB-OG1	-7.03	99.05	109.60
1	C	68	PHE	CA-CB-CG	6.67	120.47	113.80
2	D	65	ASP	CA-CB-CG	-6.40	106.20	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	169	THR	CA-CB-OG1	-6.15	100.37	109.60
2	B	1	ASP	CA-CB-CG	6.13	118.73	112.60
2	B	82	ARG	CB-CA-C	6.11	119.54	110.14
2	B	171	GLN	CB-CA-C	6.08	119.38	109.90
2	D	113	ARG	N-CA-CB	-6.01	102.06	111.56
2	B	166	GLU	CB-CA-C	-6.00	97.52	109.33
2	D	148	GLU	CB-CG-CD	5.86	122.55	112.60
2	D	64	PRO	CA-C-N	5.81	132.63	121.54
2	D	64	PRO	C-N-CA	5.81	132.63	121.54
2	B	95	GLN	CB-CA-C	5.75	119.22	109.80
1	C	30	ARG	CA-C-N	5.48	128.17	120.28
1	C	30	ARG	C-N-CA	5.48	128.17	120.28
1	C	128	THR	CB-CA-C	5.42	118.42	109.70
2	D	113	ARG	CB-CG-CD	-5.37	98.96	111.30
1	C	195	THR	CA-CB-OG1	-5.26	101.71	109.60
2	D	14	THR	CB-CA-C	5.26	116.28	108.87
1	A	12	GLU	CB-CG-CD	5.08	121.23	112.60
1	A	29	PHE	CA-C-N	5.04	127.55	120.28
1	A	29	PHE	C-N-CA	5.04	127.55	120.28

There are no chirality outliers.

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	222	ARG	Sidechain
1	A	38	ARG	Sidechain
1	A	72	ARG	Sidechain
2	B	113	ARG	Sidechain
2	B	147	ARG	Sidechain
1	C	19	ARG	Sidechain
1	C	38	ARG	Sidechain
2	D	147	ARG	Sidechain
2	D	216	ARG	Sidechain
2	D	99	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1519	0	1454	9	0
1	C	1388	0	1268	15	0
2	B	1627	0	1548	10	0
2	D	1528	0	1395	20	0
3	A	38	0	0	1	0
3	B	57	0	0	1	0
3	C	8	0	0	0	0
3	D	14	0	0	1	0
All	All	6179	0	5665	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:THR:HG22	3:A:301:HOH:O	1.77	0.83
2:D:140:LEU:HD21	2:D:142:ASN:OD1	1.97	0.65
1:C:98:LYS:O	1:C:112:LEU:HD12	2.00	0.60
1:C:112:LEU:N	2:D:41:TYR:HH	1.99	0.60
2:D:152:GLN:HG3	2:D:159:LEU:CD1	2.33	0.59
2:D:164:SER:HB2	3:D:307:HOH:O	2.04	0.57
2:D:196:VAL:HG22	2:D:215:ASN:HD22	1.70	0.56
2:B:16:GLY:HA2	2:B:82:ARG:HG3	1.88	0.56
1:A:155:LYS:HG3	1:A:189:SER:OG	2.06	0.55
2:B:21:ILE:HD12	2:B:78:LEU:HD23	1.88	0.54
1:C:47:TRP:CZ3	2:D:100:PRO:HB3	2.42	0.54
2:B:38:LEU:HD13	2:B:76:PHE:CD1	2.43	0.54
2:D:217:GLY:O	2:D:218:GLU:HG3	2.07	0.54
2:D:197:TYR:O	2:D:213:SER:HA	2.08	0.53
2:D:153:TRP:CD1	2:D:164:SER:HG	2.25	0.53
2:D:42:LEU:HD21	2:D:44:LYS:HE2	1.90	0.52
1:C:71:SER:O	1:C:79:LEU:HD12	2.10	0.52
2:D:113:ARG:HG3	2:D:176:SER:HB2	1.92	0.50
1:C:6:GLU:OE2	1:C:94:TYR:O	2.29	0.50
2:B:4:MET:HE3	2:B:23:CYS:SG	2.52	0.50
2:D:101:TYR:CD1	2:D:101:TYR:N	2.80	0.49
2:D:40:TRP:HB2	2:D:53:ILE:HB	1.94	0.49
2:D:117:ALA:HB1	2:D:206:LEU:HD13	1.94	0.49
1:C:35:THR:HG23	1:C:99:ASP:OD1	2.14	0.48
1:C:183:GLN:HG3	1:C:187:LEU:O	2.14	0.48
1:A:190:LEU:C	1:A:190:LEU:HD12	2.39	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:152:GLN:HG3	2:D:159:LEU:HD13	1.96	0.47
2:D:195:LYS:O	2:D:215:ASN:HA	2.14	0.47
1:C:5:VAL:O	1:C:22:CYS:HA	2.15	0.47
1:C:34:MET:HB3	1:C:79:LEU:HD22	1.97	0.47
2:B:42:LEU:HD13	2:B:91:TYR:CZ	2.50	0.47
2:B:59:ARG:NH1	3:B:301:HOH:O	2.48	0.47
1:A:87:ARG:O	1:A:90:ASP:HB2	2.15	0.46
1:A:198:SER:O	1:A:201:LEU:HB2	2.16	0.46
1:C:196:VAL:HG11	1:C:206:TYR:CE1	2.51	0.46
2:B:180:LEU:C	2:B:180:LEU:HD23	2.42	0.45
1:A:99:ASP:OD1	1:A:100:ARG:N	2.42	0.45
1:C:190:LEU:HD12	1:C:190:LEU:C	2.41	0.44
1:C:179:PRO:HD2	2:D:167:SER:OG	2.18	0.43
1:A:29:PHE:CD2	1:A:77:ASN:HA	2.53	0.43
2:B:88:VAL:HG23	2:B:109:LEU:O	2.18	0.43
2:D:217:GLY:C	2:D:218:GLU:HG3	2.44	0.42
1:C:201:LEU:HD12	1:C:201:LEU:HA	1.92	0.42
1:A:179:PRO:HD2	2:B:167:SER:OG	2.19	0.42
1:A:196:VAL:HG11	1:A:206:TYR:CE1	2.55	0.41
2:D:7:SER:HA	2:D:8:PRO:HA	1.84	0.41
2:B:117:ALA:HB1	2:B:206:LEU:HD13	2.03	0.41
1:C:40:ALA:O	1:C:41:PRO:C	2.62	0.41
2:D:152:GLN:CG	2:D:159:LEU:HD13	2.50	0.41
1:C:204:LYS:HD2	1:C:206:TYR:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	200/234 (86%)	195 (98%)	4 (2%)	1 (0%)	24 21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	190/234 (81%)	178 (94%)	12 (6%)	0	100	100
2	B	211/219 (96%)	205 (97%)	6 (3%)	0	100	100
2	D	200/219 (91%)	190 (95%)	8 (4%)	2 (1%)	12	8
All	All	801/906 (88%)	768 (96%)	30 (4%)	3 (0%)	30	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	65	ASP
2	D	97	LEU
1	A	65	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/198 (83%)	155 (94%)	10 (6%)	17	14
1	C	139/198 (70%)	128 (92%)	11 (8%)	11	8
2	B	181/193 (94%)	168 (93%)	13 (7%)	13	10
2	D	163/193 (84%)	149 (91%)	14 (9%)	10	6
All	All	648/782 (83%)	600 (93%)	48 (7%)	13	9

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	12	GLU
1	A	100	ARG
1	A	117	GLN
1	A	171	LEU
1	A	184	SER
1	A	190	LEU
1	A	200	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	201	LEU
1	A	213	LYS
2	B	1	ASP
2	B	10	SER
2	B	22	SER
2	B	52	LEU
2	B	65	ASP
2	B	99	THR
2	B	112	LYS
2	B	130	LEU
2	B	131	LYS
2	B	193	LYS
2	B	206	LEU
2	B	208	SER
2	B	215	ASN
1	C	35	THR
1	C	43	LYS
1	C	49	SER
1	C	99	ASP
1	C	112	LEU
1	C	159	PRO
1	C	162	VAL
1	C	201	LEU
1	C	204	LYS
1	C	205	THR
1	C	221	LYS
2	D	3	VAL
2	D	7	SER
2	D	38	LEU
2	D	52	LEU
2	D	65	ASP
2	D	86	GLU
2	D	113	ARG
2	D	130	LEU
2	D	142	ASN
2	D	157	ASN
2	D	164	SER
2	D	185	THR
2	D	206	LEU
2	D	218	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	39	GLN
2	D	43	GLN
2	D	50	GLN
2	D	152	GLN
2	D	215	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	208/234 (88%)	-0.95	0 100 100	16, 29, 44, 59	0
1	C	198/234 (84%)	-0.65	0 100 100	18, 37, 64, 79	0
2	B	215/219 (98%)	-1.01	0 100 100	18, 29, 47, 56	0
2	D	208/219 (94%)	-0.82	0 100 100	18, 33, 61, 82	0
All	All	829/906 (91%)	-0.86	0 100 100	16, 32, 57, 82	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.