



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:08 AM UTC

PDB ID : 6WGJ / pdb_00006wgj
Title : Fab portion of dupilumab with Crystal Kappa design and no interchain disulfide
Authors : Druzina, Z.; Atwell, S.; Pustilnik, A.; Antonysamy, S.; Ho, C.; Lieu, R.; Hendle, J.; Benach, J.; Wang, J.
Deposited on : 2020-04-05
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	234	
1	C	234	
1	E	234	
1	G	234	
2	B	217	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	217	 89%9%..
2	F	217	 88%11%. .
2	H	217	 85%11%.. .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12677 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dupilumab Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	216	Total	C	N	O	S	0	1	0
			1593	1001	268	317	7			
1	C	215	Total	C	N	O	S	0	1	0
			1576	992	265	312	7			
1	E	211	Total	C	N	O	S	0	1	0
			1493	940	251	295	7			
1	G	199	Total	C	N	O	S	0	2	0
			1416	895	236	278	7			

- Molecule 2 is a protein called Dupilumab Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	216	Total	C	N	O	S	0	2	0
			1638	1031	266	334	7			
2	D	215	Total	C	N	O	S	0	2	0
			1617	1020	261	329	7			
2	F	213	Total	C	N	O	S	0	1	0
			1541	969	250	315	7			
2	H	210	Total	C	N	O	S	0	1	0
			1520	960	245	308	7			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	44	Total	O	0	0
			44	44		
3	B	53	Total	O	0	0
			53	53		
3	C	28	Total	O	0	0
			28	28		
3	D	25	Total	O	0	0
			25	25		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	22	Total 22	O 22	0	0
3	F	40	Total 40	O 40	0	0
3	G	28	Total 28	O 28	0	0
3	H	43	Total 43	O 43	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dupilumab Fab heavy chain

Chain A: 




- Molecule 1: Dupilumab Fab heavy chain

Chain C: 




- Molecule 1: Dupilumab Fab heavy chain

Chain E: 




- Molecule 1: Dupilumab Fab heavy chain

Chain G: 




- Molecule 2: Dupilumab Fab light chain

Chain B:  92% 6% •



• Molecule 2: Dupilumab Fab light chain

Chain D:  89% 9% ••




• Molecule 2: Dupilumab Fab light chain

Chain F:  88% 11% •



• Molecule 2: Dupilumab Fab light chain

Chain H:  85% 11% ••



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	108.90Å 78.65Å 109.30Å 90.00° 91.40° 90.00°	Depositor
Resolution (Å)	30.00 – 1.90 30.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.8 (30.00-1.90) 95.8 (30.00-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.235 , 0.289 0.236 , 0.289	Depositor DCC
R_{free} test set	6710 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	29.0	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 24.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.054 for l,k,-h 0.058 for h,-k,-l 0.229 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12677	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 44.35 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5453e-04.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/1631	1.15	2/2221 (0.1%)
1	C	0.61	0/1612	1.11	2/2195 (0.1%)
1	E	0.60	0/1530	1.15	3/2095 (0.1%)
1	G	0.61	0/1454	1.15	5/1988 (0.3%)
2	B	0.60	0/1679	1.11	1/2287 (0.0%)
2	D	0.57	0/1658	1.12	2/2259 (0.1%)
2	F	0.61	0/1577	1.17	3/2155 (0.1%)
2	H	0.62	0/1556	1.13	1/2128 (0.0%)
All	All	0.61	0/12697	1.14	19/17328 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	E	0	3
1	G	0	1
2	B	0	2
2	D	0	2
All	All	0	9

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	PHE	CA-CB-CG	6.51	120.31	113.80
1	G	99	ASP	CA-CB-CG	6.30	118.90	112.60
1	C	68	PHE	CA-CB-CG	5.91	119.71	113.80
2	D	27	GLN	CB-CA-C	-5.77	99.89	110.62
1	G	68	PHE	CA-CB-CG	5.62	119.42	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	118	PRO	CB-CA-C	-5.61	103.63	110.98
2	F	39	ASP	CA-CB-CG	5.46	118.06	112.60
2	B	127	ASP	CB-CA-C	5.43	119.81	110.79
1	E	220	ASP	CA-CB-CG	5.41	118.01	112.60
1	E	99	ASP	CA-CB-CG	5.38	117.97	112.60
1	G	161	PRO	N-CA-CB	-5.37	96.70	102.60
1	G	172	THR	CA-CB-OG1	-5.33	101.61	109.60
1	C	176	HIS	CA-CB-CG	5.12	118.92	113.80
2	F	6	GLN	CB-CA-C	5.10	118.56	109.38
2	F	59	ARG	CB-CA-C	5.08	118.13	109.80
1	E	89	GLU	CB-CA-C	-5.07	100.43	110.27
1	A	31	ASP	CA-CB-CG	5.02	117.62	112.60
1	G	77	ASN	CB-CA-C	-5.01	104.49	111.80
2	H	202	THR	CA-CB-OG1	-5.01	102.09	109.60

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	24	ARG	Sidechain
2	B	66	ARG	Sidechain
1	C	100	ARG	Sidechain
2	D	147	ARG	Sidechain
2	D	59	ARG	Sidechain
1	E	100	ARG	Sidechain
1	E	19	ARG	Sidechain
1	E	72	ARG	Sidechain
1	G	87	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1593	0	1523	6	0
1	C	1576	0	1498	7	1
1	E	1493	0	1347	4	0
1	G	1416	0	1269	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1638	0	1561	5	0
2	D	1617	0	1536	7	0
2	F	1541	0	1390	10	0
2	H	1520	0	1374	11	1
3	A	44	0	0	0	0
3	B	53	0	0	0	0
3	C	28	0	0	0	0
3	D	25	0	0	0	0
3	E	22	0	0	0	0
3	F	40	0	0	0	0
3	G	28	0	0	0	0
3	H	43	0	0	0	0
All	All	12677	0	11498	56	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:62:ASP:OD1	1:C:65:LYS:NZ	2.19	0.70
2:D:42:LEU:HD21	2:D:44:LYS:HE3	1.80	0.64
1:C:13:GLN:OE1	1:C:125:SER:HB3	1.99	0.63
1:G:6:GLU:HG3	1:G:118:GLY:H	1.67	0.59
2:F:12:PRO:HB2	2:F:112:LYS:HE3	1.85	0.59
2:B:4:MET:HE3	2:B:23:CYS:SG	2.45	0.56
2:D:59:ARG:NH1	2:D:67:PHE:O	2.38	0.56
1:C:162:VAL:HG11	1:C:210:VAL:HG13	1.86	0.56
2:F:11:LEU:HB3	2:F:109:LEU:HD12	1.87	0.55
1:G:210:VAL:O	1:G:218:LYS:HA	2.07	0.54
1:C:2:VAL:HG12	1:C:114:VAL:HG11	1.89	0.54
2:H:4:MET:HE3	2:H:23:CYS:SG	2.49	0.53
1:A:71:SER:OG	1:A:80:TYR:HB2	2.09	0.53
1:C:6:GLU:OE2	1:C:116:GLY:HA3	2.12	0.50
1:C:34:MET:HB3	1:C:79:LEU:HD22	1.94	0.49
2:B:171:GLN:HG3	2:B:178:TYR:CZ	2.47	0.49
2:H:11:LEU:HD22	2:H:13:VAL:HG22	1.95	0.49
1:A:6:GLU:OE1	1:A:116:GLY:HA3	2.12	0.48
2:F:141:LEU:HD13	2:F:180:LEU:HD22	1.98	0.45
2:B:141:LEU:HD12	2:B:141:LEU:N	2.31	0.45
1:A:11:LEU:HB2	1:A:159:PRO:HG3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:85:ALA:HA	2:F:111:ILE:HG13	1.98	0.45
1:C:98:LYS:NZ	1:C:113:ASP:OD2	2.47	0.45
2:H:99:THR:HA	2:H:100:PRO:C	2.41	0.44
2:H:141:LEU:HD12	2:H:141:LEU:N	2.32	0.44
2:D:171:GLN:HG3	2:D:178:TYR:CZ	2.52	0.44
2:H:66:ARG:HD2	2:H:82:ARG:O	2.17	0.44
2:D:66:ARG:O	2:D:80:ILE:HA	2.17	0.44
2:D:125:PRO:HG3	2:D:135:ALA:HB1	1.98	0.44
2:H:180:LEU:C	2:H:180:LEU:HD23	2.42	0.44
1:G:68:PHE:CZ	1:G:83:MET:HE3	2.53	0.43
2:B:53:ILE:HD13	2:B:59:ARG:HA	2.00	0.43
2:F:4:MET:HE3	2:F:23:CYS:SG	2.58	0.43
2:H:15:PRO:HD3	2:H:111:ILE:HG23	2.00	0.43
2:H:36:ASN:O	2:H:55:LEU:HA	2.19	0.43
2:B:42:LEU:O	2:B:50:GLN:HG2	2.19	0.43
2:F:88:VAL:HG13	2:F:111:ILE:HG12	2.01	0.42
1:G:151:GLY:HA2	1:G:166:TRP:CZ2	2.54	0.42
1:A:34:MET:HB3	1:A:79:LEU:HD22	2.02	0.42
2:D:4:MET:HE3	2:D:23:CYS:SG	2.58	0.42
1:E:6:GLU:OE1	1:E:116:GLY:HA3	2.19	0.42
2:H:125:PRO:HD3	2:H:137:VAL:HG22	2.00	0.42
1:E:162:VAL:HG23	1:E:212:HIS:CD2	2.55	0.42
1:A:138:PRO:HD3	1:A:150:LEU:HD23	2.02	0.41
1:G:166:TRP:CH2	1:G:208:CYS:HB3	2.55	0.41
2:D:11:LEU:CD2	2:D:13:VAL:HG22	2.51	0.41
2:H:188:LYS:O	2:H:192:GLU:HG3	2.21	0.41
1:E:158:PHE:HA	1:E:159:PRO:HA	1.88	0.41
1:E:68:PHE:CZ	1:E:83:MET:HE3	2.55	0.41
2:F:153:TRP:HB2	2:F:160:GLN:HB2	2.03	0.41
1:G:6:GLU:OE1	1:G:116:GLY:HA3	2.20	0.41
1:A:101:LEU:O	1:A:107:PRO:HB3	2.21	0.41
2:F:180:LEU:HD23	2:F:180:LEU:C	2.46	0.41
2:F:40:TRP:CZ3	2:F:93:CYS:HB3	2.57	0.40
2:F:88:VAL:CG1	2:F:111:ILE:HG12	2.51	0.40
2:H:119:SER:HB2	2:H:142:ASN:HB3	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:69:THR:OG1	2:H:20:SER:OG[1_455]	2.16	0.04

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/234 (90%)	205 (97%)	6 (3%)	0	100	100
1	C	208/234 (89%)	202 (97%)	6 (3%)	0	100	100
1	E	206/234 (88%)	197 (96%)	7 (3%)	2 (1%)	12	5
1	G	193/234 (82%)	184 (95%)	7 (4%)	2 (1%)	12	5
2	B	216/217 (100%)	211 (98%)	4 (2%)	1 (0%)	24	16
2	D	215/217 (99%)	210 (98%)	5 (2%)	0	100	100
2	F	210/217 (97%)	203 (97%)	7 (3%)	0	100	100
2	H	207/217 (95%)	202 (98%)	5 (2%)	0	100	100
All	All	1666/1804 (92%)	1614 (97%)	47 (3%)	5 (0%)	36	29

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	105	ILE
1	E	197	PRO
2	B	65	ASP
1	G	185	SER
1	G	161	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/197 (88%)	168 (97%)	5 (3%)	37	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	169/197 (86%)	166 (98%)	3 (2%)	51	50
1	E	148/197 (75%)	144 (97%)	4 (3%)	39	34
1	G	138/197 (70%)	134 (97%)	4 (3%)	37	31
2	B	183/191 (96%)	178 (97%)	5 (3%)	39	34
2	D	179/191 (94%)	173 (97%)	6 (3%)	32	25
2	F	159/191 (83%)	154 (97%)	5 (3%)	35	29
2	H	156/191 (82%)	151 (97%)	5 (3%)	34	27
All	All	1305/1552 (84%)	1268 (97%)	37 (3%)	37	32

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	GLU
1	A	146	SER
1	A	150	LEU
1	A	162	VAL
1	A	218	LYS
2	B	11	LEU
2	B	20	SER
2	B	24	ARG
2	B	33	ILE
2	B	65	ASP
1	C	104	THR
1	C	108	ARG
1	C	218	LYS
2	D	11	LEU
2	D	22	SER
2	D	33	ILE
2	D	65	ASP
2	D	75	ASP
2	D	84	GLU
1	E	6	GLU
1	E	162	VAL
1	E	173	SER
1	E	204	LYS
2	F	13	VAL
2	F	65	ASP
2	F	75	ASP
2	F	130	LEU
2	F	194	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	6	GLU
1	G	159	PRO
1	G	161	PRO
1	G	162	VAL
2	H	11	LEU
2	H	13	VAL
2	H	38	LEU
2	H	50	GLN
2	H	84	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
2	B	27	GLN
2	B	142	ASN
1	C	82	GLN
2	F	98	GLN
2	F	142	ASN
2	H	50	GLN
2	H	165	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	216/234 (92%)	-0.90	0 100 100	14, 26, 45, 62	1 (0%)
1	C	215/234 (91%)	-0.83	0 100 100	17, 29, 50, 69	1 (0%)
1	E	211/234 (90%)	-0.62	1 (0%) 87 89	20, 34, 50, 99	1 (0%)
1	G	199/234 (85%)	-0.73	0 100 100	19, 30, 49, 64	2 (1%)
2	B	216/217 (99%)	-0.89	0 100 100	16, 28, 48, 66	2 (0%)
2	D	215/217 (99%)	-0.73	1 (0%) 87 89	17, 34, 51, 71	2 (0%)
2	F	213/217 (98%)	-0.72	0 100 100	19, 31, 54, 86	1 (0%)
2	H	210/217 (96%)	-0.84	0 100 100	16, 29, 49, 67	1 (0%)
All	All	1695/1804 (93%)	-0.78	2 (0%) 92 92	14, 30, 51, 99	11 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	104	THR	2.4
2	D	215	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.