



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 12:47 PM UTC

PDB ID : 8WG1 / pdb_00008wg1
Title : Crystal structure of GH97 glucodextranase mutant E509Q from *Flavobacterium johnsoniae* in complex with panose
Authors : Kurata, R.; Nakamura, S.; Miyazaki, T.
Deposited on : 2023-09-20
Resolution : 2.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	685	87% 12% .
1	B	685	88% 11% .
1	C	685	87% 12% .
1	D	685	88% 11% .
2	E	3	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	3	 100%
2	G	3	 67%  33%
2	H	3	 100%

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 22517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Candidate alpha-glucosidase Glycoside hydrolase family 97.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	682	Total	C	N	O	S	0	1	0
			5476	3490	919	1048	19			
1	B	682	Total	C	N	O	S	0	0	0
			5471	3487	918	1047	19			
1	C	682	Total	C	N	O	S	0	1	0
			5476	3490	919	1048	19			
1	D	685	Total	C	N	O	S	0	1	0
			5495	3501	921	1054	19			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	SER	-	expression tag	UNP A5FBI0
A	509	GLN	GLU	engineered mutation	UNP A5FBI0
B	20	SER	-	expression tag	UNP A5FBI0
B	509	GLN	GLU	engineered mutation	UNP A5FBI0
C	20	SER	-	expression tag	UNP A5FBI0
C	509	GLN	GLU	engineered mutation	UNP A5FBI0
D	20	SER	-	expression tag	UNP A5FBI0
D	509	GLN	GLU	engineered mutation	UNP A5FBI0

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	E	3	Total	C	O	0	0	0
			34	18	16			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	F	3	Total	C	O	0	0	0
			34	18	16			
2	G	3	Total	C	O	0	0	0
			34	18	16			
2	H	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	B	1	Total	Ca	0	0
			1	1		
3	C	1	Total	Ca	0	0
			1	1		
3	D	1	Total	Ca	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	C	2	Total	Mg	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	130	Total	O	0	0
			130	130		
6	B	89	Total	O	0	0
			89	89		
6	C	121	Total	O	0	0
			121	121		
6	D	92	Total	O	0	0
			92	92		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Candidate alpha-glucosidase Glycoside hydrolase family 97



- Molecule 1: Candidate alpha-glucosidase Glycoside hydrolase family 97





- Molecule 1: Candidate alpha-glucosidase Glycoside hydrolase family 97





- Molecule 1: Candidate alpha-glucosidase Glycoside hydrolase family 97




- Molecule 2: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranos e



- Molecule 2: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranos e



- Molecule 2: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranos e



- Molecule 2: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranos e



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	290.69Å 102.71Å 104.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.23 – 2.34 49.23 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.23-2.34) 99.9 (49.23-2.34)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.208 , 0.254 0.213 , 0.259	Depositor DCC
R_{free} test set	6575 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 18.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.006 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	22517	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CA, GLC, EDO, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/5623	1.31	19/7626 (0.2%)
1	B	0.81	2/5615 (0.0%)	1.31	28/7615 (0.4%)
1	C	0.85	1/5623 (0.0%)	1.33	27/7626 (0.4%)
1	D	0.81	0/5642	1.34	26/7652 (0.3%)
All	All	0.83	3/22503 (0.0%)	1.32	100/30519 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	3
1	C	0	4
1	D	0	4
All	All	0	15

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	427	HIS	CE1-NE2	7.93	1.40	1.32
1	B	414	HIS	ND1-CE1	6.10	1.38	1.32
1	B	458	HIS	CE1-NE2	5.49	1.38	1.32

The worst 5 of 100 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	579	GLU	CB-CA-C	-12.27	89.82	110.68
1	B	579	GLU	CB-CA-C	-11.46	91.20	110.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	702	ASP	CA-CB-CG	9.11	121.71	112.60
1	C	579	GLU	CB-CA-C	-8.84	96.12	110.79
1	D	214	GLN	N-CA-CB	8.77	123.72	110.22

There are no chirality outliers.

5 of 15 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	115	ARG	Sidechain
1	A	267	ARG	Sidechain
1	A	341	ARG	Sidechain
1	A	602	ARG	Sidechain
1	B	115	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5476	0	5259	40	0
1	B	5471	0	5253	31	0
1	C	5476	0	5259	29	0
1	D	5495	0	5273	30	0
2	E	34	0	27	0	0
2	F	34	0	27	0	0
2	G	34	0	29	1	0
2	H	34	0	28	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	8	0	12	2	0
4	B	4	0	6	0	0
4	C	8	0	12	0	0
4	D	4	0	6	0	0
5	A	1	0	0	0	0
5	C	2	0	0	0	0
6	A	130	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	89	0	0	4	0
6	C	121	0	0	3	0
6	D	92	0	0	3	0
All	All	22517	0	21191	125	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 125 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:674:LYS:HB3	6:D:991:HOH:O	1.46	1.14
1:B:583:ARG:NH1	6:B:901:HOH:O	2.06	0.82
1:B:583:ARG:NE	6:B:901:HOH:O	1.91	0.80
1:C:640:ASP:OD2	1:C:682:LYS:HE2	1.88	0.73
1:B:114:ASP:OD2	1:C:673:ARG:NH1	2.27	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	681/685 (99%)	641 (94%)	38 (6%)	2 (0%)	36	41
1	B	680/685 (99%)	645 (95%)	33 (5%)	2 (0%)	36	41
1	C	681/685 (99%)	645 (95%)	34 (5%)	2 (0%)	36	41
1	D	684/685 (100%)	649 (95%)	34 (5%)	1 (0%)	48	57
All	All	2726/2740 (100%)	2580 (95%)	139 (5%)	7 (0%)	36	41

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	308	TRP
1	A	308	TRP
1	B	503	GLU
1	A	141	ILE
1	B	141	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	582/582 (100%)	572 (98%)	10 (2%)	53	66
1	B	581/582 (100%)	576 (99%)	5 (1%)	70	80
1	C	582/582 (100%)	576 (99%)	6 (1%)	68	79
1	D	583/582 (100%)	578 (99%)	5 (1%)	70	80
All	All	2328/2328 (100%)	2302 (99%)	26 (1%)	65	77

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	682	LYS
1	C	207	THR
1	D	579	GLU
1	C	72	ASP
1	C	602	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	423	ASN
1	D	64	ASN
1	C	502	ASN
1	D	66	ASN
1	B	100	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BGC	E	1	2	12,12,12	0.97	1 (8%)	17,17,17	3.13	8 (47%)
2	GLC	E	2	3,2	11,11,12	0.47	0	15,15,17	1.95	5 (33%)
2	GLC	E	3	3,2	11,11,12	0.53	0	15,15,17	1.67	2 (13%)
2	BGC	F	1	2	12,12,12	0.83	0	17,17,17	1.69	5 (29%)
2	GLC	F	2	3,2	11,11,12	0.59	0	15,15,17	1.53	2 (13%)
2	GLC	F	3	3,2	11,11,12	0.31	0	15,15,17	2.02	3 (20%)
2	BGC	G	1	2	12,12,12	1.10	2 (16%)	17,17,17	1.54	5 (29%)
2	GLC	G	2	3,2	11,11,12	0.86	0	15,15,17	1.51	2 (13%)
2	GLC	G	3	3,2	11,11,12	0.84	0	15,15,17	2.33	7 (46%)
2	BGC	H	1	2	12,12,12	0.96	0	17,17,17	1.93	4 (23%)
2	GLC	H	2	3,2	11,11,12	0.51	0	15,15,17	2.35	3 (20%)
2	GLC	H	3	3,2	11,11,12	0.67	0	15,15,17	2.01	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	E	1	2	-	2/2/22/22	0/1/1/1
2	GLC	E	2	3,2	-	2/2/19/22	0/1/1/1
2	GLC	E	3	3,2	-	0/2/19/22	0/1/1/1
2	BGC	F	1	2	-	0/2/22/22	0/1/1/1
2	GLC	F	2	3,2	-	2/2/19/22	0/1/1/1
2	GLC	F	3	3,2	-	0/2/19/22	0/1/1/1
2	BGC	G	1	2	-	2/2/22/22	0/1/1/1
2	GLC	G	2	3,2	-	1/2/19/22	0/1/1/1
2	GLC	G	3	3,2	-	0/2/19/22	0/1/1/1
2	BGC	H	1	2	-	2/2/22/22	0/1/1/1
2	GLC	H	2	3,2	-	1/2/19/22	0/1/1/1
2	GLC	H	3	3,2	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	BGC	O1-C1	-2.46	1.31	1.39
2	G	1	BGC	C1-C2	2.04	1.57	1.52
2	G	1	BGC	O4-C4	2.01	1.47	1.43

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	2	GLC	C1-O5-C5	7.20	121.83	112.19
2	E	1	BGC	O5-C1-C2	6.98	122.58	110.30
2	F	3	GLC	C1-O5-C5	5.73	119.86	112.19
2	H	3	GLC	C2-C3-C4	-5.48	101.22	110.86
2	E	1	BGC	O1-C1-C2	-4.77	95.13	108.98

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	2	GLC	O5-C5-C6-O6
2	E	1	BGC	C4-C5-C6-O6
2	F	2	GLC	O5-C5-C6-O6
2	E	1	BGC	O5-C5-C6-O6
2	E	2	GLC	C4-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	3	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 7 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	B	802	-	3,3,3	0.35	0	2,2,2	0.67	0
4	EDO	C	802	-	3,3,3	1.34	1 (33%)	2,2,2	0.70	0
4	EDO	A	802	-	3,3,3	0.73	0	2,2,2	1.06	0
4	EDO	D	802	-	3,3,3	1.93	1 (33%)	2,2,2	0.94	0
4	EDO	A	803	-	3,3,3	0.78	0	2,2,2	0.87	0
4	EDO	C	803	-	3,3,3	1.76	2 (66%)	2,2,2	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	802	-	-	0/1/1/1	-
4	EDO	C	802	-	-	0/1/1/1	-
4	EDO	A	802	-	-	1/1/1/1	-
4	EDO	D	802	-	-	1/1/1/1	-
4	EDO	A	803	-	-	1/1/1/1	-
4	EDO	C	803	-	-	0/1/1/1	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	802	EDO	O1-C1	2.76	1.56	1.42
4	C	803	EDO	O1-C1	2.13	1.52	1.42
4	C	802	EDO	O1-C1	2.03	1.52	1.42
4	C	803	EDO	O2-C2	2.01	1.52	1.42

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	802	EDO	O1-C1-C2-O2
4	A	803	EDO	O1-C1-C2-O2
4	D	802	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	802	EDO	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	682/685 (99%)	-0.03	6 (0%) 81 84	17, 31, 48, 72	1 (0%)
1	B	682/685 (99%)	0.12	9 (1%) 75 79	23, 35, 53, 84	0
1	C	682/685 (99%)	0.00	6 (0%) 81 84	15, 32, 49, 71	1 (0%)
1	D	685/685 (100%)	0.10	9 (1%) 75 79	23, 34, 52, 75	1 (0%)
All	All	2731/2740 (99%)	0.05	30 (1%) 78 81	15, 33, 51, 84	3 (0%)

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	683	LEU	7.2
1	D	704	ALA	4.5
1	C	701	ALA	4.2
1	B	701	ALA	3.6
1	B	20	SER	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	E	1	12/12	0.82	0.11	29,39,43,43	0
2	BGC	G	1	12/12	0.85	0.12	41,52,59,60	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BGC	F	1	12/12	0.87	0.11	35,46,50,53	0
2	BGC	H	1	12/12	0.88	0.12	32,43,49,55	0
2	GLC	G	3	11/12	0.94	0.07	24,28,33,36	0
2	GLC	H	2	11/12	0.94	0.07	27,29,31,33	0
2	GLC	F	2	11/12	0.95	0.07	23,28,30,32	0
2	GLC	E	3	11/12	0.95	0.07	24,26,28,29	0
2	GLC	G	2	11/12	0.95	0.07	30,38,43,44	0
2	GLC	F	3	11/12	0.96	0.06	25,31,33,34	0
2	GLC	H	3	11/12	0.96	0.06	23,30,33,34	0
2	GLC	E	2	11/12	0.97	0.05	22,26,30,33	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain G:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	C	803	4/4	0.59	0.29	41,43,53,54	0
4	EDO	A	803	4/4	0.80	0.18	45,47,49,54	0
4	EDO	D	802	4/4	0.80	0.16	27,30,31,35	0
4	EDO	C	802	4/4	0.84	0.12	27,27,27,31	0
4	EDO	A	802	4/4	0.93	0.09	28,29,29,30	0
4	EDO	B	802	4/4	0.95	0.08	32,36,38,41	0
3	CA	C	801	1/1	0.95	0.15	45,45,45,45	0
5	MG	C	804	1/1	0.95	0.04	30,30,30,30	0
3	CA	A	801	1/1	0.97	0.10	37,37,37,37	0
5	MG	C	805	1/1	0.97	0.06	44,44,44,44	0
5	MG	A	804	1/1	0.98	0.04	27,27,27,27	0
3	CA	B	801	1/1	0.98	0.08	46,46,46,46	0
3	CA	D	801	1/1	0.98	0.04	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.