



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 09:12 PM UTC

PDB ID : 7WH5 / pdb_00007wh5
Title : Crystal structure of human ClpP in complex with ZG180
Authors : Wei, B.Y.; Gan, J.H.; Yang, C.-G.
Deposited on : 2021-12-29
Resolution : 2.13 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	6% 73% • 23%
1	B	221	9% 67% 10% 22%
1	C	221	6% 70% 6% • 23%
1	D	221	5% 71% 7% 23%
1	E	221	10% 74% • 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	221	
1	G	221	
1	H	221	
1	I	221	
1	J	221	
1	K	221	
1	L	221	
1	M	221	
1	N	221	

2 Entry composition


There are 4 unique types of molecules in this entry. The entry contains 19072 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent Clp protease proteolytic subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	170	Total	C	N	O	S	0	0	0
			1306	836	221	236	13			
1	B	172	Total	C	N	O	S	0	0	0
			1293	830	215	236	12			
1	C	170	Total	C	N	O	S	0	0	0
			1302	834	217	238	13			
1	D	171	Total	C	N	O	S	0	0	0
			1316	842	221	240	13			
1	E	172	Total	C	N	O	S	0	0	0
			1322	846	223	240	13			
1	F	173	Total	C	N	O	S	0	0	0
			1301	835	215	238	13			
1	G	171	Total	C	N	O	S	0	0	0
			1303	835	219	236	13			
1	H	172	Total	C	N	O	S	0	0	0
			1322	848	221	240	13			
1	I	177	Total	C	N	O	S	0	0	0
			1343	862	227	242	12			
1	J	171	Total	C	N	O	S	0	0	0
			1300	834	220	234	12			
1	K	171	Total	C	N	O	S	0	0	0
			1280	821	212	235	12			
1	L	170	Total	C	N	O	S	0	0	0
			1295	829	218	235	13			
1	M	169	Total	C	N	O	S	0	0	0
			1305	834	222	236	13			
1	N	170	Total	C	N	O	S	0	0	0
			1311	838	221	239	13			

- Molecule 2 is (6S,9aS)-6-[(2S)-butan-2-yl]-8-(naphthalen-1-ylmethyl)-4,7-bis(oxidanylidene)-N-[4,4,4-tris(fluoranyl)butyl]-3,6,9,9a-tetrahydro-2H-pyrazino[1,2-a]pyrimidine-1-carboxamide (CCD ID: 9DF) (formula: C₂₇H₃₃F₃N₄O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	A	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	C	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	C	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	D	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	E	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	F	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	H	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	H	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	I	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	J	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	K	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	L	1	Total	C	F	N	O	0	0
			37	27	3	4	3		
2	N	1	Total	C	F	N	O	0	0
			37	27	3	4	3		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	N	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	14	Total 14	O 14	0	0
4	B	14	Total 14	O 14	0	0
4	C	23	Total 23	O 23	0	0
4	D	36	Total 36	O 36	0	0
4	E	9	Total 9	O 9	0	0
4	F	13	Total 13	O 13	0	0
4	G	13	Total 13	O 13	0	0
4	H	16	Total 16	O 16	0	0
4	I	8	Total 8	O 8	0	0
4	J	5	Total 5	O 5	0	0
4	K	8	Total 8	O 8	0	0
4	L	19	Total 19	O 19	0	0
4	M	37	Total 37	O 37	0	0
4	N	39	Total 39	O 39	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP-dependent Clp protease proteolytic subunit, mitochondrial




- Molecule 1: ATP-dependent Clp protease proteolytic subunit, mitochondrial



- Molecule 1: ATP-dependent Clp protease proteolytic subunit, mitochondrial



- Molecule 1: ATP-dependent Clp protease proteolytic subunit, mitochondrial



GLU ASP GLU PRO THR LEU VAL GLN LYS GLU PRO VAL GLU ALA ALA PRO PRO VAL PRO VAL PRO SER THR

- Molecule 1: ATP-dependent Clp protease proteolytic subunit, mitochondrial

Chain E: 10% 74% 22%

PRO L58 L59 P60 I61 V62 V63 GLU GLN THR GLY ARG GLY ARG A72 Y73 D74 I75 M88 L98 L103 L104 F105 L106 Q107 M117 L140 P180 SER GLY GLY ALA ARG GLY GLN ALA THR ASP ILE I193 L217 S222 M233 Q236 D242 L245

- Molecule 1: ATP-dependent Clp protease proteolytic subunit, mitochondrial

Chain F: 10% 71% 6% 22%

PRO L58 L59 V62 V63 GLU GLN THR GLY ARG GLY R71 A72 M88 L98 Q107 Y118 L140 V148 P164 P180 SER GLY GLY ALA ARG GLY GLN THR ASP ILE I193 Q194 A195 I198 A210 R214 L217 M224 M233 Q236

- Molecule 1: ATP-dependent Clp protease proteolytic subunit, mitochondrial

Chain G: 6% 72% 23%

PRO L58 L59 P60 V62 V63 GLU GLN THR GLY ARG GLU A72 Y73 L79 T84 M88 Q107 L140 P180 SER GLY GLY ALA ARG GLY GLN THR ASP ILE I193 N207 L217 M230 A235 T240 K243 V244 L245 P249 GLN ASP


- Molecule 1: ATP-dependent Clp protease proteolytic subunit, mitochondrial


Chain H: 4% 71% 6% 22%


PRO L58 V62 V63 GLN THR GLY ARG GLU A72 Y73 D74 S77 E82 M88 L98 L140 H178 Q179 P180 SER GLY GLY ALA ARG GLY GLN THR ASP ILE I193 Q194 A195 I198 K203 M230 Q236 T240 V244 D242 P249 GLN


- Molecule 1: ATP-dependent Clp protease proteolytic subunit, mitochondrial


Chain I: 7% 76% 20%




- Chain J:  7% 75% 23%





- Chain K:  9% 73% 23%




- Chain L:



- Chain M:  5% 73% 24%



- 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	111.89Å 97.03Å 131.42Å 90.00° 93.70° 90.00°	Depositor
Resolution (Å)	54.33 – 2.13 54.33 – 2.13	Depositor EDS
% Data completeness (in resolution range)	99.4 (54.33-2.13) 99.7 (54.33-2.13)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.20 (at 2.12Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.230 , 0.249 0.242 , 0.257	Depositor DCC
R_{free} test set	7770 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	42.3	Xtriage
Anisotropy	0.202	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	19072	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 9DF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	0/1331	0.86	1/1805 (0.1%)
1	B	0.82	1/1317 (0.1%)	0.89	2/1791 (0.1%)
1	C	0.75	0/1327	0.87	1/1799 (0.1%)
1	D	0.89	0/1341	0.92	1/1818 (0.1%)
1	E	0.76	0/1347	0.89	1/1827 (0.1%)
1	F	0.74	0/1326	0.88	0/1803
1	G	0.74	1/1328 (0.1%)	0.89	3/1801 (0.2%)
1	H	0.74	0/1347	0.88	1/1827 (0.1%)
1	I	0.80	1/1369 (0.1%)	0.88	1/1858 (0.1%)
1	J	0.73	0/1326	0.87	1/1800 (0.1%)
1	K	0.77	1/1306 (0.1%)	0.85	1/1776 (0.1%)
1	L	0.75	0/1319	0.84	0/1788
1	M	0.78	0/1329	0.88	0/1800
1	N	0.84	0/1335	0.92	0/1807
All	All	0.78	4/18648 (0.0%)	0.88	13/25300 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	63	VAL	CA-C	5.93	1.59	1.52
1	G	207	ASN	CG-ND2	5.62	1.45	1.33
1	K	177	ILE	C-O	-5.39	1.19	1.24
1	I	63	VAL	CA-C	5.30	1.59	1.52

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	75	ILE	N-CA-C	8.02	119.77	110.62
1	G	62	VAL	N-CA-C	7.70	118.52	108.35
1	D	75	ILE	N-CA-C	7.32	118.97	110.62

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	75	ILE	N-CA-C	7.18	118.81	110.62
1	B	63	VAL	N-CA-C	7.00	119.14	108.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1306	0	1316	6	0
1	B	1293	0	1278	17	0
1	C	1302	0	1309	12	0
1	D	1316	0	1327	11	0
1	E	1322	0	1334	3	0
1	F	1301	0	1287	18	0
1	G	1303	0	1309	7	0
1	H	1322	0	1341	7	0
1	I	1343	0	1353	7	0
1	J	1300	0	1301	6	0
1	K	1280	0	1255	7	0
1	L	1295	0	1303	6	0
1	M	1305	0	1322	6	0
1	N	1311	0	1330	4	0
2	A	74	0	0	1	0
2	C	74	0	0	1	0
2	D	37	0	0	0	0
2	E	37	0	0	0	0
2	F	37	0	0	0	0
2	H	74	0	0	1	0
2	I	37	0	0	0	0
2	J	37	0	0	0	0
2	K	37	0	0	1	0
2	L	37	0	0	1	0
2	N	37	0	0	0	0
3	N	1	0	0	0	0
4	A	14	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	14	0	0	2	0
4	C	23	0	0	0	0
4	D	36	0	0	1	0
4	E	9	0	0	0	0
4	F	13	0	0	0	0
4	G	13	0	0	2	0
4	H	16	0	0	0	0
4	I	8	0	0	0	0
4	J	5	0	0	0	0
4	K	8	0	0	0	0
4	L	19	0	0	0	0
4	M	37	0	0	0	0
4	N	39	0	0	0	0
All	All	19072	0	18365	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 102 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:198:ILE:HG21	1:J:198:ILE:HD11	1.29	1.10
1:F:198:ILE:HG21	1:J:198:ILE:CD1	1.89	1.02
1:B:118:TYR:CE2	1:B:148:VAL:HG21	2.14	0.82
1:A:94:SER:O	1:A:98:LEU:HD23	1.85	0.76
1:F:198:ILE:CG2	1:J:198:ILE:HD11	2.16	0.72

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/221 (74%)	162 (99%)	2 (1%)	0	100	100
1	B	166/221 (75%)	164 (99%)	2 (1%)	0	100	100
1	C	164/221 (74%)	162 (99%)	2 (1%)	0	100	100
1	D	165/221 (75%)	162 (98%)	3 (2%)	0	100	100
1	E	166/221 (75%)	164 (99%)	2 (1%)	0	100	100
1	F	167/221 (76%)	165 (99%)	2 (1%)	0	100	100
1	G	165/221 (75%)	163 (99%)	2 (1%)	0	100	100
1	H	166/221 (75%)	164 (99%)	2 (1%)	0	100	100
1	I	171/221 (77%)	169 (99%)	2 (1%)	0	100	100
1	J	165/221 (75%)	163 (99%)	2 (1%)	0	100	100
1	K	165/221 (75%)	163 (99%)	2 (1%)	0	100	100
1	L	164/221 (74%)	162 (99%)	2 (1%)	0	100	100
1	M	163/221 (74%)	160 (98%)	3 (2%)	0	100	100
1	N	164/221 (74%)	162 (99%)	2 (1%)	0	100	100
All	All	2315/3094 (75%)	2285 (99%)	30 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/185 (77%)	140 (99%)	2 (1%)	59	65
1	B	136/185 (74%)	131 (96%)	5 (4%)	30	29
1	C	142/185 (77%)	138 (97%)	4 (3%)	38	39
1	D	144/185 (78%)	142 (99%)	2 (1%)	59	65
1	E	144/185 (78%)	140 (97%)	4 (3%)	38	39
1	F	138/185 (75%)	131 (95%)	7 (5%)	21	17
1	G	141/185 (76%)	137 (97%)	4 (3%)	38	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	145/185 (78%)	141 (97%)	4 (3%)	38	39
1	I	144/185 (78%)	141 (98%)	3 (2%)	47	51
1	J	139/185 (75%)	138 (99%)	1 (1%)	76	81
1	K	134/185 (72%)	131 (98%)	3 (2%)	45	49
1	L	140/185 (76%)	137 (98%)	3 (2%)	47	51
1	M	143/185 (77%)	141 (99%)	2 (1%)	59	65
1	N	145/185 (78%)	142 (98%)	3 (2%)	47	51
All	All	1977/2590 (76%)	1930 (98%)	47 (2%)	43	45

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	88	MET
1	J	98	LEU
1	H	98	LEU
1	I	78	ARG
1	K	88	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	168	HIS
1	L	168	HIS
1	N	247	HIS
1	L	179	GLN
1	K	247	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 1 is monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	9DF	A	301	-	39,40,40	0.97	3 (7%)	50,58,58	1.10	5 (10%)
2	9DF	N	301	-	39,40,40	0.89	1 (2%)	50,58,58	1.52	7 (14%)
2	9DF	F	301	-	39,40,40	0.96	2 (5%)	50,58,58	1.56	8 (16%)
2	9DF	H	301	-	39,40,40	0.98	4 (10%)	50,58,58	1.30	8 (16%)
2	9DF	H	302	-	39,40,40	0.85	0	50,58,58	1.12	4 (8%)
2	9DF	I	301	-	39,40,40	0.81	0	50,58,58	1.16	7 (14%)
2	9DF	A	302	-	39,40,40	1.01	2 (5%)	50,58,58	1.20	5 (10%)
2	9DF	E	301	-	39,40,40	0.84	1 (2%)	50,58,58	1.31	6 (12%)
2	9DF	J	301	-	39,40,40	0.80	0	50,58,58	1.22	4 (8%)
2	9DF	C	302	-	39,40,40	0.95	2 (5%)	50,58,58	1.16	4 (8%)
2	9DF	L	301	-	39,40,40	0.78	0	50,58,58	1.15	3 (6%)
2	9DF	K	301	-	39,40,40	0.87	1 (2%)	50,58,58	1.30	10 (20%)
2	9DF	D	301	-	39,40,40	0.84	0	50,58,58	1.40	7 (14%)
2	9DF	C	301	-	39,40,40	0.80	1 (2%)	50,58,58	1.19	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9DF	A	301	-	-	7/22/55/55	0/4/4/4
2	9DF	N	301	-	-	0/22/55/55	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9DF	F	301	-	-	5/22/55/55	0/4/4/4
2	9DF	H	301	-	-	7/22/55/55	0/4/4/4
2	9DF	H	302	-	-	6/22/55/55	0/4/4/4
2	9DF	I	301	-	-	9/22/55/55	0/4/4/4
2	9DF	A	302	-	-	5/22/55/55	0/4/4/4
2	9DF	E	301	-	-	6/22/55/55	0/4/4/4
2	9DF	J	301	-	-	6/22/55/55	0/4/4/4
2	9DF	C	302	-	-	3/22/55/55	0/4/4/4
2	9DF	L	301	-	-	2/22/55/55	0/4/4/4
2	9DF	K	301	-	-	5/22/55/55	0/4/4/4
2	9DF	D	301	-	-	5/22/55/55	0/4/4/4
2	9DF	C	301	-	-	5/22/55/55	0/4/4/4

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	301	9DF	C18-N19	3.38	1.50	1.46
2	C	302	9DF	C17-N3	3.03	1.50	1.46
2	A	302	9DF	C17-N3	2.93	1.50	1.46
2	A	301	9DF	C18-N19	2.82	1.50	1.46
2	A	302	9DF	C18-N19	2.69	1.49	1.46

The worst 5 of 82 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	301	9DF	C37-C36-C35	-6.25	101.80	112.61
2	N	301	9DF	C37-C36-C35	-5.91	102.40	112.61
2	D	301	9DF	C5-C4-N3	-4.05	99.25	111.52
2	L	301	9DF	C10-C9-C8	3.73	118.72	111.51
2	E	301	9DF	C10-C9-C8	3.72	118.70	111.51

There are no chirality outliers.

5 of 71 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	301	9DF	C35-C36-C37-C38
2	E	301	9DF	C36-C37-C38-F42
2	H	302	9DF	O1-C2-N3-C17
2	I	301	9DF	C36-C37-C38-F41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	I	301	9DF	C11-C10-C9-C46


There are no ring outliers.

5 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	301	9DF	1	0
2	A	302	9DF	1	0
2	L	301	9DF	1	0
2	K	301	9DF	1	0
2	C	301	9DF	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand 9DF A 301




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF N 301




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF F 301




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF H 301




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF H 302




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF I 301




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF A 302




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF E 301




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF J 301




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF C 302




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF L 301




Bond lengths



Bond angles




Torsions




Rings

Ligand 9DF K 301




Bond lengths



Bond angles




Torsions




Rings


Ligand 9DF D 301




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	170/221 (76%)	0.80	14 (8%) 17 20	36, 51, 79, 114	0
1	B	172/221 (77%)	0.90	19 (11%) 10 12	38, 54, 76, 93	0
1	C	170/221 (76%)	0.70	14 (8%) 17 20	32, 47, 73, 107	0
1	D	171/221 (77%)	0.52	10 (5%) 29 33	28, 42, 61, 84	0
1	E	172/221 (77%)	0.90	21 (12%) 8 10	36, 51, 76, 92	0
1	F	173/221 (78%)	1.15	23 (13%) 7 8	38, 54, 77, 93	0
1	G	171/221 (77%)	0.79	13 (7%) 20 23	37, 52, 73, 101	0
1	H	172/221 (77%)	0.78	8 (4%) 36 41	33, 47, 69, 95	0
1	I	177/221 (80%)	0.84	16 (9%) 15 17	37, 54, 85, 102	0
1	J	171/221 (77%)	1.04	16 (9%) 14 16	43, 58, 81, 100	0
1	K	171/221 (77%)	1.00	19 (11%) 10 12	43, 58, 80, 125	0
1	L	170/221 (76%)	0.77	15 (8%) 15 18	35, 51, 78, 120	0
1	M	169/221 (76%)	0.45	10 (5%) 28 32	28, 41, 62, 81	0
1	N	170/221 (76%)	0.47	8 (4%) 36 41	27, 38, 57, 108	0
All	All	2399/3094 (77%)	0.79	206 (8%) 16 18	27, 51, 76, 125	0

The worst 5 of 206 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	62	VAL	7.0
1	E	72	ALA	5.6
1	K	61	ILE	5.1
1	N	62	VAL	5.1
1	K	140	LEU	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	9DF	L	301	37/37	0.81	0.16	61,87,115,123	0
2	9DF	A	302	37/37	0.82	0.18	74,106,155,158	0
2	9DF	E	301	37/37	0.84	0.17	68,79,120,125	0
3	MG	N	302	1/1	0.86	0.10	57,57,57,57	0
2	9DF	K	301	37/37	0.87	0.15	71,77,137,144	0
2	9DF	I	301	37/37	0.88	0.15	76,93,117,129	0
2	9DF	H	302	37/37	0.89	0.12	57,63,92,97	0
2	9DF	J	301	37/37	0.90	0.13	66,79,111,123	0
2	9DF	A	301	37/37	0.90	0.12	47,65,87,94	0
2	9DF	C	301	37/37	0.90	0.12	44,60,87,94	0
2	9DF	D	301	37/37	0.90	0.12	57,65,90,95	0
2	9DF	C	302	37/37	0.91	0.13	59,70,117,124	0
2	9DF	F	301	37/37	0.91	0.11	50,63,85,100	0
2	9DF	H	301	37/37	0.93	0.09	41,52,74,75	0
2	9DF	N	301	37/37	0.94	0.09	40,45,64,79	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 9DF L 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 9DF A 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9DF E 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around 9DF K 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 9DF I 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9DF H 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 9DF J 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9DF A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9DF C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around 9DF D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 9DF C 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9DF F 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 9DF H 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9DF N 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.