



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:09 PM UTC

PDB ID : 2WIC / pdb_00002wic
Title : Crystal Structures of the N-terminal Intracellular Domain of FeoB from Klebsiella Pneumoniae in GMPPNP binding state
Authors : Hung, K.-W.; Chang, Y.-W.; Chen, J.-H.; Chen, Y.-C.; Sun, Y.-J.; Hsiao, C.-D.; Huang, T.-H.
Deposited on : 2009-05-09
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	21% 70% 16% • • 7%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1977 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FERROUS IRON TRANSPORT PROTEIN B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	0	0
			1883	1182	329	363	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	129	GLN	LYS	conflict	UNP A6TF32

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (CCD ID: GNP) (formula: $C_{10}H_{17}N_6O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			32	10	6	13	3		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	60	Total	O	0	0
			60	60		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FERROUS IRON TRANSPORT PROTEIN B



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	106.90Å 106.90Å 106.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.93 – 2.05 25.93 – 2.05	Depositor EDS
% Data completeness (in resolution range)	83.2 (25.93-2.05) 94.7 (25.93-2.05)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 2.04Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.215 , 0.218 0.231 , 0.235	Depositor DCC
R_{free} test set	1392 reflections (2.88%)	wwPDB-VP
Wilson B-factor (Å ²)	32.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 53.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.038 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1977	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GNP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.16	3/1906 (0.2%)	1.22	15/2593 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	146	ILE	CA-CB	5.91	1.61	1.54
1	A	197	PRO	CA-C	5.67	1.54	1.51
1	A	133	ASP	CA-C	5.26	1.59	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	192	MET	CA-C-N	8.03	136.87	121.54
1	A	192	MET	C-N-CA	8.03	136.87	121.54
1	A	74	GLU	N-CA-C	7.65	120.95	108.55
1	A	221	ASP	CA-CB-CG	7.31	119.91	112.60
1	A	224	ASP	CB-CA-C	7.12	121.04	111.14
1	A	224	ASP	CA-C-O	6.49	131.30	120.65
1	A	75	GLN	N-CA-C	-6.43	104.06	112.41
1	A	225	LYS	N-CA-C	-5.95	106.00	113.20
1	A	195	GLN	CA-C-O	5.68	126.93	120.00
1	A	193	SER	CA-C-N	5.53	132.10	121.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	193	SER	C-N-CA	5.53	132.10	121.54
1	A	224	ASP	O-C-N	-5.47	113.31	121.63
1	A	194	ALA	CA-C-N	5.25	132.82	121.64
1	A	194	ALA	C-N-CA	5.25	132.82	121.64
1	A	173	GLU	N-CA-C	-5.19	100.62	108.67

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	220	GLY	Peptide
1	A	221	ASP	Mainchain
1	A	224	ASP	Peptide
1	A	74	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1883	0	1903	34	0
2	A	32	0	13	1	0
3	A	2	0	0	0	0
4	A	60	0	0	0	0
All	All	1977	0	1916	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:ARG:HG3	1:A:202:ARG:HH11	1.20	1.02
1:A:202:ARG:HG3	1:A:202:ARG:NH1	1.91	0.83
1:A:41:LYS:HG2	1:A:41:LYS:O	1.84	0.78
1:A:193:SER:HB2	1:A:219:ALA:O	1.86	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:ARG:HH11	1:A:202:ARG:CG	2.02	0.71
1:A:41:LYS:O	1:A:41:LYS:CG	2.42	0.66
1:A:111:LEU:HG	1:A:113:ILE:HD11	1.80	0.63
1:A:4:LEU:HD21	1:A:88:MET:HE3	1.81	0.61
1:A:1:MET:HE1	1:A:48:THR:O	2.01	0.61
1:A:111:LEU:HG	1:A:113:ILE:CD1	2.33	0.58
1:A:111:LEU:HD13	1:A:203:TRP:CD2	2.38	0.58
1:A:58:PRO:HB2	1:A:74:GLU:CB	2.34	0.58
1:A:216:ARG:CD	1:A:216:ARG:H	2.21	0.54
1:A:224:ASP:HB3	1:A:227:ASP:OD1	2.08	0.53
1:A:120:ASN:HD21	1:A:150:SER:H	1.57	0.53
1:A:124:ILE:O	1:A:128:GLN:HG2	2.08	0.53
1:A:11:ASN:HD21	1:A:101:ASN:HA	1.74	0.51
1:A:224:ASP:C	1:A:226:LEU:H	2.18	0.51
1:A:50:HIS:HD2	1:A:164:ASP:OD1	1.93	0.51
1:A:44:ILE:HD11	1:A:51:GLN:HG3	1.94	0.50
1:A:210:GLU:HG2	1:A:241:ALA:HA	1.95	0.49
1:A:183:ARG:NH2	1:A:236:GLU:OE2	2.33	0.47
1:A:189:ALA:O	1:A:201:ARG:HD2	2.14	0.47
1:A:119:LEU:HD23	1:A:134:ILE:HD12	1.98	0.45
1:A:224:ASP:HA	1:A:226:LEU:H	1.82	0.45
1:A:77:ALA:O	1:A:81:ILE:HG12	2.18	0.44
1:A:123:ASP:O	1:A:127:LYS:HD3	2.19	0.42
1:A:202:ARG:NH1	1:A:202:ARG:CG	2.68	0.42
1:A:13:ASN:HA	2:A:1265:GNP:HNB3	1.84	0.41
1:A:11:ASN:ND2	1:A:101:ASN:HA	2.36	0.41
1:A:20:PHE:CZ	1:A:41:LYS:HE2	2.56	0.41
1:A:187:LEU:O	1:A:190:GLN:HB2	2.21	0.40
1:A:216:ARG:H	1:A:216:ARG:HD3	1.85	0.40
1:A:192:MET:HB2	1:A:219:ALA:C	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	242/267 (91%)	226 (93%)	9 (4%)	7 (3%)	3	0

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	193	SER
1	A	223	ALA
1	A	219	ALA
1	A	221	ASP
1	A	232	ASN
1	A	192	MET
1	A	194	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/221 (91%)	186 (92%)	15 (8%)	12	6

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	LYS
1	A	44	ILE
1	A	65	THR
1	A	75	GLN
1	A	111	LEU
1	A	127	LYS
1	A	128	GLN
1	A	191	GLN
1	A	193	SER
1	A	202	ARG
1	A	210	GLU
1	A	215	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	216	ARG
1	A	226	LEU
1	A	230	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	50	HIS
1	A	75	GLN
1	A	97	ASN
1	A	120	ASN
1	A	128	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GNP	A	1265	3	34,34,34	2.54	15 (44%)	47,54,54	2.64	18 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GNP	A	1265	3	-	2/18/38/38	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1265	GNP	PB-O3A	6.43	1.67	1.59
2	A	1265	GNP	PG-O1G	5.26	1.54	1.46
2	A	1265	GNP	PA-O3A	4.48	1.64	1.59
2	A	1265	GNP	PB-O2B	4.09	1.67	1.56
2	A	1265	GNP	O4'-C4'	4.03	1.53	1.45
2	A	1265	GNP	PB-N3B	3.89	1.73	1.63
2	A	1265	GNP	C6-N1	3.50	1.45	1.38
2	A	1265	GNP	PG-O3G	3.11	1.65	1.56
2	A	1265	GNP	C4-N9	-2.89	1.30	1.38
2	A	1265	GNP	C5-N7	-2.57	1.33	1.39
2	A	1265	GNP	O4'-C1'	-2.39	1.36	1.42
2	A	1265	GNP	PG-O2G	-2.26	1.50	1.56
2	A	1265	GNP	PA-O5'	-2.23	1.50	1.59
2	A	1265	GNP	C2'-C1'	2.22	1.60	1.53
2	A	1265	GNP	C5-C4	2.17	1.44	1.38

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1265	GNP	O2B-PB-O1B	8.27	127.61	109.87
2	A	1265	GNP	O1B-PB-N3B	7.95	123.47	111.77
2	A	1265	GNP	O2G-PG-O3G	5.13	121.37	107.59
2	A	1265	GNP	O2G-PG-O1G	-4.29	102.70	113.45
2	A	1265	GNP	C2-N3-C4	3.43	118.20	112.30
2	A	1265	GNP	C4-C5-N7	-3.27	105.48	110.67
2	A	1265	GNP	C6-C5-N7	3.24	136.19	130.29
2	A	1265	GNP	O2A-PA-O3A	-3.19	98.64	107.27
2	A	1265	GNP	O1G-PG-N3B	-3.13	107.16	111.77
2	A	1265	GNP	O3'-C3'-C2'	-2.98	102.26	111.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1265	GNP	O6-C6-N1	-2.91	114.64	120.11
2	A	1265	GNP	C5-C4-N3	-2.79	123.95	128.39
2	A	1265	GNP	N2-C2-N3	2.55	124.65	119.67
2	A	1265	GNP	O2B-PB-O3A	-2.55	96.14	104.64
2	A	1265	GNP	O4'-C1'-N9	-2.21	103.36	108.36
2	A	1265	GNP	N9-C8-N7	-2.20	109.32	113.40
2	A	1265	GNP	C8-N7-C5	2.19	108.17	104.26
2	A	1265	GNP	O3A-PB-N3B	-2.17	100.57	106.59

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1265	GNP	PB-N3B-PG-O1G
2	A	1265	GNP	PA-O3A-PB-O1B

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1265	GNP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/267 (92%)	0.94	56 (22%) 2 1	21, 39, 86, 95	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	74	GLU	6.3
1	A	73	ASP	5.9
1	A	67	SER	5.7
1	A	221	ASP	5.5
1	A	229	ALA	5.4
1	A	222	ALA	5.3
1	A	223	ALA	5.0
1	A	218	TYR	4.8
1	A	230	LEU	4.7
1	A	228	ILE	4.6
1	A	217	ALA	4.4
1	A	213	ILE	4.3
1	A	68	SER	4.2
1	A	188	LEU	4.1
1	A	220	GLY	4.1
1	A	189	ALA	4.1
1	A	28	GLN	4.1
1	A	264	THR	4.0
1	A	226	LEU	3.7
1	A	235	ASP	3.6
1	A	219	ALA	3.6
1	A	233	LEU	3.4
1	A	231	ALA	3.3
1	A	237	ILE	3.3
1	A	232	ASN	3.2
1	A	187	LEU	3.1
1	A	196	ILE	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	195	GLN	3.1
1	A	227	ASP	3.0
1	A	216	ARG	2.9
1	A	75	GLN	2.9
1	A	131	ARG	2.8
1	A	76	ILE	2.8
1	A	66	ILE	2.7
1	A	240	PRO	2.6
1	A	191	GLN	2.6
1	A	185	ALA	2.5
1	A	215	SER	2.5
1	A	129	GLN	2.5
1	A	263	LEU	2.5
1	A	192	MET	2.4
1	A	262	THR	2.3
1	A	171	ASP	2.3
1	A	224	ASP	2.3
1	A	152	ARG	2.2
1	A	184	GLU	2.2
1	A	190	GLN	2.2
1	A	199	ARG	2.1
1	A	206	LEU	2.1
1	A	173	GLU	2.1
1	A	197	PRO	2.1
1	A	214	TYR	2.1
1	A	225	LYS	2.1
1	A	77	ALA	2.1
1	A	236	GLU	2.0
1	A	26	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	A	1267	1/1	0.26	0.29	1,1,1,1	1
2	GNP	A	1265	32/32	0.95	0.09	36,41,44,47	0
3	MG	A	1266	1/1	0.99	0.05	40,40,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.