



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:05 PM UTC

PDB ID : 3WIIQ / pdb_00003wiq
Title : Crystal structure of kojibiose phosphorylase complexed with kojibiose
Authors : Okada, S.; Yamamoto, T.; Watanabe, H.; Nishimoto, T.; Chaen, H.; Fukuda, S.; Wakagi, T.; Fushinobu, S.
Deposited on : 2013-09-24
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	764	 3% 72% 22% ...
2	B	2	 50% 50%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6241 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Kojibiose phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	751	6148	3960	1013	1154	21	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	757	GLY	-	expression tag	UNP A4XGP2
A	758	SER	-	expression tag	UNP A4XGP2
A	759	HIS	-	expression tag	UNP A4XGP2
A	760	HIS	-	expression tag	UNP A4XGP2
A	761	HIS	-	expression tag	UNP A4XGP2
A	762	HIS	-	expression tag	UNP A4XGP2
A	763	HIS	-	expression tag	UNP A4XGP2
A	764	HIS	-	expression tag	UNP A4XGP2

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-2)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	2	23	12	11	0	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	65	Total	O	0	0
			65	65		

4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	192.51Å 192.51Å 202.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.13 – 2.80 48.13 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.6 (48.13-2.80) 98.7 (48.13-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.204 , 0.259 0.206 , 0.258	Depositor DCC
R_{free} test set	2338 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	55.9	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 32.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	0.000 for -l,-k,-h 0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6241	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, SO4, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	3/6289 (0.0%)	1.21	30/8497 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	54	VAL	C-O	-7.68	1.14	1.23
1	A	197	LYS	CA-C	-5.04	1.46	1.52
1	A	170	VAL	N-CA	-5.01	1.40	1.46

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	709	GLU	N-CA-C	-8.25	98.27	110.23
1	A	648	THR	N-CA-C	7.70	122.11	112.87
1	A	539	GLU	N-CA-C	-7.37	103.19	111.07
1	A	658	THR	N-CA-CB	6.71	120.67	110.28
1	A	54	VAL	CB-CA-C	6.63	119.67	111.25
1	A	7	GLU	N-CA-C	-6.57	104.37	112.38
1	A	745	VAL	N-CA-C	6.33	117.59	108.48
1	A	131	LYS	N-CA-C	6.17	120.06	112.54
1	A	323	GLU	N-CA-C	5.98	120.27	113.15
1	A	307	VAL	N-CA-C	-5.96	104.50	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	154	PHE	N-CA-C	5.91	118.27	108.99
1	A	666	ASN	CB-CA-C	-5.71	99.07	110.42
1	A	338	LYS	N-CA-C	5.56	119.10	111.54
1	A	491	ASN	N-CA-C	-5.27	102.11	109.96
1	A	625	LYS	CA-CB-CG	5.26	124.63	114.10
1	A	729	ILE	CB-CA-C	-5.25	103.90	111.19
1	A	34	ILE	CB-CA-C	-5.23	103.36	110.42
1	A	626	ARG	N-CA-C	5.23	118.97	112.59
1	A	377	GLU	N-CA-C	-5.17	105.73	111.36
1	A	53	GLY	CA-C-N	-5.16	115.93	122.43
1	A	53	GLY	C-N-CA	-5.16	115.93	122.43
1	A	496	TYR	N-CA-C	-5.16	105.55	111.07
1	A	549	HIS	CA-C-N	5.09	126.21	119.84
1	A	549	HIS	C-N-CA	5.09	126.21	119.84
1	A	304	ASN	CA-C-N	5.07	127.04	120.44
1	A	304	ASN	C-N-CA	5.07	127.04	120.44
1	A	625	LYS	CG-CD-CE	5.07	122.97	111.30
1	A	347	ILE	N-CA-C	5.06	116.16	111.81
1	A	284	GLU	N-CA-C	-5.05	105.90	111.71
1	A	228	GLU	N-CA-C	-5.04	100.25	108.76

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	551	ASP	Peptide
1	A	749	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6148	0	6079	99	0
2	B	23	0	21	1	0
3	A	5	0	0	0	0
4	A	65	0	0	1	0
All	All	6241	0	6100	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:GLU:O	1:A:6:ARG:HB2	1.62	0.97
1:A:352:PHE:O	1:A:356:THR:HB	1.67	0.94
1:A:733:ASN:HA	1:A:750:LYS:HE2	1.65	0.78
1:A:5:GLU:O	1:A:6:ARG:CB	2.30	0.77
1:A:29:ASN:HA	1:A:128:MET:HE2	1.72	0.72
1:A:750:LYS:CB	1:A:751:GLY:HA3	2.22	0.70
1:A:247:VAL:C	1:A:248:LEU:HD12	2.18	0.69
1:A:636:SER:HB3	1:A:658:THR:HG21	1.76	0.68
1:A:750:LYS:HB3	1:A:751:GLY:HA3	1.75	0.68
1:A:248:LEU:HD12	1:A:248:LEU:N	2.11	0.65
1:A:430:TYR:HA	1:A:500:TRP:CH2	2.33	0.63
1:A:21:PHE:O	1:A:25:PHE:HB2	1.97	0.63
1:A:709:GLU:O	1:A:710:LYS:CB	2.47	0.63
1:A:337:TYR:C	1:A:338:LYS:HG2	2.23	0.63
1:A:286:TYR:OH	1:A:316:SER:OG	2.14	0.62
1:A:750:LYS:CB	1:A:751:GLY:CA	2.78	0.61
1:A:239:ARG:HG3	1:A:239:ARG:HH21	1.66	0.61
1:A:747:LEU:HG	1:A:749:ILE:HD11	1.82	0.60
1:A:704:ASN:N	1:A:705:PRO:HD3	2.16	0.60
1:A:286:TYR:HH	1:A:316:SER:HG	1.47	0.59
1:A:709:GLU:O	1:A:710:LYS:HB2	2.04	0.57
1:A:472:ARG:HD3	1:A:545:TYR:HA	1.86	0.57
1:A:703:VAL:HG12	1:A:705:PRO:HD3	1.85	0.57
1:A:63:THR:HG22	1:A:400:GLU:CD	2.30	0.57
1:A:117:ILE:HG22	1:A:143:GLU:HG3	1.86	0.57
1:A:39:GLU:OE2	1:A:138:TYR:OH	2.22	0.57
1:A:257:GLY:N	4:A:924:HOH:O	2.38	0.55
1:A:204:THR:HG22	1:A:205:LYS:N	2.22	0.55
1:A:551:ASP:O	1:A:552:LYS:HB2	2.07	0.55
1:A:450:SER:O	1:A:454:ILE:HG13	2.07	0.54
1:A:239:ARG:HH21	1:A:239:ARG:CG	2.20	0.54
1:A:580:TRP:CD2	1:A:628:MET:HE2	2.42	0.54
1:A:187:LEU:HD23	1:A:187:LEU:C	2.33	0.54
1:A:749:ILE:HG22	1:A:750:LYS:H	1.73	0.52
1:A:352:PHE:CZ	1:A:356:THR:HG21	2.45	0.52
1:A:624:GLU:HG3	1:A:625:LYS:N	2.24	0.51
1:A:580:TRP:CE3	1:A:628:MET:HE2	2.45	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:310:ASN:HA	1:A:679:LEU:HD22	1.92	0.51
1:A:248:LEU:N	1:A:248:LEU:CD1	2.72	0.51
1:A:6:ARG:O	1:A:305:LYS:NZ	2.43	0.51
1:A:634:SER:O	1:A:635:PRO:C	2.50	0.50
1:A:344:ASP:OD1	2:B:2:GLC:O6	2.25	0.49
1:A:317:MET:HE3	1:A:327:LEU:HD13	1.95	0.48
1:A:352:PHE:CE2	1:A:356:THR:HG21	2.49	0.48
1:A:68:LEU:HB3	1:A:69:PRO:HD2	1.96	0.48
1:A:350:LEU:HD23	1:A:431:ALA:HB1	1.95	0.48
1:A:750:LYS:HB2	1:A:751:GLY:C	2.40	0.47
1:A:128:MET:HE3	1:A:316:SER:HA	1.96	0.47
1:A:116:ARG:HB3	1:A:145:TYR:CD2	2.49	0.47
1:A:343:TRP:O	1:A:344:ASP:C	2.56	0.47
1:A:156:ASP:OD1	1:A:156:ASP:C	2.58	0.47
1:A:193:ASP:O	1:A:194:LYS:HB2	2.16	0.47
1:A:720:TRP:CD1	1:A:745:VAL:HG21	2.49	0.46
1:A:749:ILE:O	1:A:750:LYS:C	2.59	0.46
1:A:747:LEU:HD12	1:A:748:LYS:H	1.80	0.46
1:A:350:LEU:HB3	1:A:351:PRO:HD3	1.98	0.46
1:A:22:GLU:HB3	1:A:35:ARG:NH2	2.31	0.46
1:A:570:LYS:O	1:A:571:GLU:HB2	2.16	0.46
1:A:129:ASN:HB3	1:A:285:GLU:HG3	1.97	0.45
1:A:29:ASN:HA	1:A:128:MET:CE	2.43	0.45
1:A:187:LEU:HD23	1:A:188:GLY:N	2.32	0.45
1:A:280:PHE:O	1:A:281:GLU:C	2.59	0.45
1:A:621:ASP:OD1	1:A:647:GLU:OE2	2.34	0.45
1:A:569:ILE:O	1:A:570:LYS:O	2.35	0.45
1:A:734:VAL:O	1:A:734:VAL:CG2	2.64	0.44
1:A:533:GLU:HB2	1:A:534:PRO:HD3	1.98	0.44
1:A:432:VAL:HB	1:A:453:ILE:HD11	1.98	0.44
1:A:749:ILE:HG22	1:A:750:LYS:N	2.32	0.44
1:A:750:LYS:HB2	1:A:751:GLY:CA	2.47	0.44
1:A:510:ASN:O	1:A:511:LEU:C	2.59	0.44
1:A:548:TYR:HB3	1:A:549:HIS:CA	2.48	0.44
1:A:496:TYR:CD1	1:A:603:LEU:HD21	2.52	0.44
1:A:336:GLY:C	1:A:338:LYS:H	2.26	0.44
1:A:204:THR:CG2	1:A:205:LYS:N	2.82	0.43
1:A:166:PRO:HA	1:A:169:ARG:NH1	2.33	0.43
1:A:435:TYR:CE1	1:A:439:THR:HG21	2.54	0.43
1:A:180:ASP:HB2	1:A:219:THR:HG21	2.01	0.43
1:A:533:GLU:N	1:A:534:PRO:CD	2.82	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:ASP:OD1	1:A:98:ASP:C	2.63	0.42
1:A:235:LYS:O	1:A:236:GLN:C	2.62	0.42
1:A:137:LYS:HA	1:A:242:ILE:O	2.20	0.42
1:A:42:PHE:O	1:A:43:CYS:C	2.60	0.41
1:A:649:ASN:O	1:A:650:ARG:C	2.64	0.41
1:A:128:MET:CE	1:A:316:SER:HA	2.50	0.41
1:A:445:LEU:O	1:A:450:SER:OG	2.38	0.41
1:A:548:TYR:HB3	1:A:549:HIS:C	2.45	0.41
1:A:613:ASP:OD1	1:A:613:ASP:N	2.50	0.41
1:A:77:TYR:O	1:A:149:LEU:HA	2.21	0.41
1:A:108:LEU:HD23	1:A:108:LEU:HA	1.89	0.41
1:A:630:LYS:HD3	1:A:630:LYS:HA	1.78	0.41
1:A:654:ASN:OD1	1:A:657:ARG:NH2	2.54	0.41
1:A:543:LYS:O	1:A:544:ILE:C	2.63	0.41
1:A:330:LYS:HA	1:A:674:ILE:O	2.21	0.41
1:A:446:LEU:HD23	1:A:512:LEU:HD21	2.03	0.40
1:A:319:ASN:HA	1:A:320:PRO:HD2	1.88	0.40
1:A:510:ASN:O	1:A:514:GLU:HG2	2.21	0.40
1:A:565:LYS:HG3	1:A:566:ASP:N	2.36	0.40
1:A:717:SER:O	1:A:718:ILE:HG23	2.20	0.40
1:A:658:THR:HB	1:A:682:THR:OG1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	749/764 (98%)	688 (92%)	46 (6%)	15 (2%)	6	21

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	569	ILE
1	A	570	LYS
1	A	609	GLU
1	A	750	LYS
1	A	6	ARG
1	A	545	TYR
1	A	552	LYS
1	A	710	LYS
1	A	469	GLU
1	A	706	TRP
1	A	625	LYS
1	A	721	LYS
1	A	291	ASP
1	A	571	GLU
1	A	41	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	665/676 (98%)	604 (91%)	61 (9%)	8 27

All (61) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	LYS
1	A	14	LYS
1	A	15	LEU
1	A	54	VAL
1	A	74	LEU
1	A	89	GLU
1	A	127	SER
1	A	170	VAL
1	A	174	GLU
1	A	219	THR
1	A	226	ILE
1	A	229	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	239	ARG
1	A	241	GLU
1	A	261	GLU
1	A	264	THR
1	A	266	LYS
1	A	277	LYS
1	A	287	LYS
1	A	306	SER
1	A	338	LYS
1	A	351	PRO
1	A	356	THR
1	A	440	ASP
1	A	450	SER
1	A	453	ILE
1	A	472	ARG
1	A	478	VAL
1	A	483	GLU
1	A	488	CYS
1	A	497	LEU
1	A	507	GLU
1	A	508	LEU
1	A	513	LEU
1	A	514	GLU
1	A	530	SER
1	A	539	GLU
1	A	543	LYS
1	A	552	LYS
1	A	565	LYS
1	A	568	VAL
1	A	570	LYS
1	A	574	GLN
1	A	582	GLU
1	A	609	GLU
1	A	615	THR
1	A	624	GLU
1	A	653	ILE
1	A	658	THR
1	A	664	GLU
1	A	678	SER
1	A	693	ILE
1	A	697	LYS
1	A	702	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	710	LYS
1	A	731	LYS
1	A	734	VAL
1	A	735	GLU
1	A	742	LYS
1	A	746	LYS
1	A	750	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	GLN
1	A	136	GLN
1	A	319	ASN
1	A	382	ASN
1	A	476	ASN
1	A	501	ASN
1	A	597	GLN
1	A	667	GLN
1	A	671	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	B	1	2	12,12,12	0.48	0	17,17,17	1.55	3 (17%)
2	GLC	B	2	2	11,11,12	0.85	0	15,15,17	1.79	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	GLC	C1-O5-C5	3.55	116.94	112.19
2	B	1	BGC	C1-C2-C3	-3.51	103.20	110.36
2	B	2	GLC	O3-C3-C4	3.13	117.75	110.38
2	B	1	BGC	O2-C2-C1	3.00	116.17	109.25
2	B	2	GLC	C2-C3-C4	-2.96	105.65	110.86
2	B	2	GLC	O4-C4-C3	2.56	116.41	110.38
2	B	1	BGC	O5-C1-C2	-2.42	106.04	110.30

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	803	-	4,4,4	0.93	0	6,6,6	1.13	1 (16%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	803	SO4	O4-S-O1	-2.04	98.88	109.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	751/764 (98%)	0.03	26 (3%)	47 38	35, 52, 82, 150	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	547	PRO	7.9
1	A	548	TYR	6.9
1	A	751	GLY	5.7
1	A	549	HIS	5.2
1	A	625	LYS	5.2
1	A	552	LYS	4.7
1	A	550	PRO	4.7
1	A	568	VAL	4.3
1	A	697	LYS	4.2
1	A	570	LYS	3.8
1	A	744	ASN	3.6
1	A	709	GLU	3.3
1	A	525	LYS	2.9
1	A	224	TYR	2.8
1	A	745	VAL	2.8
1	A	551	ASP	2.7
1	A	750	LYS	2.6
1	A	6	ARG	2.5
1	A	698	ASP	2.5
1	A	176	ASP	2.3
1	A	743	GLY	2.3
1	A	416	TRP	2.1
1	A	738	LYS	2.0
1	A	747	LEU	2.0
1	A	414	ARG	2.0
1	A	284	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BGC	B	1	12/12	0.95	0.09	47,55,59,71	0
2	GLC	B	2	11/12	0.96	0.07	40,45,54,54	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	803	5/5	0.91	0.13	55,55,74,83	0

6.5 Other polymers [i](#)

There are no such residues in this entry.