



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 04:05 PM UTC

PDB ID : 3WIR / pdb_00003wir
Title : Crystal structure of kojibiose phosphorylase complexed with glucose
Authors : Okada, S.; Yamamoto, T.; Watanabe, H.; Nishimoto, T.; Chaen, H.; Fukuda, S.; Wakagi, T.; Fushinobu, S.
Deposited on : 2013-09-24
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	764	
1	B	764	
1	C	764	
1	D	764	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 26306 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kojibiose phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	756	Total	C	N	O	S	0	0	0
			6187	3985	1019	1162	21			
1	B	756	Total	C	N	O	S	0	0	0
			6187	3985	1019	1162	21			
1	C	756	Total	C	N	O	S	0	0	0
			6187	3985	1019	1162	21			
1	D	756	Total	C	N	O	S	0	0	0
			6187	3985	1019	1162	21			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	757	GLY	-	expression tag	UNP A4XGP2
A	758	SER	-	expression tag	UNP A4XGP2
A	759	HIS	-	expression tag	UNP A4XGP2
A	760	HIS	-	expression tag	UNP A4XGP2
A	761	HIS	-	expression tag	UNP A4XGP2
A	762	HIS	-	expression tag	UNP A4XGP2
A	763	HIS	-	expression tag	UNP A4XGP2
A	764	HIS	-	expression tag	UNP A4XGP2
B	757	GLY	-	expression tag	UNP A4XGP2
B	758	SER	-	expression tag	UNP A4XGP2
B	759	HIS	-	expression tag	UNP A4XGP2
B	760	HIS	-	expression tag	UNP A4XGP2
B	761	HIS	-	expression tag	UNP A4XGP2
B	762	HIS	-	expression tag	UNP A4XGP2
B	763	HIS	-	expression tag	UNP A4XGP2
B	764	HIS	-	expression tag	UNP A4XGP2
C	757	GLY	-	expression tag	UNP A4XGP2
C	758	SER	-	expression tag	UNP A4XGP2
C	759	HIS	-	expression tag	UNP A4XGP2
C	760	HIS	-	expression tag	UNP A4XGP2
C	761	HIS	-	expression tag	UNP A4XGP2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	762	HIS	-	expression tag	UNP A4XGP2
C	763	HIS	-	expression tag	UNP A4XGP2
C	764	HIS	-	expression tag	UNP A4XGP2
D	757	GLY	-	expression tag	UNP A4XGP2
D	758	SER	-	expression tag	UNP A4XGP2
D	759	HIS	-	expression tag	UNP A4XGP2
D	760	HIS	-	expression tag	UNP A4XGP2
D	761	HIS	-	expression tag	UNP A4XGP2
D	762	HIS	-	expression tag	UNP A4XGP2
D	763	HIS	-	expression tag	UNP A4XGP2
D	764	HIS	-	expression tag	UNP A4XGP2

- Molecule 2 is beta-D-glucopyranose (CCD ID: BGC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			12	6	6		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	C	O	0	0
			12	6	6		
2	D	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	C	1	Total	O	P	0	0
			5	4	1		
4	C	1	Total	O	P	0	0
			5	4	1		
4	D	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	371	Total	O	0	0
			371	371		
5	B	387	Total	O	0	0
			387	387		
5	C	299	Total	O	0	0
			299	299		
5	D	321	Total	O	0	0
			321	321		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Kojibiose phosphorylase




• Molecule 1: Kojibiose phosphorylase





• Molecule 1: Kojibiose phosphorylase





• Molecule 1: Kojibiose phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	71.58Å 104.46Å 124.16Å 68.83° 86.02° 90.06°	Depositor
Resolution (Å)	46.26 – 2.05 46.26 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.0 (46.26-2.05) 97.1 (46.26-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.191 , 0.251 0.196 , 0.254	Depositor DCC
R_{free} test set	10235 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	26.3	Xtriage
Anisotropy	0.153	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	26306	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.80 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.0405e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, PO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.22	14/6328 (0.2%)	1.12	7/8550 (0.1%)
1	B	1.19	15/6328 (0.2%)	1.14	10/8550 (0.1%)
1	C	1.08	0/6328	1.08	14/8550 (0.2%)
1	D	1.10	8/6328 (0.1%)	1.10	18/8550 (0.2%)
All	All	1.15	37/25312 (0.1%)	1.11	49/34200 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
All	All	0	2

The worst 5 of 37 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	679	LEU	C-O	6.97	1.32	1.24
1	B	252	ARG	CA-C	6.95	1.62	1.52
1	B	68	LEU	CA-C	-6.93	1.44	1.52
1	B	251	SER	N-CA	-6.92	1.36	1.46
1	A	95	ARG	CA-C	-6.45	1.45	1.52

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	658	THR	CB-CA-C	-8.37	96.62	110.85
1	B	259	VAL	CB-CA-C	-7.64	101.82	112.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	350	LEU	CA-C-N	-7.52	111.01	119.28
1	C	350	LEU	C-N-CA	-7.52	111.01	119.28
1	D	743	GLY	N-CA-C	7.41	122.28	111.24

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	551	ASP	Peptide
1	C	255	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6187	0	6120	51	0
1	B	6187	0	6119	61	0
1	C	6187	0	6120	62	0
1	D	6187	0	6119	71	0
2	A	24	0	24	0	0
2	B	24	0	24	0	0
2	C	24	0	24	0	0
2	D	24	0	24	0	0
3	A	12	0	16	0	0
3	B	6	0	8	1	0
3	C	12	0	16	0	0
3	D	24	0	32	2	0
4	A	5	0	0	0	0
4	B	10	0	0	1	0
4	C	10	0	0	1	0
4	D	5	0	0	0	0
5	A	371	0	0	6	0
5	B	387	0	0	2	0
5	C	299	0	0	4	0
5	D	321	0	0	10	0
All	All	26306	0	24646	244	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 244 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:MET:HE3	1:B:103:ILE:HD11	1.31	1.11
1:B:303:ALA:HB2	1:B:656:MET:HE1	1.29	1.09
1:D:636:SER:HB3	1:D:658:THR:HG21	1.54	0.87
1:B:303:ALA:CB	1:B:656:MET:HE1	2.06	0.85
1:B:303:ALA:HB2	1:B:656:MET:CE	2.07	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	754/764 (99%)	724 (96%)	27 (4%)	3 (0%)	30	22
1	B	754/764 (99%)	729 (97%)	23 (3%)	2 (0%)	36	30
1	C	754/764 (99%)	726 (96%)	23 (3%)	5 (1%)	18	10
1	D	754/764 (99%)	717 (95%)	33 (4%)	4 (0%)	24	16
All	All	3016/3056 (99%)	2896 (96%)	106 (4%)	14 (0%)	24	16

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	742	LYS
1	D	259	VAL
1	D	741	GLU
1	D	258	ASP
1	B	742	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	669/676 (99%)	643 (96%)	26 (4%)	28	23
1	B	669/676 (99%)	643 (96%)	26 (4%)	28	23
1	C	669/676 (99%)	633 (95%)	36 (5%)	20	12
1	D	669/676 (99%)	628 (94%)	41 (6%)	17	10
All	All	2676/2704 (99%)	2547 (95%)	129 (5%)	23	16

5 of 129 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	650	ARG
1	D	701	LEU
1	B	734	VAL
1	B	715	LYS
1	D	709	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	136	GLN
1	D	538	GLN
1	D	597	GLN
1	D	549	HIS
1	B	101	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	A	802	-	12,12,12	0.60	0	17,17,17	1.85	5 (29%)
3	GOL	C	804	-	5,5,5	0.42	0	5,5,5	0.29	0
4	PO4	B	805	-	4,4,4	1.28	0	6,6,6	1.21	1 (16%)
3	GOL	D	805	-	5,5,5	1.16	1 (20%)	5,5,5	1.03	0
4	PO4	A	805	-	4,4,4	0.62	0	6,6,6	1.26	1 (16%)
3	GOL	A	804	-	5,5,5	0.86	0	5,5,5	1.26	1 (20%)
4	PO4	D	807	-	4,4,4	0.94	0	6,6,6	1.11	1 (16%)
2	BGC	D	801	-	12,12,12	0.66	0	17,17,17	1.00	1 (5%)
2	BGC	B	801	-	12,12,12	1.18	1 (8%)	17,17,17	1.19	2 (11%)
4	PO4	C	806	-	4,4,4	1.00	0	6,6,6	0.91	0
2	BGC	D	802	-	12,12,12	1.29	2 (16%)	17,17,17	1.71	6 (35%)
3	GOL	B	803	-	5,5,5	0.49	0	5,5,5	1.03	0
3	GOL	A	803	-	5,5,5	0.79	0	5,5,5	0.62	0
3	GOL	C	803	-	5,5,5	0.72	0	5,5,5	0.48	0
2	BGC	B	802	-	12,12,12	0.58	0	17,17,17	1.22	2 (11%)
4	PO4	C	805	-	4,4,4	0.91	0	6,6,6	1.42	1 (16%)
3	GOL	D	803	-	5,5,5	0.76	0	5,5,5	0.44	0
3	GOL	D	804	-	5,5,5	0.81	0	5,5,5	1.15	0
2	BGC	C	801	-	12,12,12	0.53	0	17,17,17	1.34	2 (11%)
3	GOL	D	806	-	5,5,5	0.63	0	5,5,5	0.48	0
2	BGC	A	801	-	12,12,12	0.68	0	17,17,17	1.57	5 (29%)
4	PO4	B	804	-	4,4,4	1.94	2 (50%)	6,6,6	1.60	1 (16%)
2	BGC	C	802	-	12,12,12	1.03	1 (8%)	17,17,17	1.07	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	802	-	-	0/2/22/22	0/1/1/1
2	BGC	C	801	-	-	0/2/22/22	0/1/1/1
2	BGC	A	802	-	-	0/2/22/22	0/1/1/1
3	GOL	D	806	-	-	4/4/4/4	-
2	BGC	D	801	-	-	0/2/22/22	0/1/1/1
3	GOL	D	803	-	-	1/4/4/4	-
3	GOL	C	804	-	-	2/4/4/4	-
2	BGC	A	801	-	-	0/2/22/22	0/1/1/1
2	BGC	B	801	-	-	0/2/22/22	0/1/1/1
3	GOL	D	805	-	-	2/4/4/4	-
2	BGC	D	802	-	-	0/2/22/22	0/1/1/1
3	GOL	A	804	-	-	1/4/4/4	-
3	GOL	B	803	-	-	3/4/4/4	-
3	GOL	D	804	-	-	2/4/4/4	-
3	GOL	A	803	-	-	0/4/4/4	-
2	BGC	C	802	-	-	0/2/22/22	0/1/1/1
3	GOL	C	803	-	-	4/4/4/4	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	804	PO4	P-O1	3.13	1.57	1.50
2	D	802	BGC	O5-C5	-3.11	1.36	1.44
3	D	805	GOL	O2-C2	2.46	1.50	1.43
2	B	801	BGC	O4-C4	2.46	1.49	1.43
2	C	802	BGC	O1-C1	2.44	1.47	1.39

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	802	BGC	O2-C2-C1	-3.54	101.07	109.25
4	B	804	PO4	O4-P-O3	3.44	118.62	107.91
2	B	802	BGC	O1-C1-O5	-3.23	100.82	110.41
2	C	801	BGC	C4-C3-C2	-3.13	105.33	110.83
2	D	802	BGC	C1-C2-C3	3.09	116.66	110.36

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	804	GOL	O1-C1-C2-C3
3	D	804	GOL	O1-C1-C2-C3
3	D	805	GOL	O1-C1-C2-C3
3	D	806	GOL	O1-C1-C2-C3
3	D	804	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	805	PO4	1	0
3	D	805	GOL	2	0
3	B	803	GOL	1	0
4	C	805	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	756/764 (98%)	0.13	19 (2%) 58 60	13, 27, 53, 93	0
1	B	756/764 (98%)	0.08	11 (1%) 72 73	13, 26, 51, 69	0
1	C	756/764 (98%)	0.22	13 (1%) 69 70	17, 30, 54, 72	0
1	D	756/764 (98%)	0.24	17 (2%) 62 64	16, 30, 55, 75	0
All	All	3024/3056 (98%)	0.17	60 (1%) 65 67	13, 29, 53, 93	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	550	PRO	4.2
1	A	548	TYR	4.0
1	D	408	TYR	3.7
1	A	755	ILE	3.5
1	D	496	TYR	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	D	804	6/6	0.61	0.26	55,57,60,64	0
3	GOL	C	803	6/6	0.71	0.20	46,55,59,62	0
3	GOL	D	803	6/6	0.72	0.19	44,54,61,61	0
3	GOL	A	804	6/6	0.81	0.18	33,35,40,41	0
3	GOL	A	803	6/6	0.82	0.15	29,41,45,47	0
3	GOL	D	805	6/6	0.83	0.13	27,36,38,50	0
3	GOL	D	806	6/6	0.84	0.13	46,47,49,49	0
3	GOL	C	804	6/6	0.87	0.11	47,49,50,55	0
4	PO4	B	805	5/5	0.89	0.13	42,43,47,47	0
3	GOL	B	803	6/6	0.90	0.14	30,40,42,42	0
4	PO4	C	806	5/5	0.90	0.12	45,46,46,50	0
2	BGC	D	802	12/12	0.94	0.07	21,26,27,27	0
2	BGC	A	801	12/12	0.94	0.06	19,21,23,24	0
2	BGC	A	802	12/12	0.94	0.07	21,26,29,32	0
2	BGC	B	802	12/12	0.94	0.07	19,26,29,31	0
2	BGC	B	801	12/12	0.95	0.06	20,22,24,26	0
2	BGC	C	801	12/12	0.95	0.06	19,24,25,25	0
2	BGC	C	802	12/12	0.95	0.06	20,25,29,30	0
2	BGC	D	801	12/12	0.95	0.06	21,24,25,25	0
4	PO4	B	804	5/5	0.98	0.04	18,21,23,26	0
4	PO4	D	807	5/5	0.98	0.06	22,25,27,28	0
4	PO4	A	805	5/5	0.99	0.05	20,20,23,24	0
4	PO4	C	805	5/5	0.99	0.06	22,24,26,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.