



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 19, 2026 – 07:34 PM UTC

PDB ID : 3WJL / pdb_00003wj
Title : Crystal structure of IIb selective Fc variant, Fc(V12), in complex with FcgRIIb
Authors : Kadono, S.; Mimoto, F.; Katada, H.; Igawa, T.; Kuramochi, T.; Muraoka, M.;
Wada, Y.; Haraya, K.; Miyazaki, T.; Hattori, K.
Deposited on : 2013-10-11
Resolution : 2.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	7% 85% 7% 8%
1	B	230	2% 78% 11% 10%
2	C	179	9% 80% 9% 11%
3	D	8	50% 50%
4	E	9	89% 11%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	0	0
			1626	1039	265	316	6			
1	B	208	Total	C	N	O	S	0	0	0
			1643	1041	272	324	6			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	220	SER	CYS	engineered mutation	UNP P01857
A	233	ASP	GLU	engineered mutation	UNP P01857
A	237	ASP	GLY	engineered mutation	UNP P01857
A	238	ASP	PRO	engineered mutation	UNP P01857
A	268	ASP	HIS	engineered mutation	UNP P01857
A	271	GLY	PRO	engineered mutation	UNP P01857
A	330	ARG	ALA	engineered mutation	UNP P01857
B	220	SER	CYS	engineered mutation	UNP P01857
B	233	ASP	GLU	engineered mutation	UNP P01857
B	237	ASP	GLY	engineered mutation	UNP P01857
B	238	ASP	PRO	engineered mutation	UNP P01857
B	268	ASP	HIS	engineered mutation	UNP P01857
B	271	GLY	PRO	engineered mutation	UNP P01857
B	330	ARG	ALA	engineered mutation	UNP P01857


- Molecule 2 is a protein called Low affinity immunoglobulin gamma Fc region receptor II-c.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	160	Total	C	N	O	S	0	0	0
			1233	785	210	234	4			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	173	HIS	-	expression tag	UNP P31995
C	174	HIS	-	expression tag	UNP P31995
C	175	HIS	-	expression tag	UNP P31995
C	176	HIS	-	expression tag	UNP P31995
C	177	HIS	-	expression tag	UNP P31995
C	178	HIS	-	expression tag	UNP P31995

- Molecule 3 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	8	Total	C	N	O	0	0	0
			100	56	4	40			

- Molecule 4 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	9	Total	C	N	O	0	0	0
			110	62	4	44			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	17	Total	O	0	0
			17	17		
6	B	21	Total	O	0	0
			21	21		
6	C	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ig gamma-1 chain C region



- Molecule 1: Ig gamma-1 chain C region



- Molecule 2: Low affinity immunoglobulin gamma Fc region receptor II-c



- Molecule 3: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  50% 50%



- Molecule 4: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  89% 11%



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	156.71Å 260.20Å 56.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.86 25.00 – 2.86	Depositor EDS
% Data completeness (in resolution range)	99.4 (25.00-2.86) 99.2 (25.00-2.86)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.86Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.230 , 0.270 0.231 , 0.270	Depositor DCC
R_{free} test set	1366 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	64.8	Xtriage
Anisotropy	0.287	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 31.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.016 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.023 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4768	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, MAN, NAG, BMA, FUL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1669	0.74	0/2285
1	B	0.43	0/1686	0.76	0/2303
2	C	0.42	0/1267	0.68	0/1732
All	All	0.42	0/4622	0.73	0/6320

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1626	0	1517	10	0
1	B	1643	0	1553	15	0
2	C	1233	0	1153	6	0
3	D	100	0	85	0	0
4	E	110	0	94	1	0
5	C	14	0	13	0	0
6	A	17	0	0	0	0
6	B	21	0	0	0	0
6	C	4	0	0	0	0
All	All	4768	0	4415	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:346:PRO:HB3	1:B:372:PHE:HB3	1.68	0.76
2:C:110:TRP:O	2:C:113:LYS:HG3	1.92	0.70
1:A:294:GLU:HG3	1:A:295:GLN:H	1.59	0.67
1:A:429:HIS:CD2	1:A:431:ALA:H	2.17	0.61
1:B:292:ARG:HG3	1:B:300:TYR:CD1	2.40	0.57
2:C:10:GLU:HB3	2:C:23:THR:HB	1.87	0.56
1:B:344:ARG:NH2	1:B:401:ASP:O	2.39	0.56
4:E:2:NAG:HN2	4:E:9:FUL:H61	1.70	0.55
1:A:429:HIS:HD2	1:A:431:ALA:H	1.57	0.52
1:B:367:CYS:HB2	1:B:381:TRP:CZ2	2.46	0.50
1:B:429:HIS:CD2	1:B:431:ALA:H	2.30	0.50
1:B:277:TRP:CE3	1:B:306:LEU:HD22	2.47	0.49
1:B:418:GLN:HA	1:B:443:LEU:HD22	1.94	0.49
1:A:294:GLU:HG3	1:A:295:GLN:N	2.27	0.48
1:B:325:ASN:HB3	1:B:328:LEU:HD22	1.95	0.48
1:A:378:ALA:HB3	1:A:428:MET:HB2	1.96	0.48
1:A:248:LYS:HE3	1:A:255:ARG:HD2	1.95	0.48
1:B:266:VAL:HB	1:B:300:TYR:HB2	1.96	0.47
1:B:430:GLU:HA	1:B:435:HIS:CD2	2.49	0.47
2:C:24:LEU:HD12	2:C:54:TYR:HD2	1.81	0.46
1:B:443:LEU:HG	1:B:444:SER:N	2.32	0.44
1:B:312:ASP:O	1:B:317:LYS:HB2	2.17	0.44
1:A:330:ARG:HG3	2:C:85:SER:O	2.19	0.43
1:B:406:LEU:HD12	1:B:406:LEU:C	2.44	0.42
1:B:386:GLN:HA	1:B:387:PRO:HD3	1.92	0.42
1:A:379:VAL:HG21	1:A:406:LEU:HD11	2.02	0.42
2:C:5:ALA:HB3	2:C:76:SER:HB3	2.01	0.41
1:A:276:ASN:HB2	1:A:322:LYS:HB3	2.03	0.41
1:B:367:CYS:HB2	1:B:381:TRP:CH2	2.56	0.41
1:A:256:THR:HA	1:A:257:PRO:HD3	1.86	0.40
2:C:90:LEU:HD21	2:C:149:TYR:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/230 (91%)	202 (96%)	8 (4%)	0	100	100
1	B	206/230 (90%)	198 (96%)	8 (4%)	0	100	100
2	C	156/179 (87%)	151 (97%)	5 (3%)	0	100	100
All	All	572/639 (90%)	551 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/215 (82%)	173 (98%)	3 (2%)	53	75
1	B	186/215 (86%)	179 (96%)	7 (4%)	29	54
2	C	136/164 (83%)	131 (96%)	5 (4%)	30	55
All	All	498/594 (84%)	483 (97%)	15 (3%)	36	61

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	250	THR
1	A	256	THR
1	A	442	SER
1	B	268	ASP
1	B	292	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	302	VAL
1	B	328	LEU
1	B	355	ARG
1	B	382	GLU
1	B	386	GLN
2	C	9	LEU
2	C	67	THR
2	C	116	VAL
2	C	167	ILE
2	C	168	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	276	ASN
1	A	384	ASN
1	A	389	ASN
1	A	390	ASN
1	A	418	GLN
1	A	429	HIS
1	B	285	HIS
1	B	295	GLN
1	B	390	ASN
1	B	434	ASN
1	B	435	HIS
2	C	59	ASN
2	C	80	HIS
2	C	94	HIS
2	C	108	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

17 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	1	3,1	14,14,15	0.49	0	17,19,21	1.08	1 (5%)
3	NAG	D	2	3	14,14,15	0.52	0	17,19,21	0.91	0
3	BMA	D	3	3	11,11,12	0.54	0	15,15,17	0.96	1 (6%)
3	MAN	D	4	3	11,11,12	0.59	0	15,15,17	0.76	1 (6%)
3	NAG	D	5	3	14,14,15	0.49	0	17,19,21	0.76	0
3	GAL	D	6	3	11,11,12	0.59	0	15,15,17	0.61	0
3	MAN	D	7	3	11,11,12	0.50	0	15,15,17	1.06	1 (6%)
3	NAG	D	8	3	14,14,15	0.56	0	17,19,21	0.77	0
4	NAG	E	1	1,4	14,14,15	0.41	0	17,19,21	1.07	1 (5%)
4	NAG	E	2	4	14,14,15	0.54	0	17,19,21	0.89	1 (5%)
4	BMA	E	3	4	11,11,12	0.34	0	15,15,17	1.06	2 (13%)
4	MAN	E	4	4	11,11,12	0.48	0	15,15,17	1.44	1 (6%)
4	NAG	E	5	4	14,14,15	0.66	0	17,19,21	1.05	1 (5%)
4	GAL	E	6	4	11,11,12	0.60	0	15,15,17	1.18	3 (20%)
4	MAN	E	7	4	11,11,12	0.50	0	15,15,17	1.48	2 (13%)
4	NAG	E	8	4	14,14,15	0.56	0	17,19,21	1.03	2 (11%)
4	FUL	E	9	4	10,10,11	0.57	0	14,14,16	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	1/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	NAG	D	5	3	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GAL	D	6	3	-	1/2/19/22	0/1/1/1
3	MAN	D	7	3	-	2/2/19/22	0/1/1/1
3	NAG	D	8	3	-	2/6/23/26	0/1/1/1
4	NAG	E	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	E	2	4	-	0/6/23/26	0/1/1/1
4	BMA	E	3	4	-	0/2/19/22	0/1/1/1
4	MAN	E	4	4	-	0/2/19/22	0/1/1/1
4	NAG	E	5	4	-	0/6/23/26	0/1/1/1
4	GAL	E	6	4	-	2/2/19/22	0/1/1/1
4	MAN	E	7	4	-	0/2/19/22	0/1/1/1
4	NAG	E	8	4	-	2/6/23/26	0/1/1/1
4	FUL	E	9	4	-	-	0/1/1/1

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	4	MAN	C1-O5-C5	4.94	118.81	112.19
4	E	7	MAN	C1-O5-C5	3.88	117.39	112.19
4	E	1	NAG	O5-C5-C6	3.23	113.94	107.66
4	E	5	NAG	C4-C3-C2	3.03	115.46	111.02
4	E	7	MAN	C1-C2-C3	-2.94	105.37	109.64
3	D	3	BMA	C1-O5-C5	2.84	116.00	112.19
4	E	2	NAG	C1-O5-C5	2.84	115.99	112.19
4	E	3	BMA	C1-O5-C5	2.77	115.90	112.19
3	D	7	MAN	C1-O5-C5	2.70	115.80	112.19
3	D	1	NAG	C4-C3-C2	2.67	114.93	111.02
4	E	6	GAL	C1-O5-C5	2.44	115.45	112.19
4	E	8	NAG	C4-C3-C2	2.40	114.53	111.02
4	E	8	NAG	C3-C4-C5	2.37	114.52	110.23
3	D	4	MAN	C1-O5-C5	2.26	115.22	112.19
4	E	6	GAL	O5-C5-C6	2.16	111.87	107.66
4	E	6	GAL	C1-C2-C3	2.12	112.73	109.64
4	E	3	BMA	C1-C2-C3	2.11	112.71	109.64

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	1	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	D	5	NAG	O5-C5-C6-O6
4	E	6	GAL	O5-C5-C6-O6
3	D	5	NAG	C4-C5-C6-O6
3	D	8	NAG	O5-C5-C6-O6
3	D	8	NAG	C4-C5-C6-O6
4	E	1	NAG	C4-C5-C6-O6
4	E	8	NAG	C4-C5-C6-O6
3	D	7	MAN	C4-C5-C6-O6
4	E	6	GAL	C4-C5-C6-O6
4	E	8	NAG	O5-C5-C6-O6
3	D	7	MAN	O5-C5-C6-O6
3	D	6	GAL	O5-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	9	FUL	1	0
4	E	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	C	1001	2	14,14,15	0.55	0	17,19,21	1.26	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	1001	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	1001	NAG	C3-C4-C5	2.92	115.52	110.23
5	C	1001	NAG	C4-C3-C2	2.30	114.39	111.02
5	C	1001	NAG	C1-O5-C5	2.07	114.96	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	1001	NAG	C4-C5-C6-O6
5	C	1001	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	212/230 (92%)	0.51	15 (7%) 22 16	31, 65, 122, 147	1 (0%)
1	B	208/230 (90%)	0.34	5 (2%) 59 51	28, 58, 78, 89	1 (0%)
2	C	160/179 (89%)	0.61	17 (10%) 11 9	41, 67, 107, 127	0
All	All	580/639 (90%)	0.47	37 (6%) 25 19	28, 62, 107, 147	2 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	3	PRO	5.0
1	A	300	TYR	4.9
1	A	301	ARG	4.3
2	C	72	GLN	4.2
1	A	291	PRO	4.1
2	C	38	GLN	4.0
1	A	232	PRO	3.9
2	C	49	HIS	3.8
1	A	290	LYS	3.6
1	A	292	ARG	3.5
2	C	75	LEU	3.5
1	A	311	GLN	3.4
2	C	170	GLN	3.2
1	A	252	MET	3.0
1	A	268	ASP	3.0
1	A	293	GLU	2.9
2	C	74	SER	2.9
2	C	69	GLN	2.8
2	C	48	THR	2.8
1	A	264	VAL	2.7
2	C	6	VAL	2.7
1	B	444	SER	2.7
2	C	52	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	436	TYR	2.6
1	A	296	TYR	2.5
1	B	248	LYS	2.5
2	C	27	ARG	2.3
1	A	272	GLU	2.3
2	C	39	TRP	2.3
2	C	54	TYR	2.3
2	C	36	SER	2.3
2	C	68	CYS	2.2
2	C	51	GLN	2.2
1	B	373	TYR	2.1
1	B	237	ASP	2.0
1	B	318	GLU	2.0
1	A	269	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GAL	D	6	11/12	0.52	0.19	125,129,132,137	0
3	NAG	D	1	14/15	0.69	0.19	120,137,152,154	0
3	NAG	D	8	14/15	0.70	0.19	120,126,132,133	0
4	NAG	E	8	14/15	0.73	0.18	102,109,111,112	0
3	NAG	D	5	14/15	0.78	0.18	108,114,120,127	0
3	MAN	D	4	11/12	0.78	0.14	97,106,109,109	0
4	MAN	E	7	11/12	0.79	0.12	73,79,83,94	0
4	FUL	E	9	10/11	0.79	0.16	66,70,73,75	0
3	NAG	D	2	14/15	0.83	0.17	106,116,126,127	0
3	BMA	D	3	11/12	0.84	0.12	101,106,110,113	0
3	MAN	D	7	11/12	0.87	0.10	111,114,122,123	0
4	MAN	E	4	11/12	0.91	0.10	53,59,59,61	0
4	NAG	E	5	14/15	0.92	0.10	56,65,69,70	0
4	NAG	E	1	14/15	0.93	0.11	41,49,58,61	0

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GAL	E	6	11/12	0.94	0.09	56,60,62,62	0
4	NAG	E	2	14/15	0.95	0.08	35,40,41,43	0
4	BMA	E	3	11/12	0.95	0.08	48,49,54,62	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain D:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	C	1001	14/15	0.86	0.12	78,82,87,90	0

6.5 Other polymers [i](#)

There are no such residues in this entry.