



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 08:44 PM UTC

PDB ID : 4WJ7 / pdb_00004wj7
Title : CCM2 PTB domain in complex with KRIT1 NPxY/F3
Authors : Fisher, O.S.; Liu, W.; Zhang, R.; Stiegler, A.L.; Boggon, T.J.
Deposited on : 2014-09-29
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	
1	B	180	
1	C	180	
1	D	180	
2	W	13	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	X	13	8% 69% 23% 8%
2	Y	13	8% 69% 8% 8% 15%
2	Z	13	23% 38% 38% 8% 15%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Malcavernin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	137	Total	C	N	O	S	0	0	0
			1072	683	183	202	4			
1	B	132	Total	C	N	O	S	0	0	0
			1030	656	175	195	4			
1	C	129	Total	C	N	O	S	0	0	0
			1004	640	172	188	4			
1	D	123	Total	C	N	O	S	0	0	0
			952	607	165	176	4			

There are 8 discrepancies between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
A	49	GLY	-	expression tag	UNP Q9BSQ5
A	50	SER	-	expression tag	UNP Q9BSQ5
B	49	GLY	-	expression tag	UNP Q9BSQ5
B	50	SER	-	expression tag	UNP Q9BSQ5
C	49	GLY	-	expression tag	UNP Q9BSQ5
C	50	SER	-	expression tag	UNP Q9BSQ5
D	49	GLY	-	expression tag	UNP Q9BSQ5
D	50	SER	-	expression tag	UNP Q9BSQ5

- Molecule 2 is a protein called KRIT1 NPxY/F3.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	W	11	Total	C	N	O	0	0	0
			88	60	13	15			
2	X	12	Total	C	N	O	0	0	0
			96	66	14	16			
2	Y	11	Total	C	N	O	0	0	0
			88	60	13	15			
2	Z	11	Total	C	N	O	0	0	0
			88	60	13	15			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total 4	O 4	0	0
3	B	6	Total 6	O 6	0	0




● Molecule 2: KRIT1 NPxY/F3




● Molecule 2: KRIT1 NPxY/F3



● Molecule 2: KRIT1 NPxY/F3



● Molecule 2: KRIT1 NPxY/F3



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	110.94Å 110.94Å 315.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.67 – 2.75 29.67 – 2.75	Depositor EDS
% Data completeness (in resolution range)	90.4 (29.67-2.75) 88.2 (29.67-2.75)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.76Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.220 , 0.251 0.221 , 0.253	Depositor DCC
R_{free} test set	1994 reflections (6.48%)	wwPDB-VP
Wilson B-factor (Å ²)	66.0	Xtriage
Anisotropy	0.457	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 58.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4428	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/1091	0.79	1/1483 (0.1%)
1	B	0.30	0/1048	0.80	3/1425 (0.2%)
1	C	0.29	0/1021	0.77	0/1388
1	D	0.40	0/968	0.85	1/1315 (0.1%)
2	W	0.26	0/90	0.56	0/122
2	X	0.26	0/98	0.57	0/133
2	Y	0.26	0/90	0.67	0/122
2	Z	0.35	0/90	0.70	0/122
All	All	0.32	0/4496	0.79	5/6110 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	101	LEU	CA-C-N	7.67	127.83	120.31
1	B	101	LEU	C-N-CA	7.67	127.83	120.31
1	B	103	GLY	N-CA-C	-5.31	103.59	111.03
1	A	105	LEU	N-CA-C	5.14	116.68	108.41
1	D	119	ASN	N-CA-C	5.06	115.99	108.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1072	0	1077	17	0
1	B	1030	0	1030	16	0
1	C	1004	0	1012	18	0
1	D	952	0	967	31	0
2	W	88	0	88	0	0
2	X	96	0	99	2	0
2	Y	88	0	88	2	0
2	Z	88	0	88	7	0
3	A	4	0	0	1	0
3	B	6	0	0	1	0
All	All	4428	0	4449	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:66:GLU:OE2	1:C:125:ARG:NH2	2.20	0.75
1:B:85:ARG:NH2	1:B:146:ARG:O	2.21	0.74
1:D:140:ALA:HB3	1:D:156:LYS:HG3	1.71	0.71
1:A:85:ARG:NH2	1:A:146:ARG:O	2.24	0.71
1:D:106:THR:HG23	1:D:108:GLU:H	1.56	0.70
1:B:126:ASP:OD1	1:B:127:GLY:N	2.26	0.67
1:C:85:ARG:NH2	1:C:146:ARG:O	2.29	0.66
1:C:108:GLU:HG3	1:D:102:PRO:HB3	1.78	0.65
1:A:151:HIS:NE2	3:A:302:HOH:O	2.29	0.65
1:D:119:ASN:HB3	1:D:135:PRO:HA	1.79	0.65
1:D:142:VAL:HG21	1:D:213:LEU:HD13	1.78	0.64
1:C:121:LYS:HD2	1:C:133:ARG:HH21	1.63	0.64
1:A:106:THR:OG1	1:A:107:GLN:N	2.26	0.64
1:A:68:LYS:NZ	1:A:110:ASP:OD1	2.32	0.62
1:B:107:GLN:NE2	1:B:110:ASP:OD2	2.32	0.62
1:A:66:GLU:CD	1:A:125:ARG:HH22	2.07	0.62
1:D:139:ILE:HD11	1:D:217:PHE:HZ	1.65	0.61
1:D:203:LYS:NZ	1:D:207:GLU:OE2	2.33	0.61
1:B:220:VAL:HG13	1:B:221:TYR:H	1.66	0.61
1:D:159:GLN:OE1	1:D:159:GLN:N	2.37	0.58
1:D:115:LEU:HD13	1:D:120:VAL:HG12	1.86	0.58
1:D:121:LYS:NZ	1:D:133:ARG:HB3	2.19	0.58
1:A:107:GLN:N	1:A:107:GLN:OE1	2.37	0.57
2:Z:251:PRO:HG2	2:Z:252:TYR:HD2	1.69	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:155:LEU:HB2	1:D:196:VAL:HB	1.87	0.56
1:B:68:LYS:HB2	1:B:199:ALA:HB3	1.86	0.56
1:D:138:ASP:O	1:D:158:ALA:N	2.22	0.55
1:A:68:LYS:HB2	1:A:199:ALA:HB3	1.90	0.53
1:A:155:LEU:HB2	1:A:196:VAL:HB	1.91	0.53
1:B:137:HIS:HB2	2:X:255:LEU:H	1.74	0.52
1:B:155:LEU:HB2	1:B:196:VAL:HB	1.92	0.52
1:B:191:GLU:N	1:B:191:GLU:OE1	2.42	0.52
1:D:63:ILE:HG21	1:D:212:LEU:HD23	1.92	0.52
1:D:207:GLU:OE2	2:Z:246:LYS:NZ	2.29	0.51
1:A:115:LEU:HD11	1:A:216:VAL:HG21	1.91	0.51
1:D:115:LEU:HD11	1:D:216:VAL:HG21	1.92	0.51
1:C:144:TYR:HB3	2:Y:246:LYS:HB3	1.94	0.50
1:B:156:LYS:NZ	3:B:304:HOH:O	2.29	0.50
2:Z:251:PRO:HG2	2:Z:252:TYR:CD2	2.45	0.50
1:C:219:VAL:HG23	1:C:220:VAL:HG12	1.93	0.49
1:D:139:ILE:HD11	1:D:217:PHE:CZ	2.47	0.49
1:C:216:VAL:O	1:C:219:VAL:HG22	2.12	0.49
1:D:139:ILE:HA	1:D:157:THR:HA	1.95	0.49
1:D:208:GLU:O	1:D:212:LEU:HD12	2.13	0.49
1:D:83:SER:HA	2:Z:247:VAL:HG21	1.95	0.48
1:A:118:TYR:OH	2:X:245:ASP:OD1	2.22	0.48
1:C:114:SER:OG	1:C:121:LYS:HB2	2.14	0.48
1:A:122:LEU:HD12	1:A:132:LEU:HD11	1.96	0.47
1:B:69:TYR:O	1:B:109:HIS:HB3	2.15	0.47
1:B:102:PRO:HB3	1:B:109:HIS:CE1	2.49	0.47
1:D:144:TYR:HB3	2:Z:246:LYS:HB2	1.97	0.47
1:D:68:LYS:HB2	1:D:199:ALA:HB3	1.97	0.46
1:A:113:LEU:HD12	1:A:122:LEU:HD21	1.98	0.46
1:D:145:VAL:HG22	2:Z:244:VAL:HG13	1.96	0.46
1:B:123:ALA:HA	1:B:131:ILE:HG13	1.96	0.46
1:D:121:LYS:HZ2	1:D:133:ARG:HB3	1.79	0.45
1:A:62:TYR:O	1:C:146:ARG:NH2	2.49	0.45
1:D:95:ALA:O	1:D:100:GLN:N	2.45	0.45
1:C:155:LEU:HB2	1:C:196:VAL:HB	1.99	0.45
1:A:119:ASN:C	1:A:119:ASN:HD22	2.24	0.45
1:C:107:GLN:HG3	1:D:108:GLU:OE1	2.17	0.44
1:C:145:VAL:HG22	2:Y:244:VAL:HG22	1.98	0.44
1:D:214:GLY:HA2	2:Z:248:VAL:HG11	1.99	0.44
1:A:124:TRP:O	1:A:128:GLU:HA	2.18	0.44
1:C:214:GLY:O	1:C:218:GLN:HG2	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:113:LEU:HA	1:C:121:LYS:O	2.18	0.43
1:C:113:LEU:HD12	1:C:122:LEU:HD21	1.99	0.43
1:D:63:ILE:HG21	1:D:212:LEU:CD2	2.47	0.43
1:B:220:VAL:HG22	1:B:221:TYR:CD2	2.52	0.43
1:D:211:CYS:O	1:D:215:GLN:HG3	2.18	0.43
1:D:139:ILE:O	1:D:139:ILE:HG13	2.20	0.42
1:C:121:LYS:HD2	1:C:133:ARG:NH2	2.33	0.42
1:C:122:LEU:HD12	1:C:132:LEU:HD11	2.01	0.42
1:C:76:ILE:HG22	1:C:193:CYS:SG	2.61	0.41
1:D:134:VAL:HA	1:D:135:PRO:HD2	1.86	0.41
1:A:110:ASP:O	1:A:125:ARG:HB2	2.20	0.41
1:A:114:SER:HG	1:B:148:ASP:CG	2.28	0.41
1:B:81:ASN:HA	1:B:82:PRO:HD2	1.97	0.41
1:D:109:HIS:HA	1:D:124:TRP:HZ3	1.85	0.41
1:B:69:TYR:CD1	1:B:122:LEU:HD13	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/180 (74%)	128 (96%)	5 (4%)	0	100	100
1	B	128/180 (71%)	121 (94%)	7 (6%)	0	100	100
1	C	125/180 (69%)	122 (98%)	3 (2%)	0	100	100
1	D	119/180 (66%)	117 (98%)	2 (2%)	0	100	100
2	W	9/13 (69%)	9 (100%)	0	0	100	100
2	X	10/13 (77%)	10 (100%)	0	0	100	100
2	Y	9/13 (69%)	9 (100%)	0	0	100	100
2	Z	9/13 (69%)	9 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	542/772 (70%)	525 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/152 (77%)	114 (97%)	3 (3%)	40	63
1	B	112/152 (74%)	108 (96%)	4 (4%)	31	55
1	C	109/152 (72%)	107 (98%)	2 (2%)	51	71
1	D	103/152 (68%)	99 (96%)	4 (4%)	28	51
2	W	10/11 (91%)	9 (90%)	1 (10%)	7	14
2	X	11/11 (100%)	10 (91%)	1 (9%)	9	17
2	Y	10/11 (91%)	9 (90%)	1 (10%)	7	14
2	Z	10/11 (91%)	9 (90%)	1 (10%)	7	14
All	All	482/652 (74%)	465 (96%)	17 (4%)	32	56

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	LEU
1	A	106	THR
1	A	119	ASN
1	B	101	LEU
1	B	119	ASN
1	B	130	ILE
1	B	219	VAL
1	C	107	GLN
1	C	130	ILE
1	D	106	THR
1	D	130	ILE
1	D	139	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	211	CYS
2	W	244	VAL
2	X	244	VAL
2	Y	244	VAL
2	Z	244	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	107	GLN
1	B	137	HIS
1	B	218	GLN
1	C	90	HIS
1	C	109	HIS
1	C	137	HIS
1	D	90	HIS
1	D	109	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	137/180 (76%)	0.25	7 (5%) 33 33	55, 78, 138, 164	0
1	B	132/180 (73%)	0.40	8 (6%) 27 27	51, 80, 145, 175	0
1	C	129/180 (71%)	0.10	6 (4%) 36 36	66, 89, 135, 164	0
1	D	123/180 (68%)	1.01	15 (12%) 8 7	99, 142, 187, 212	0
2	W	11/13 (84%)	0.71	1 (9%) 15 14	61, 72, 95, 130	0
2	X	12/13 (92%)	0.38	1 (8%) 17 18	60, 68, 88, 89	0
2	Y	11/13 (84%)	0.63	1 (9%) 15 14	81, 98, 125, 149	0
2	Z	11/13 (84%)	1.55	3 (27%) 1 1	116, 141, 187, 209	0
All	All	566/772 (73%)	0.46	42 (7%) 20 20	51, 91, 168, 212	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	220	VAL	6.4
1	D	105	LEU	5.4
1	D	130	ILE	4.4
1	D	217	PHE	4.2
2	Y	254	GLY	4.1
1	C	220	VAL	4.0
1	B	221	TYR	3.8
1	A	55	PRO	3.6
1	A	56	ASP	3.6
2	W	254	GLY	3.5
1	B	103	GLY	3.4
2	Z	254	GLY	3.3
1	D	104	HIS	3.2
1	A	128	GLU	3.1
1	C	79	TYR	3.1
1	D	79	TYR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	137	HIS	3.0
1	D	132	LEU	3.0
1	D	142	VAL	2.9
1	A	160	ASP	2.9
1	D	118	TYR	2.8
1	C	159	GLN	2.7
1	C	126	ASP	2.7
1	C	61	ASP	2.7
1	C	219	VAL	2.6
1	D	86	THR	2.6
1	B	96	LYS	2.6
1	B	104	HIS	2.6
1	D	111	ALA	2.5
1	A	221	TYR	2.5
2	X	255	LEU	2.4
1	A	104	HIS	2.4
1	B	111	ALA	2.4
1	B	79	TYR	2.3
2	Z	251	PRO	2.3
2	Z	252	TYR	2.2
1	D	125	ARG	2.2
1	A	58	LEU	2.2
1	D	112	VAL	2.1
1	D	145	VAL	2.1
1	B	110	ASP	2.1
1	D	101	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.