



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:33 AM UTC

PDB ID : 5WJO / pdb_00005wjo
Title : Crystal structure of the unliganded PG90 TCR
Authors : Shahine, A.; Gras, S.; Rossjohn, J.
Deposited on : 2017-07-24
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	9% 75% 17% 7%
1	C	204	6% 79% 17% .
2	B	249	3% 90% 10%
2	D	249	3% 86% 9% 5%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7251 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PG90 TCR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	189	Total	C	N	O	S	12	1	0
			1486	931	255	290	10			
1	C	198	Total	C	N	O	S	15	1	0
			1542	963	264	305	10			

- Molecule 2 is a protein called PG90 TCR beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	248	Total	C	N	O	S	3	1	0
			1954	1229	343	377	5			
2	D	237	Total	C	N	O	S	3	1	0
			1880	1187	329	359	5			

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		
3	D	1	Total	Na	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	4	Total Cl 4 4	0	0
5	C	2	Total Cl 2 2	0	0
5	D	3	Total Cl 3 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	69	Total O 69 69	0	0
6	B	94	Total O 94 94	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	94	Total 94	O 94	0	0
6	D	104	Total 104	O 104	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PG90 TCR alpha chain




- Molecule 1: PG90 TCR alpha chain




- Molecule 2: PG90 TCR beta chain



- Molecule 2: PG90 TCR beta chain





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	41.15Å 128.08Å 169.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.18 – 2.50 39.18 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (39.18-2.50) 99.9 (39.18-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.208 , 0.230 0.219 , 0.243	Depositor DCC
R_{free} test set	1625 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.232	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7251	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/1518	1.13	10/2066 (0.5%)
1	C	0.66	0/1576	1.13	6/2147 (0.3%)
2	B	0.62	0/2004	1.05	2/2726 (0.1%)
2	D	0.61	0/1928	1.04	2/2619 (0.1%)
All	All	0.65	0/7026	1.08	20/9558 (0.2%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	96	GLN	CA-C-N	6.72	130.26	120.71
1	C	96	GLN	C-N-CA	6.72	130.26	120.71
2	D	120	GLU	CB-CG-CD	6.26	123.25	112.60
1	C	162	MET	CA-C-N	5.99	128.59	120.38
1	C	162	MET	C-N-CA	5.99	128.59	120.38
1	A	127	SER	CA-C-N	5.92	131.51	122.93
1	A	127	SER	C-N-CA	5.92	131.51	122.93
1	A	127	SER	N-CA-C	5.78	117.75	109.14
1	A	166	ASP	CA-CB-CG	5.78	118.38	112.60
1	A	53	ASN	CA-CB-CG	5.75	118.35	112.60
1	C	139	ASP	CA-CB-CG	5.69	118.29	112.60
1	A	161	ASP	CA-CB-CG	5.50	118.10	112.60
2	B	68	GLU	CA-C-N	5.39	127.38	122.37
2	B	68	GLU	C-N-CA	5.39	127.38	122.37
1	C	161	ASP	CA-CB-CG	5.34	117.94	112.60
1	A	162	MET	CA-C-N	5.30	127.69	120.54
1	A	162	MET	C-N-CA	5.30	127.69	120.54
1	A	67	ASP	CA-CB-CG	5.08	117.68	112.60
1	A	124	ASP	CA-CB-CG	5.06	117.66	112.60
2	D	191	SER	N-CA-C	5.02	117.18	110.35

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1486	0	1411	15	0
1	C	1542	0	1454	17	0
2	B	1954	0	1854	12	0
2	D	1880	0	1782	12	0
3	A	2	0	0	0	0
3	D	1	0	0	0	0
4	A	8	0	12	1	0
4	C	4	0	6	0	0
4	D	4	0	6	2	0
5	B	4	0	0	0	0
5	C	2	0	0	0	0
5	D	3	0	0	0	0
6	A	69	0	0	1	0
6	B	94	0	0	1	0
6	C	94	0	0	0	0
6	D	104	0	0	2	0
All	All	7251	0	6525	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:ARG:HB2	1:A:47:ILE:HD11	1.62	0.81
1:C:37:ARG:HB2	1:C:47:ILE:HD11	1.69	0.75
1:A:165:MET:HE1	2:B:202:SER:HB3	1.69	0.74
1:A:10:MET:HE1	1:A:21:PRO:HD2	1.70	0.73
1:C:10:MET:HE1	1:C:21:PRO:HD2	1.71	0.71
2:D:51:ASN:O	2:D:69:ARG:HD3	1.94	0.66
2:D:20:ALA:O	4:D:302:EDO:H12	1.96	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:PRO:HG2	1:A:10:MET:HE2	1.79	0.65
2:B:126:PHE:CD2	2:B:192:ARG:HD3	2.32	0.64
1:A:132:VAL:HG12	1:A:175:TRP:HB3	1.80	0.62
1:A:102:THR:HG23	6:A:401:HOH:O	1.98	0.62
1:C:7:PRO:HG2	1:C:10:MET:HE2	1.81	0.62
2:D:8:PRO:HD3	4:D:302:EDO:H11	1.81	0.62
2:D:159:HIS:HB3	2:D:220:TYR:HB2	1.82	0.62
1:A:81[B]:ARG:HH11	4:A:304:EDO:H21	1.65	0.61
2:B:159:HIS:HB3	2:B:220:TYR:HB2	1.82	0.61
1:C:39:ILE:HD13	1:C:84:ALA:HB2	1.82	0.60
1:A:39:ILE:HD13	1:A:84:ALA:HB2	1.82	0.60
1:C:42:GLN:HG2	2:D:112:GLU:OE1	2.04	0.57
1:C:10:MET:HE3	1:C:104:THR:HG21	1.87	0.55
1:A:10:MET:HE3	1:A:104:THR:HG21	1.89	0.53
1:A:80:LEU:HD22	1:A:161:ASP:HB2	1.91	0.51
1:A:37:ARG:HG3	1:A:86:TYR:CE2	2.45	0.51
1:C:32:TYR:HE2	1:C:93:ALA:HB2	1.74	0.51
1:C:80:LEU:HD22	1:C:161:ASP:HB2	1.94	0.50
1:A:165:MET:CE	2:B:202:SER:HB3	2.39	0.49
1:A:184:ALA:HA	1:A:198:PHE:CE1	2.48	0.48
2:B:60:LEU:HD22	2:B:65:PHE:HB3	1.96	0.48
1:C:151:ASP:HB2	1:C:178:LYS:HD3	1.95	0.48
2:D:60:LEU:HD22	2:D:65:PHE:HB3	1.95	0.48
1:A:27:ILE:HG21	1:A:68:ARG:O	2.14	0.47
1:C:6:GLN:HE22	1:C:88:CYS:H	1.61	0.47
1:C:6:GLN:NE2	1:C:88:CYS:H	2.13	0.47
1:C:132:VAL:HG12	1:C:175:TRP:HB3	1.97	0.47
2:D:135:PRO:HD3	2:D:148:LEU:HG	1.97	0.47
2:D:216:GLN:HG3	2:D:239:ILE:HG23	1.97	0.46
2:D:175:VAL:HA	2:D:198:ARG:O	2.15	0.46
2:B:135:PRO:HD3	2:B:148:LEU:HG	1.97	0.46
2:B:216:GLN:HG3	2:B:239:ILE:HG23	1.98	0.45
2:B:30:VAL:HG13	6:B:402:HOH:O	2.18	0.44
2:D:120:GLU:HB2	6:D:401:HOH:O	2.18	0.43
1:C:37:ARG:HG3	1:C:86:TYR:CE2	2.54	0.42
2:B:9:ARG:HG3	2:B:10:TYR:CD1	2.55	0.42
2:D:161:GLU:HG3	2:D:220:TYR:HE1	1.84	0.42
2:B:82[A]:THR:HG22	2:B:118:VAL:CG2	2.50	0.42
1:C:117:PRO:HG3	1:C:193:PRO:HB2	2.02	0.42
2:D:84:GLN:NE2	6:D:401:HOH:O	2.52	0.42
1:C:14:GLU:HB3	1:C:110:PRO:HA	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:4:THR:CG2	1:C:90:VAL:HG13	2.50	0.41
1:A:96:GLN:HB2	2:B:33:PHE:HZ	1.86	0.41
2:B:36:GLN:HB2	2:B:46:LEU:HD11	2.03	0.40
1:C:200:SER:HA	1:C:201:PRO:HD3	1.90	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/204 (90%)	177 (96%)	7 (4%)	0	100	100
1	C	195/204 (96%)	189 (97%)	6 (3%)	0	100	100
2	B	247/249 (99%)	239 (97%)	8 (3%)	0	100	100
2	D	232/249 (93%)	227 (98%)	5 (2%)	0	100	100
All	All	858/906 (95%)	832 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/184 (91%)	156 (93%)	11 (7%)	15	32
1	C	173/184 (94%)	168 (97%)	5 (3%)	37	65

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	209/210 (100%)	205 (98%)	4 (2%)	50	76
2	D	202/210 (96%)	197 (98%)	5 (2%)	42	69
All	All	751/788 (95%)	726 (97%)	25 (3%)	33	61

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	SER
1	A	27	ILE
1	A	30	ASN
1	A	51	LEU
1	A	130	LYS
1	A	158	CYS
1	A	159	VAL
1	A	164	SER
1	A	177	ASN
1	A	189	ASN
1	A	190	SER
2	B	100	GLN
2	B	104	ASN
2	B	161	GLU
2	B	191	SER
1	C	27	ILE
1	C	51	LEU
1	C	56	THR
1	C	159	VAL
1	C	192	ILE
2	D	52	GLU
2	D	96	LEU
2	D	123	LYS
2	D	141	SER
2	D	204	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	111	ASN
1	A	121	GLN
2	B	36	GLN
2	B	54	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	142	HIS
2	B	238	GLN
1	C	6	GLN
1	C	107	GLN
1	C	111	ASN
1	C	185	ASN
1	C	188	ASN
2	D	36	GLN
2	D	50	GLN
2	D	54	GLN
2	D	212	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 12 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	D	302	-	3,3,3	0.49	0	2,2,2	0.33	0
4	EDO	A	304	-	3,3,3	0.44	0	2,2,2	0.24	0
4	EDO	C	301	-	3,3,3	0.42	0	2,2,2	0.50	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	303	-	3,3,3	0.43	0	2,2,2	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	D	302	-	-	0/1/1/1	-
4	EDO	A	304	-	-	0/1/1/1	-
4	EDO	C	301	-	-	0/1/1/1	-
4	EDO	A	303	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	302	EDO	2	0
4	A	304	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	189/204 (92%)	0.60	19 (10%) 12 10	17, 38, 82, 101	3 (1%)
1	C	198/204 (97%)	0.38	13 (6%) 24 21	14, 33, 68, 88	4 (2%)
2	B	248/249 (99%)	0.23	7 (2%) 55 50	16, 33, 66, 90	2 (0%)
2	D	237/249 (95%)	0.11	8 (3%) 48 43	14, 29, 66, 89	2 (0%)
All	All	872/906 (96%)	0.31	47 (5%) 31 27	14, 33, 71, 101	11 (1%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	201	PRO	4.8
1	A	187	PHE	3.4
2	B	189	ASN	3.3
1	C	94	TYR	3.2
2	B	2	ALA	3.2
1	A	94	TYR	3.2
1	A	192	ILE	3.2
1	C	96	GLN	3.1
1	C	92	VAL	3.1
1	C	95	ARG	3.1
2	B	249	ASP	3.0
1	C	127	SER	2.9
1	A	95	ARG	2.9
1	A	199	PRO	2.9
2	D	186	PRO	2.8
2	D	230	GLN	2.8
2	B	188	LEU	2.7
1	A	198	PHE	2.7
2	D	97	ALA	2.7
2	D	96	LEU	2.6
2	D	229	THR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	180	ASP	2.5
1	A	96	GLN	2.5
1	A	191	ILE	2.5
1	A	53	ASN	2.5
1	C	41	SER	2.4
2	D	41[A]	GLN	2.4
1	A	194	GLU	2.4
1	A	146	GLN	2.4
1	C	147	SER	2.3
2	B	97	ALA	2.3
1	A	188	ASN	2.3
2	B	187	ALA	2.3
1	A	145	SER	2.3
1	C	200	SER	2.3
1	C	130	LYS	2.3
2	D	249	ASP	2.2
1	C	152	VAL	2.2
1	A	196	THR	2.2
1	A	177	ASN	2.2
2	B	190	ASP	2.2
1	A	197	PHE	2.2
1	A	93	ALA	2.1
1	A	129	ASP	2.1
1	A	185	ASN	2.1
2	D	59	GLY	2.0
1	C	93	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	D	302	4/4	0.80	0.14	42,42,43,43	0
3	NA	A	302	1/1	0.81	0.21	61,61,61,61	0
5	CL	B	301	1/1	0.84	0.13	52,52,52,52	0
5	CL	C	303	1/1	0.85	0.10	58,58,58,58	0
5	CL	C	302	1/1	0.86	0.11	53,53,53,53	0
4	EDO	A	304	4/4	0.86	0.16	44,44,44,44	0
5	CL	D	304	1/1	0.86	0.11	41,41,41,41	0
5	CL	D	305	1/1	0.86	0.15	49,49,49,49	0
5	CL	D	303	1/1	0.90	0.12	52,52,52,52	0
3	NA	D	301	1/1	0.92	0.10	34,34,34,34	0
5	CL	B	302	1/1	0.93	0.13	58,58,58,58	0
4	EDO	A	303	4/4	0.93	0.13	26,27,30,31	0
5	CL	B	303	1/1	0.94	0.11	49,49,49,49	0
5	CL	B	304	1/1	0.95	0.09	66,66,66,66	0
3	NA	A	301	1/1	0.96	0.11	31,31,31,31	0
4	EDO	C	301	4/4	0.96	0.08	22,22,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.