



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 12:13 AM UTC

PDB ID : 1WKR / pdb_00001wkr
Title : Crystal structure of aspartic proteinase from *Irpex lacteus*
Authors : Fujimoto, Z.; Fujii, Y.; Kaneko, S.; Kobayashi, H.; Mizuno, H.
Deposited on : 2004-06-02
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1553 (1.30-1.30)
Clashscore	190562	1595 (1.30-1.30)
Ramachandran outliers	187476	1551 (1.30-1.30)
Sidechain outliers	187428	1551 (1.30-1.30)
RSRZ outliers	180081	1549 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	2% 85% 13% .
2	I	6	67% 17% 17%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3257 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyporopepsin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	340	Total	C	N	O	0	0	0
			2472	1545	396	531			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	194	SER	THR	SEE REMARK 999	UNP P17576

- Molecule 2 is a protein called pepstatin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	6	Total	C	N	O	0	0	0
			48	34	5	9			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	693	Total O 693 693	0	0
4	I	9	Total O 9 9	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Polyporopepsin



• Molecule 2: pepstatin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	37.27Å 78.90Å 54.07Å 90.00° 96.71° 90.00°	Depositor
Resolution (Å)	26.99 – 1.30 26.99 – 1.30	Depositor EDS
% Data completeness (in resolution range)	98.8 (26.99-1.30) 99.0 (26.99-1.30)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.46 (at 1.30Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.147 , 0.171 0.155 , 0.178	Depositor DCC
R_{free} test set	3791 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	7.0	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3257	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, STA, IVA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.96	70/2516 (2.8%)	1.37	23/3443 (0.7%)
2	I	0.98	0/17	0.93	0/21
All	All	1.95	70/2533 (2.8%)	1.37	23/3464 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
2	I	0	2
All	All	0	6

All (70) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	94	ARG	NE-CZ	13.42	1.47	1.33
1	A	285	SER	CB-OG	11.41	1.65	1.42
1	A	285	SER	CA-CB	-11.01	1.36	1.53
1	A	238	ASN	CA-C	-10.20	1.40	1.52
1	A	94	ARG	CZ-NH1	10.10	1.46	1.32
1	A	54	ALA	C-N	9.68	1.49	1.33
1	A	152	SER	CA-C	-9.65	1.40	1.52
1	A	85	PRO	C-N	-9.11	1.21	1.33
1	A	196	ARG	CB-CG	-8.55	1.26	1.52
1	A	121	THR	CB-OG1	-8.46	1.30	1.43
1	A	120	HIS	ND1-CE1	8.44	1.41	1.32
1	A	85	PRO	CA-C	8.23	1.63	1.52
1	A	238	ASN	C-O	7.93	1.33	1.24
1	A	283	GLY	C-O	-7.93	1.13	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	229	LYS	CB-CG	-7.84	1.28	1.52
1	A	80	GLY	C-O	7.67	1.35	1.24
1	A	244	ARG	CD-NE	7.54	1.56	1.46
1	A	167	SER	CA-CB	7.52	1.66	1.53
1	A	167	SER	CB-OG	-7.48	1.27	1.42
1	A	54	ALA	CA-C	-7.44	1.43	1.52
1	A	239	ASN	CA-C	7.28	1.62	1.52
1	A	285	SER	N-CA	7.25	1.54	1.45
1	A	154	SER	C-O	7.23	1.32	1.23
1	A	86	LYS	CE-NZ	7.17	1.70	1.49
1	A	280	THR	CB-OG1	-7.08	1.32	1.43
1	A	196	ARG	CG-CD	6.83	1.73	1.52
1	A	152	SER	C-N	6.81	1.43	1.34
1	A	154	SER	CA-CB	6.76	1.61	1.52
1	A	236	ALA	C-O	6.69	1.31	1.23
1	A	329	LEU	CB-CG	-6.56	1.40	1.53
1	A	120	HIS	CG-CD2	6.52	1.43	1.35
1	A	263	GLN	CD-OE1	6.48	1.35	1.23
1	A	236	ALA	CA-CB	6.42	1.64	1.53
1	A	236	ALA	CA-C	-6.40	1.44	1.52
1	A	93	SER	N-CA	6.38	1.54	1.46
1	A	17	ASN	CG-ND2	-6.24	1.20	1.33
1	A	152	SER	N-CA	6.14	1.54	1.45
1	A	21	GLY	N-CA	6.02	1.52	1.45
1	A	137	THR	C-O	6.01	1.31	1.24
1	A	64	GLY	C-O	6.00	1.31	1.23
1	A	199	SER	CB-OG	5.95	1.54	1.42
1	A	190	GLY	CA-C	-5.95	1.48	1.52
1	A	55	THR	C-O	-5.86	1.15	1.23
1	A	285	SER	C-O	5.85	1.31	1.23
1	A	17	ASN	CG-OD1	5.81	1.34	1.23
1	A	153	GLU	CA-C	-5.70	1.45	1.52
1	A	196	ARG	NE-CZ	-5.66	1.26	1.33
1	A	287	SER	C-N	-5.66	1.25	1.33
1	A	284	GLY	CA-C	5.64	1.61	1.52
1	A	263	GLN	CD-NE2	-5.63	1.21	1.33
1	A	277	ASN	CA-C	-5.59	1.45	1.52
1	A	165	THR	C-O	5.48	1.30	1.23
1	A	122	SER	CA-CB	5.45	1.63	1.53
1	A	284	GLY	C-N	-5.41	1.25	1.33
1	A	238	ASN	CA-CB	5.39	1.61	1.53
1	A	86	LYS	N-CA	-5.33	1.38	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	83	THR	C-O	5.31	1.30	1.23
1	A	242	LEU	C-O	5.23	1.30	1.23
1	A	120	HIS	CG-ND1	-5.20	1.32	1.38
1	A	47	VAL	CA-CB	-5.18	1.48	1.53
1	A	192	ASN	CG-ND2	-5.17	1.22	1.33
1	A	42	ALA	C-O	5.14	1.30	1.24
1	A	123	THR	CA-C	5.14	1.59	1.52
1	A	121	THR	C-N	-5.12	1.26	1.33
1	A	151	THR	N-CA	5.11	1.51	1.46
1	A	192	ASN	CG-OD1	5.11	1.33	1.23
1	A	237	ASP	C-N	5.11	1.40	1.33
1	A	58	LYS	CD-CE	5.11	1.67	1.52
1	A	81	SER	CA-C	-5.09	1.44	1.52
1	A	153	GLU	CG-CD	-5.09	1.39	1.52

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	196	ARG	NE-CZ-NH2	10.78	128.91	119.20
1	A	238	ASN	O-C-N	-9.38	112.27	122.03
1	A	284	GLY	O-C-N	8.83	134.50	122.84
1	A	152	SER	CA-C-O	8.40	131.33	121.58
1	A	254	GLN	OE1-CD-NE2	-7.09	115.51	122.60
1	A	238	ASN	CA-C-N	6.68	129.90	120.28
1	A	238	ASN	C-N-CA	6.68	129.90	120.28
1	A	120	HIS	CG-CD2-NE2	-6.65	100.55	107.20
1	A	254	GLN	CB-CG-CD	-6.58	101.42	112.60
1	A	120	HIS	CE1-NE2-CD2	6.48	115.48	109.00
1	A	277	ASN	OD1-CG-ND2	-6.27	116.33	122.60
1	A	154	SER	O-C-N	-6.09	116.52	123.40
1	A	152	SER	O-C-N	-5.88	114.88	122.94
1	A	239	ASN	N-CA-C	-5.75	105.10	111.71
1	A	98	PHE	CA-CB-CG	-5.61	108.19	113.80
1	A	94	ARG	CD-NE-CZ	-5.59	116.57	124.40
1	A	236	ALA	O-C-N	-5.55	116.05	122.87
1	A	284	GLY	CA-C-N	-5.45	113.32	122.92
1	A	284	GLY	C-N-CA	-5.45	113.32	122.92
1	A	17	ASN	CB-CG-ND2	5.38	124.47	116.40
1	A	120	HIS	ND1-CE1-NE2	-5.36	103.04	108.40
1	A	285	SER	CA-CB-OG	5.11	121.31	111.10
1	A	151	THR	N-CA-C	-5.05	105.44	112.30

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	154	SER	Mainchain
1	A	238	ASN	Mainchain
1	A	54	ALA	Mainchain
1	A	94	ARG	Sidechain
2	I	804	STA	Peptide,Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2472	0	2408	10	0
2	I	48	0	60	0	0
3	A	35	0	0	0	0
4	A	693	0	0	4	5
4	I	9	0	0	0	0
All	All	3257	0	2468	10	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:LYS:NZ	1:A:86:LYS:CE	1.70	1.53
1:A:285:SER:OG	1:A:285:SER:CB	1.65	1.44
1:A:78:THR:HG23	4:A:1595:HOH:O	1.79	0.83
1:A:176:TYR:OH	4:A:1546:HOH:O	2.01	0.78
1:A:293:VAL:HG23	4:A:1513:HOH:O	1.91	0.71
1:A:285:SER:CB	1:A:285:SER:HG	2.08	0.60
1:A:78:THR:CG2	4:A:1595:HOH:O	2.45	0.59
1:A:86:LYS:NZ	1:A:86:LYS:CD	2.62	0.57
1:A:152:SER:HB2	1:A:155:SER:OG	2.12	0.49
1:A:310:LEU:HD23	1:A:310:LEU:C	2.44	0.42

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1431:HOH:O	4:A:1574:HOH:O[1_455]	2.02	0.18
4:A:1033:HOH:O	4:A:1504:HOH:O[1_455]	2.12	0.08
4:A:1082:HOH:O	4:A:1504:HOH:O[1_455]	2.15	0.05
4:A:1189:HOH:O	4:A:1579:HOH:O[1_554]	2.19	0.01
4:A:1445:HOH:O	4:A:1576:HOH:O[2_546]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/340 (99%)	332 (98%)	6 (2%)	0	100	100
2	I	3/6 (50%)	2 (67%)	1 (33%)	0	100	100
All	All	341/346 (99%)	334 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/278 (100%)	278 (100%)	0	100	100
2	I	2/2 (100%)	2 (100%)	0	100	100
All	All	280/280 (100%)	280 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	277	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	STA	I	806	2	11,11,11	1.93	3 (27%)	11,14,14	0.78	0
2	STA	I	804	2	10,10,11	1.12	1 (10%)	9,12,14	1.12	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	STA	I	806	2	-	2/12/12/12	-
2	STA	I	804	2	-	1/11/11/12	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	806	STA	O-C	4.79	1.37	1.22
2	I	804	STA	CH-CA	-2.57	1.50	1.53
2	I	806	STA	OH-CH	-2.53	1.38	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	806	STA	CM-C	2.18	1.56	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	804	STA	CH-CM-C	-2.57	108.63	113.02

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	804	STA	O-C-CM-CH
2	I	806	STA	CA-CB-CG-CD2
2	I	806	STA	CA-CB-CG-CD1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	906	-	4,4,4	1.79	2 (50%)	6,6,6	1.00	1 (16%)
3	SO4	A	904	-	4,4,4	1.53	1 (25%)	6,6,6	0.80	0
3	SO4	A	903	-	4,4,4	0.39	0	6,6,6	0.66	0
3	SO4	A	902	-	4,4,4	1.94	1 (25%)	6,6,6	0.54	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	907	-	4,4,4	2.72	2 (50%)	6,6,6	0.91	0
3	SO4	A	905	-	4,4,4	2.96	3 (75%)	6,6,6	1.06	0
3	SO4	A	901	-	4,4,4	0.72	0	6,6,6	0.60	0

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	905	SO4	O1-S	3.98	1.68	1.44
3	A	907	SO4	O2-S	3.85	1.67	1.44
3	A	905	SO4	O2-S	3.79	1.67	1.44
3	A	902	SO4	O1-S	3.62	1.66	1.44
3	A	907	SO4	O1-S	3.51	1.65	1.44
3	A	906	SO4	O3-S	2.71	1.70	1.48
3	A	904	SO4	O1-S	2.40	1.59	1.44
3	A	906	SO4	O4-S	2.19	1.66	1.48
3	A	905	SO4	O4-S	2.13	1.65	1.48

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	906	SO4	O4-S-O3	-2.02	97.39	108.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/340 (100%)	-0.37	6 (1%) 67 72	3, 6, 13, 23	10 (2%)
2	I	3/6 (50%)	-0.50	0 100 100	5, 5, 6, 6	0
All	All	343/346 (99%)	-0.37	6 (1%) 69 73	3, 6, 13, 23	10 (2%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	238	ASN	3.6
1	A	99	ASP	3.1
1	A	287	SER	2.3
1	A	154	SER	2.3
1	A	54	ALA	2.2
1	A	239	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	STA	I	806	12/12	0.92	0.08	7,11,14,14	0
2	STA	I	804	11/12	0.97	0.05	3,4,5,6	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	905	5/5	0.76	0.14	32,32,35,36	0
3	SO4	A	907	5/5	0.83	0.15	27,28,31,35	0
3	SO4	A	906	5/5	0.89	0.12	24,33,37,43	0
3	SO4	A	901	5/5	0.89	0.19	9,13,15,17	0
3	SO4	A	904	5/5	0.93	0.09	27,27,29,32	0
3	SO4	A	902	5/5	0.93	0.12	15,19,24,27	0
3	SO4	A	903	5/5	0.97	0.11	11,12,19,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.