



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:05 PM UTC

PDB ID : 5WK4 / pdb_00005wk4
Title : Crystal structure of an anti-idiotypic VLR
Authors : Collins, B.C.; Nakahara, H.; Cooper, M.D.; Herrin, B.R.; Wilson, I.A.
Deposited on : 2017-07-24
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	201	
1	B	201	
1	C	201	
1	D	201	
1	E	201	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	201	9% 85% 10% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Variable lymphocyte receptor 39.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	192	Total	C	N	O	S	0	0	0
			1459	918	259	274	8			
1	A	190	Total	C	N	O	S	0	0	0
			1443	911	254	270	8			
1	B	192	Total	C	N	O	S	0	0	0
			1452	915	259	270	8			
1	C	192	Total	C	N	O	S	0	0	0
			1447	911	256	272	8			
1	E	191	Total	C	N	O	S	0	0	0
			1438	910	250	270	8			
1	F	191	Total	C	N	O	S	0	0	0
			1426	903	250	265	8			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		
2	E	1	Total	Mg	0	0
			1	1		
2	F	1	Total	Mg	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	175	Total	O	0	0
			175	175		
3	A	182	Total	O	0	0
			182	182		

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	180	Total 180	O 180	0	0
3	C	140	Total 140	O 140	0	0
3	E	142	Total 142	O 142	0	0
3	F	124	Total 124	O 124	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Variable lymphocyte receptor 39




- Molecule 1: Variable lymphocyte receptor 39




- Molecule 1: Variable lymphocyte receptor 39





- Molecule 1: Variable lymphocyte receptor 39



- Molecule 1: Variable lymphocyte receptor 39



- Molecule 1: Variable lymphocyte receptor 39



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.41Å 61.38Å 80.56Å 110.09° 99.79° 93.52°	Depositor
Resolution (Å)	34.72 – 1.50 34.72 – 1.50	Depositor EDS
% Data completeness (in resolution range)	85.1 (34.72-1.50) 85.1 (34.72-1.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 1.50Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.193 , 0.223 (Not available) , 0.237	Depositor DCC
R_{free} test set	7060 reflections (4.13%)	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 25.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9612	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/1477	0.51	0/2021
1	B	0.28	0/1486	0.50	0/2032
1	C	0.27	0/1481	0.48	0/2027
1	D	0.30	0/1493	0.51	0/2041
1	E	0.26	0/1473	0.47	0/2019
1	F	0.24	0/1461	0.44	0/2004
All	All	0.27	0/8871	0.49	0/12144

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1443	0	1417	9	0
1	B	1452	0	1428	4	0
1	C	1447	0	1412	6	0
1	D	1459	0	1434	11	0
1	E	1438	0	1397	10	0
1	F	1426	0	1384	12	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	182	0	0	2	0
3	B	180	0	0	1	0
3	C	140	0	0	2	0
3	D	175	0	0	0	0
3	E	142	0	0	3	0
3	F	124	0	0	1	0
All	All	9612	0	8472	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:108:GLN:NE2	3:E:402:HOH:O	2.21	0.73
1:C:110:THR:O	3:C:301:HOH:O	2.11	0.67
1:A:136:ILE:HG12	1:A:167:LEU:HB2	1.76	0.67
1:E:114:ALA:O	3:E:401:HOH:O	2.13	0.66
1:A:49:THR:HG22	1:A:71:SER:HB2	1.79	0.65
1:D:58:ASN:ND2	1:D:82:ASP:OD2	2.30	0.64
1:B:207:SER:HB3	1:B:210:LYS:HD2	1.78	0.64
1:B:138:ARG:NH2	3:B:303:HOH:O	2.31	0.62
1:F:58:ASN:ND2	1:F:82:ASP:OD2	2.32	0.62
1:F:48:PRO:O	1:F:51:ARG:HG2	2.04	0.58
1:A:210:LYS:NZ	3:A:403:HOH:O	2.37	0.57
1:A:80:VAL:HG22	1:A:104:LEU:HD23	1.88	0.56
1:D:175:ALA:O	1:D:178:VAL:HG12	2.06	0.55
1:E:39:SER:HB2	1:F:38:ARG:HG2	1.91	0.53
1:F:48:PRO:HB2	1:F:51:ARG:HD3	1.91	0.52
1:D:170:TRP:CH2	1:D:177:ILE:HD11	2.44	0.52
1:E:89:PRO:HG2	1:E:92:VAL:HB	1.92	0.52
1:E:160:GLU:HB3	1:E:210:LYS:HD3	1.93	0.51
1:F:136:ILE:HG12	1:F:167:LEU:HB2	1.93	0.50
1:A:136:ILE:HD13	1:A:167:LEU:HD13	1.94	0.50
1:C:89:PRO:HG2	1:C:92:VAL:HB	1.95	0.48
1:F:28:CYS:SG	1:F:33:VAL:HG22	2.53	0.48
1:E:136:ILE:HG12	1:E:167:LEU:HB2	1.96	0.48
1:D:136:ILE:HD13	1:D:141:PHE:CE2	2.49	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:ARG:NH1	3:A:408:HOH:O	2.47	0.47
1:F:33:VAL:CG2	1:F:51:ARG:HG3	2.44	0.47
1:D:80:VAL:HG22	1:D:104:LEU:HD23	1.97	0.46
1:A:72:LEU:HB3	1:A:75:LEU:HG	1.97	0.46
1:C:95:ARG:NH1	3:C:303:HOH:O	2.40	0.45
1:F:210:LYS:HB3	1:F:210:LYS:HE2	1.72	0.45
1:D:80:VAL:CG2	1:D:104:LEU:HD23	2.46	0.45
1:E:210:LYS:HE2	1:E:210:LYS:HB3	1.76	0.43
1:E:68:VAL:HG11	3:E:511:HOH:O	2.17	0.43
1:B:65:GLU:HB2	1:B:68:VAL:HG13	1.99	0.43
1:C:175:ALA:O	1:C:178:VAL:HG12	2.19	0.43
1:D:136:ILE:CG1	1:D:167:LEU:HB2	2.49	0.42
1:F:100:THR:HA	1:F:123:LEU:HA	2.01	0.42
1:D:136:ILE:HD12	1:D:166:TYR:CE2	2.55	0.42
1:D:55:TRP:CZ2	1:E:29:ARG:HG2	2.55	0.42
1:A:159:CYS:SG	1:A:198:VAL:HG22	2.60	0.42
1:A:170:TRP:CH2	1:A:177:ILE:HD11	2.55	0.42
1:E:40:LEU:HD13	1:E:44:PRO:HG3	2.02	0.42
1:F:25:GLN:NE2	3:F:408:HOH:O	2.53	0.42
1:C:80:VAL:CG2	1:C:104:LEU:HD23	2.50	0.42
1:D:72:LEU:HB3	1:D:75:LEU:HG	2.02	0.41
1:D:136:ILE:HG12	1:D:167:LEU:HB2	2.02	0.41
1:F:56:LEU:HB2	1:F:80:VAL:HG12	2.03	0.40
1:B:80:VAL:CG2	1:B:104:LEU:HD23	2.52	0.40
1:C:178:VAL:HG11	1:C:186:VAL:HG23	2.02	0.40
1:F:33:VAL:HG23	1:F:51:ARG:HG3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/201 (94%)	181 (96%)	7 (4%)	0	100	100
1	B	190/201 (94%)	183 (96%)	7 (4%)	0	100	100
1	C	190/201 (94%)	182 (96%)	8 (4%)	0	100	100
1	D	190/201 (94%)	183 (96%)	7 (4%)	0	100	100
1	E	189/201 (94%)	181 (96%)	8 (4%)	0	100	100
1	F	189/201 (94%)	182 (96%)	7 (4%)	0	100	100
All	All	1136/1206 (94%)	1092 (96%)	44 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/176 (93%)	160 (98%)	4 (2%)	43	15
1	B	164/176 (93%)	160 (98%)	4 (2%)	43	15
1	C	163/176 (93%)	160 (98%)	3 (2%)	51	24
1	D	166/176 (94%)	162 (98%)	4 (2%)	43	15
1	E	161/176 (92%)	156 (97%)	5 (3%)	35	9
1	F	159/176 (90%)	152 (96%)	7 (4%)	25	4
All	All	977/1056 (92%)	950 (97%)	27 (3%)	38	11

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	60	GLN
1	D	99	LEU
1	D	108	GLN
1	D	207	SER
1	A	33	VAL
1	A	99	LEU
1	A	168	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	210	LYS
1	B	99	LEU
1	B	147	LEU
1	B	178	VAL
1	B	210	LYS
1	C	99	LEU
1	C	147	LEU
1	C	210	LYS
1	E	70	ASP
1	E	99	LEU
1	E	147	LEU
1	E	178	VAL
1	E	210	LYS
1	F	60	GLN
1	F	84	GLN
1	F	90	ILE
1	F	99	LEU
1	F	108	GLN
1	F	178	VAL
1	F	210	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	58	ASN
1	D	143	ASN
1	A	58	ASN
1	A	143	ASN
1	B	106	HIS
1	B	132	GLN
1	C	52	GLN
1	E	174	HIS
1	F	34	ASN
1	F	154	ASN
1	F	174	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	190/201 (94%)	0.31	7 (3%)	45 49	13, 20, 34, 52	0
1	B	192/201 (95%)	0.28	5 (2%)	57 61	13, 20, 32, 40	0
1	C	192/201 (95%)	0.44	6 (3%)	51 55	14, 22, 33, 44	0
1	D	192/201 (95%)	0.22	7 (3%)	46 50	14, 19, 31, 41	0
1	E	191/201 (95%)	0.51	6 (3%)	51 55	17, 24, 32, 47	0
1	F	191/201 (95%)	0.88	18 (9%)	14 14	16, 27, 39, 48	0
All	All	1148/1206 (95%)	0.44	49 (4%)	40 44	13, 22, 35, 52	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	136	ILE	4.9
1	E	136	ILE	4.3
1	D	211	CYS	3.7
1	E	212	PRO	3.6
1	A	211	CYS	3.6
1	F	114	ALA	3.5
1	C	211	CYS	3.5
1	D	114	ALA	3.4
1	C	20	GLY	3.3
1	F	136	ILE	3.2
1	A	209	SER	3.2
1	F	22	CYS	3.1
1	B	211	CYS	3.1
1	F	212	PRO	2.8
1	A	22	CYS	2.7
1	B	20	GLY	2.7
1	A	136	ILE	2.7
1	A	159	CYS	2.7
1	D	20	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	46	GLY	2.7
1	D	82	ASP	2.6
1	F	195	ASN	2.6
1	E	114	ALA	2.6
1	F	193	GLY	2.5
1	F	194	THR	2.5
1	C	68	VAL	2.5
1	F	165	LEU	2.4
1	B	114	ALA	2.4
1	F	90	ILE	2.4
1	E	115	GLY	2.4
1	A	77	PHE	2.3
1	F	206	THR	2.3
1	D	204	ALA	2.3
1	F	178	VAL	2.3
1	F	201	VAL	2.3
1	F	167	LEU	2.3
1	F	204	ALA	2.2
1	F	94	ASP	2.2
1	C	90	ILE	2.2
1	C	58	ASN	2.2
1	B	68	VAL	2.2
1	B	209	SER	2.1
1	D	58	ASN	2.1
1	F	39	SER	2.1
1	E	211	CYS	2.1
1	C	209	SER	2.1
1	A	210	LYS	2.0
1	F	113	PRO	2.0
1	E	28	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	F	300	1/1	0.80	0.15	46,46,46,46	0
2	MG	A	300	1/1	0.85	0.10	35,35,35,35	0
2	MG	D	300	1/1	0.90	0.12	36,36,36,36	0
2	MG	E	300	1/1	0.98	0.09	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.