



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 12:10 PM UTC

PDB ID : 5WKH / pdb_00005wkh
Title : D30 TCR in complex with HLA-A*11:01-GTS3
Authors : Gras, S.; Rossjohn, J.
Deposited on : 2017-07-25
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	85% 14%
1	F	274	81% 17%
2	B	100	77% 19%
2	G	100	82% 17%
3	C	10	80% 20%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	10	70%30%
4	D	198	5% 72%23%5% •
4	I	198	4% 67%26%6% •
5	E	244	76%23% •
5	J	244	% 80%17% •

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 13218 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-11 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	S	0	0	0
			2235	1389	407	430	9			
1	F	274	Total	C	N	O	S	0	0	0
			2235	1389	407	430	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			829	528	140	158	3			
2	G	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
G	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called GTS3 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	10	Total	C	N	O	0	0	0
			70	41	14	15			
3	H	10	Total	C	N	O	0	0	0
			70	41	14	15			

- Molecule 4 is a protein called T-cell receptor alpha variable 30,T-cell receptor, sp3.4 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	198	Total 1535	C 961	N 259	O 305	S 10	0	0	0
4	I	198	Total 1535	C 961	N 259	O 305	S 10	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	51	ILE	VAL	engineered mutation	UNP A0A087WSZ9
D	102	LEU	-	linker	UNP A0A087WSZ9
D	103	GLY	-	linker	UNP A0A087WSZ9
D	105	ASP	-	linker	UNP A0A087WSZ9
D	106	ALA	-	linker	UNP A0A087WSZ9
D	107	GLY	-	linker	UNP A0A087WSZ9
D	108	ASN	-	linker	UNP A0A087WSZ9
D	109	MET	-	linker	UNP A0A087WSZ9
D	110	LEU	-	linker	UNP A0A087WSZ9
D	111	THR	-	linker	UNP A0A087WSZ9
D	112	PHE	-	linker	UNP A0A087WSZ9
D	113	GLY	-	linker	UNP A0A087WSZ9
D	114	GLY	-	linker	UNP A0A087WSZ9
D	115	GLY	-	linker	UNP A0A087WSZ9
D	116	THR	-	linker	UNP A0A087WSZ9
D	117	ARG	-	linker	UNP A0A087WSZ9
D	118	LEU	-	linker	UNP A0A087WSZ9
D	119	MET	-	linker	UNP A0A087WSZ9
D	120	VAL	-	linker	UNP A0A087WSZ9
D	121	LYS	-	linker	UNP A0A087WSZ9
D	122	PRO	-	linker	UNP A0A087WSZ9
D	123	HIS	-	linker	UNP A0A087WSZ9
I	51	ILE	VAL	engineered mutation	UNP A0A087WSZ9
I	102	LEU	-	linker	UNP A0A087WSZ9
I	103	GLY	-	linker	UNP A0A087WSZ9
I	105	ASP	-	linker	UNP A0A087WSZ9
I	106	ALA	-	linker	UNP A0A087WSZ9
I	107	GLY	-	linker	UNP A0A087WSZ9
I	108	ASN	-	linker	UNP A0A087WSZ9
I	109	MET	-	linker	UNP A0A087WSZ9
I	110	LEU	-	linker	UNP A0A087WSZ9
I	111	THR	-	linker	UNP A0A087WSZ9
I	112	PHE	-	linker	UNP A0A087WSZ9
I	113	GLY	-	linker	UNP A0A087WSZ9
I	114	GLY	-	linker	UNP A0A087WSZ9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	115	GLY	-	linker	UNP A0A087WSZ9
I	116	THR	-	linker	UNP A0A087WSZ9
I	117	ARG	-	linker	UNP A0A087WSZ9
I	118	LEU	-	linker	UNP A0A087WSZ9
I	119	MET	-	linker	UNP A0A087WSZ9
I	120	VAL	-	linker	UNP A0A087WSZ9
I	121	LYS	-	linker	UNP A0A087WSZ9
I	122	PRO	-	linker	UNP A0A087WSZ9
I	123	HIS	-	linker	UNP A0A087WSZ9

- Molecule 5 is a protein called D30 TCR beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	244	Total	C	N	O	S	0	0	0
			1935	1229	336	365	5			
5	J	244	Total	C	N	O	S	0	0	0
			1935	1229	336	365	5			


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	E	1	Total	O	0	0
			1	1		
6	F	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, A-11 alpha chain




- Molecule 1: HLA class I histocompatibility antigen, A-11 alpha chain




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: GTS3 peptide





- Molecule 3: GTS3 peptide



- Molecule 4: T-cell receptor alpha variable 30,T-cell receptor, sp3.4 alpha chain




- Molecule 4: T-cell receptor alpha variable 30,T-cell receptor, sp3.4 alpha chain




- Molecule 5: D30 TCR beta chain



- Molecule 5: D30 TCR beta chain





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	77.39Å 170.54Å 85.69Å 90.00° 113.89° 90.00°	Depositor
Resolution (Å)	46.01 – 3.20 46.01 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.01-3.20) 99.8 (46.01-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 3.19Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.177 , 0.246 (Not available) , 0.229	Depositor DCC
R_{free} test set	1663 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	38.6	Xtriage
Anisotropy	0.683	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.030 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13218	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/2296	1.30	6/3117 (0.2%)
1	F	0.73	0/2296	1.23	1/3117 (0.0%)
2	B	0.68	0/852	1.20	1/1152 (0.1%)
2	G	0.71	0/860	1.16	2/1162 (0.2%)
3	C	0.76	0/70	1.47	0/92
3	H	0.78	0/70	1.23	0/92
4	D	0.85	0/1568	1.32	7/2116 (0.3%)
4	I	0.78	0/1568	1.30	14/2116 (0.7%)
5	E	0.74	0/1986	1.21	0/2702
5	J	0.76	0/1986	1.20	2/2702 (0.1%)
All	All	0.75	0/13552	1.25	33/18368 (0.2%)

There are no bond length outliers.

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	I	210	PHE	CA-CB-CG	6.84	120.64	113.80
4	D	76	PHE	CA-CB-CG	6.69	120.49	113.80
4	I	105	ASP	CA-CB-CG	6.44	119.04	112.60
4	D	105	ASP	CA-CB-CG	6.32	118.92	112.60
4	D	12	ILE	N-CA-CB	6.18	119.44	111.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2235	0	2085	15	0
1	F	2235	0	2089	24	0
2	B	829	0	794	10	0
2	G	837	0	805	5	0
3	C	70	0	73	1	0
3	H	70	0	73	5	0
4	D	1535	0	1477	35	0
4	I	1535	0	1476	37	0
5	E	1935	0	1878	28	0
5	J	1935	0	1879	23	0
6	E	1	0	0	0	0
6	F	1	0	0	0	0
All	All	13218	0	12629	154	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 154 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:I:60:GLY:HA2	4:I:61:GLU:HB2	1.13	1.12
4:D:60:GLY:HA2	4:D:61:GLU:HB2	1.09	1.07
4:D:60:GLY:CA	4:D:61:GLU:HB2	1.99	0.93
1:F:155:GLN:NE2	5:J:114:GLY:O	2.02	0.93
4:I:60:GLY:HA2	4:I:61:GLU:CB	2.01	0.89

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	272/274 (99%)	260 (96%)	11 (4%)	1 (0%)	30 62

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	272/274 (99%)	254 (93%)	17 (6%)	1 (0%)	30	62
2	B	97/100 (97%)	95 (98%)	2 (2%)	0	100	100
2	G	98/100 (98%)	91 (93%)	6 (6%)	1 (1%)	12	45
3	C	8/10 (80%)	7 (88%)	1 (12%)	0	100	100
3	H	8/10 (80%)	7 (88%)	1 (12%)	0	100	100
4	D	196/198 (99%)	173 (88%)	18 (9%)	5 (3%)	4	26
4	I	196/198 (99%)	174 (89%)	17 (9%)	5 (3%)	4	26
5	E	242/244 (99%)	227 (94%)	13 (5%)	2 (1%)	16	50
5	J	242/244 (99%)	219 (90%)	20 (8%)	3 (1%)	10	42
All	All	1631/1652 (99%)	1507 (92%)	106 (6%)	18 (1%)	11	43

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	61	GLU
4	D	135	ARG
5	J	15	LYS
4	D	139	SER
5	E	15	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	216 (94%)	15 (6%)	15	47
1	F	231/231 (100%)	217 (94%)	14 (6%)	17	49
2	B	81/95 (85%)	73 (90%)	8 (10%)	7	31
2	G	95/95 (100%)	88 (93%)	7 (7%)	13	43
3	C	8/8 (100%)	7 (88%)	1 (12%)	4	21
3	H	8/8 (100%)	8 (100%)	0	100	100
4	D	173/173 (100%)	158 (91%)	15 (9%)	9	36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	I	173/173 (100%)	149 (86%)	24 (14%)	3	17
5	E	209/209 (100%)	195 (93%)	14 (7%)	15	47
5	J	209/209 (100%)	180 (86%)	29 (14%)	3	17
All	All	1418/1432 (99%)	1291 (91%)	127 (9%)	9	34

5 of 127 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	61	ASP
5	J	96	LEU
2	G	17	ASN
5	J	81	LEU
5	J	162	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 50 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	62	GLN
4	I	46	HIS
5	J	249	GLN
1	F	151	HIS
2	G	83	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	274/274 (100%)	-0.05	2 (0%) 84 69	13, 33, 71, 94	0
1	F	274/274 (100%)	-0.01	4 (1%) 72 52	9, 32, 73, 87	0
2	B	99/100 (99%)	-0.20	0 100 100	16, 27, 52, 70	0
2	G	100/100 (100%)	0.18	3 (3%) 52 33	19, 42, 70, 85	0
3	C	10/10 (100%)	-0.25	0 100 100	11, 21, 29, 54	0
3	H	10/10 (100%)	-0.51	0 100 100	17, 22, 27, 34	0
4	D	198/198 (100%)	0.36	9 (4%) 38 24	21, 47, 81, 89	1 (0%)
4	I	198/198 (100%)	0.43	8 (4%) 42 26	15, 46, 83, 95	1 (0%)
5	E	244/244 (100%)	-0.16	1 (0%) 88 79	11, 31, 58, 92	6 (2%)
5	J	244/244 (100%)	-0.08	2 (0%) 82 67	13, 33, 60, 91	6 (2%)
All	All	1651/1652 (99%)	0.04	29 (1%) 67 48	9, 35, 75, 95	14 (0%)

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	I	176	SER	4.9
4	I	175	ARG	4.3
4	I	173	ASP	4.1
2	G	0	MET	3.7
4	D	207	ASP	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.