



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2026 – 10:36 AM UTC

PDB ID : 5WKY / pdb_00005wky
Title : Bromide sites in the structure of an acid sensing ion channel in a resting state
Authors : Yoder, N.; Gouaux, E.
Deposited on : 2017-07-25
Resolution : 4.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1082 (4.20-3.80)
Clashscore	190562	1129 (4.20-3.80)
Ramachandran outliers	187476	1064 (4.20-3.80)
Sidechain outliers	187428	1055 (4.20-3.80)
RSRZ outliers	180081	1082 (4.20-3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	439	5% 79% 15% • 5%
1	B	439	6% 83% 13% 5%
1	C	439	10% 83% 12% 6%
2	D	2	100%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9355 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acid-sensing ion channel 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	417	Total	C	N	O	S	0	0	0
			3140	2019	515	579	27			
1	C	414	Total	C	N	O	S	0	0	0
			3012	1933	494	559	26			
1	B	418	Total	C	N	O	S	0	0	0
			3115	2000	508	580	27			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	C	1	Total	Cl	0	0
			1	1		
4	B	1	Total	Cl	0	0
			1	1		


- Molecule 5 is BARIUM ION (CCD ID: BA) (formula: Ba).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ba	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acid-sensing ion channel 1




- Molecule 1: Acid-sensing ion channel 1



- Molecule 1: Acid-sensing ion channel 1





● Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	109.55Å 134.49Å 159.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.15 – 4.00 43.15 – 4.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.15-4.00) 99.8 (43.15-4.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 4.00Å)	Xtriage
Refinement program	PHENIX (dev_2597: ???)	Depositor
R, R_{free}	0.281 , 0.299 0.282 , 0.300	Depositor DCC
R_{free} test set	1017 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	162.5	Xtriage
Anisotropy	0.640	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.22 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.82	EDS
Total number of atoms	9355	wwPDB-VP
Average B, all atoms (Å ²)	132.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NAG, BA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/3215	0.42	0/4376
1	B	0.20	0/3189	0.42	0/4349
1	C	0.17	0/3081	0.39	0/4206
All	All	0.20	0/9485	0.41	0/12931

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3140	0	2895	48	0
1	B	3115	0	2836	39	0
1	C	3012	0	2680	32	0
2	D	28	0	25	0	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
3	C	28	0	26	0	0
4	A	1	0	0	1	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	1	0	0	0	0
All	All	9355	0	8488	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:ARG:HH22	1:A:349:LEU:HD13	1.46	0.81
1:A:192:TYR:HE2	1:A:247:LYS:HD3	1.53	0.73
1:A:291:CYS:SG	1:A:292:LYS:N	2.64	0.70
1:C:96:LEU:HD13	1:C:243:GLU:HG3	1.72	0.70
1:A:133:GLU:O	1:A:137:GLU:HG2	1.95	0.66
1:B:192:TYR:HE2	1:B:247:LYS:HD3	1.63	0.63
1:C:192:TYR:HE2	1:C:247:LYS:HD3	1.62	0.63
1:A:365:PRO:HG2	1:A:368:VAL:HG22	1.80	0.62
1:C:275:SER:O	1:C:373:LYS:HA	1.99	0.62
1:A:96:LEU:HD13	1:A:243:GLU:HG3	1.80	0.62
1:B:276:CYS:HA	1:B:372:GLY:O	1.99	0.62
1:B:192:TYR:CE2	1:B:247:LYS:HD3	2.34	0.61
1:C:325:ARG:HD3	1:C:333:ALA:HB3	1.83	0.61
1:A:99:PHE:CE2	1:A:231:LEU:HD21	2.36	0.60
1:B:130:THR:O	1:B:130:THR:HG22	2.00	0.60
1:B:325:ARG:HH22	1:B:331:GLY:C	2.09	0.60
1:B:326:MET:HE3	1:B:345:ALA:HB1	1.85	0.59
1:A:76:THR:HG21	1:C:78:LEU:HD12	1.84	0.59
1:A:78:LEU:HD12	1:B:76:THR:HG21	1.85	0.59
1:A:326:MET:HE3	1:A:345:ALA:HB1	1.85	0.59
1:A:276:CYS:HA	1:A:372:GLY:O	2.03	0.58
1:C:276:CYS:HA	1:C:372:GLY:O	2.04	0.57
1:B:275:SER:O	1:B:373:LYS:HA	2.05	0.56
1:C:76:THR:HG21	1:B:78:LEU:HD12	1.88	0.55
1:A:275:SER:O	1:A:373:LYS:HA	2.07	0.55
1:A:422:LYS:NZ	4:A:502:CL:CL	2.77	0.55
1:A:378:VAL:HG11	1:B:96:LEU:HD11	1.87	0.54
1:C:365:PRO:HG2	1:C:368:VAL:HG22	1.90	0.54
1:C:192:TYR:CE2	1:C:247:LYS:HD3	2.43	0.53
1:C:156:LEU:HD11	1:C:327:VAL:HA	1.91	0.53
1:C:99:PHE:CZ	1:C:231:LEU:HD21	2.44	0.52
1:A:96:LEU:HD11	1:C:378:VAL:HG11	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:PHE:CE1	1:A:116:LEU:HD22	2.44	0.52
1:A:227:GLN:NE2	1:A:402:GLU:O	2.43	0.51
1:B:451:GLU:O	1:B:454:ASP:HB3	2.11	0.51
1:A:304:TYR:OH	1:A:363:GLU:O	2.17	0.51
1:A:113:GLY:HA3	1:A:119:LEU:HD12	1.93	0.51
1:A:268:PRO:HA	1:A:405:LEU:HB3	1.93	0.50
1:C:158:PHE:O	1:C:162:ALA:HB3	2.11	0.50
1:C:190:THR:OG1	1:C:195:CYS:SG	2.70	0.50
1:B:156:LEU:HD12	1:B:327:VAL:HG12	1.93	0.50
1:B:99:PHE:CZ	1:B:231:LEU:HD21	2.46	0.50
1:C:227:GLN:NE2	1:C:402:GLU:O	2.45	0.49
1:A:88:PHE:CG	1:A:89:PRO:HD2	2.48	0.49
1:A:385:SER:HB2	1:B:242:PHE:HD1	1.77	0.49
1:A:268:PRO:HD2	1:C:379:LYS:HB2	1.95	0.49
1:B:281:LEU:HD23	1:B:283:TYR:OH	2.13	0.49
1:A:118:LEU:HD21	1:A:231:LEU:HD22	1.95	0.49
1:A:192:TYR:CE2	1:A:247:LYS:HD3	2.40	0.48
1:A:210:THR:HG21	1:A:411:PHE:HD2	1.77	0.48
1:B:88:PHE:CG	1:B:89:PRO:HD2	2.48	0.48
1:B:318:LEU:HD12	1:B:329:MET:HE1	1.95	0.48
1:A:379:LYS:HB2	1:B:268:PRO:HD2	1.94	0.48
1:C:414:LEU:HD23	1:C:414:LEU:HA	1.74	0.48
1:A:109:LEU:HD23	1:A:143:ALA:HB2	1.95	0.47
1:A:130:THR:HG22	1:C:387:LYS:HB3	1.97	0.47
1:C:88:PHE:CG	1:C:89:PRO:HD2	2.50	0.47
1:B:220:GLU:HA	1:B:407:LEU:O	2.15	0.46
1:A:377:MET:SD	1:B:377:MET:HE1	2.55	0.46
1:C:211:MET:C	1:C:414:LEU:HD11	2.41	0.46
1:A:365:PRO:HG2	1:A:368:VAL:CG2	2.45	0.46
1:C:198:PHE:CD2	1:C:219:LEU:HD22	2.51	0.46
1:C:268:PRO:HA	1:C:405:LEU:HB3	1.99	0.45
1:A:435:GLY:HA3	1:B:436:GLY:O	2.16	0.45
1:A:388:TYR:N	1:B:130:THR:HG21	2.31	0.45
1:C:62:CYS:HA	1:C:437:GLN:NE2	2.32	0.45
1:A:445:SER:H	1:A:448:THR:HB	1.81	0.44
1:C:280:ARG:HD2	1:C:416:TYR:CD2	2.52	0.44
1:A:112:ALA:O	1:A:116:LEU:HG	2.17	0.44
1:B:398:GLN:O	1:B:402:GLU:HG2	2.17	0.44
1:B:61:VAL:O	1:B:64:ASN:HB3	2.17	0.44
1:B:268:PRO:HA	1:B:405:LEU:HB3	1.98	0.44
1:B:325:ARG:NH2	1:B:333:ALA:H	2.15	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:ARG:HD2	1:A:416:TYR:CD1	2.53	0.44
1:C:94:CYS:SG	1:C:259:ILE:HG21	2.57	0.43
1:B:93:PHE:HZ	1:B:166:ILE:HD13	1.82	0.43
1:B:96:LEU:HD13	1:B:243:GLU:HG3	2.00	0.43
1:B:168:GLU:HB3	1:B:226:GLN:NE2	2.34	0.43
1:B:365:PRO:HG2	1:B:368:VAL:HG22	1.99	0.43
1:B:325:ARG:CG	1:B:329:MET:HB2	2.48	0.43
1:A:72:TYR:CD1	1:A:287:PRO:HG2	2.53	0.43
1:C:94:CYS:SG	1:C:259:ILE:HD13	2.59	0.43
1:B:274:VAL:HG22	1:B:375:LEU:CD2	2.49	0.42
1:B:96:LEU:HD23	1:B:96:LEU:HA	1.89	0.42
1:A:62:CYS:HA	1:A:437:GLN:NE2	2.35	0.42
1:C:381:PRO:HB3	1:C:389:LEU:HD12	2.00	0.42
1:C:281:LEU:HD21	1:C:370:ARG:HH21	1.85	0.42
1:A:92:THR:CG2	1:A:259:ILE:HD11	2.50	0.41
1:C:434:ILE:HA	1:C:437:GLN:NE2	2.34	0.41
1:A:227:GLN:HE21	1:A:227:GLN:HB2	1.62	0.41
1:A:54:SER:OG	1:A:441:PHE:O	2.34	0.41
1:A:99:PHE:CZ	1:A:231:LEU:HD11	2.55	0.41
1:A:215:THR:HA	1:A:410:PHE:CD1	2.56	0.41
1:C:280:ARG:HD2	1:C:416:TYR:CE2	2.55	0.41
1:B:129:GLN:H	1:B:129:GLN:HG3	1.71	0.41
1:A:435:GLY:HA3	1:B:440:LEU:HB2	2.02	0.41
1:C:242:PHE:CE2	1:B:389:LEU:HD21	2.56	0.41
1:A:301:TYR:CE2	1:A:311:ILE:HD11	2.56	0.41
1:A:391:LYS:HE2	1:B:129:GLN:HB3	2.03	0.40
1:A:106:LYS:HG3	1:A:142:LYS:O	2.21	0.40
1:B:381:PRO:HB3	1:B:389:LEU:HD12	2.03	0.40
1:A:454:ASP:O	1:A:457:TYR:HB3	2.21	0.40
1:C:65:ARG:NH1	1:B:426:GLU:OE1	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/439 (94%)	409 (99%)	6 (1%)	0	100	100
1	B	416/439 (95%)	408 (98%)	8 (2%)	0	100	100
1	C	410/439 (93%)	403 (98%)	7 (2%)	0	100	100
All	All	1241/1317 (94%)	1220 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/384 (80%)	303 (98%)	5 (2%)	55	70
1	B	303/384 (79%)	300 (99%)	3 (1%)	68	76
1	C	278/384 (72%)	275 (99%)	3 (1%)	65	74
All	All	889/1152 (77%)	878 (99%)	11 (1%)	63	73

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	THR
1	A	180	CYS
1	A	291	CYS
1	A	292	LYS
1	A	366	CYS
1	C	180	CYS
1	C	291	CYS
1	C	366	CYS
1	B	180	CYS
1	B	291	CYS
1	B	366	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	135	GLN
1	A	277	GLN
1	A	328	HIS
1	A	415	ASN
1	C	74	HIS
1	C	227	GLN
1	B	74	HIS
1	B	227	GLN
1	B	279	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	D	1	2,1	14,14,15	0.25	0	17,19,21	0.44	0
2	NAG	D	2	2	14,14,15	0.28	0	17,19,21	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	2	NAG	C4-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	503	1	14,14,15	0.33	0	17,19,21	0.47	0
3	NAG	A	501	1	14,14,15	0.25	0	17,19,21	0.41	0
3	NAG	C	502	1	14,14,15	0.31	0	17,19,21	0.47	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	501	1	14,14,15	0.24	0	17,19,21	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	503	1	-	2/6/23/26	0/1/1/1
3	NAG	A	501	1	-	0/6/23/26	0/1/1/1
3	NAG	C	502	1	-	1/6/23/26	0/1/1/1
3	NAG	C	501	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	503	NAG	C4-C5-C6-O6
3	B	503	NAG	O5-C5-C6-O6
3	C	501	NAG	C1-C2-N2-C7
3	C	501	NAG	C3-C2-N2-C7
3	C	502	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	417/439 (94%)	0.34	23 (5%)	30	26	61, 116, 191, 227	0
1	B	418/439 (95%)	0.44	27 (6%)	25	23	66, 124, 194, 237	0
1	C	414/439 (94%)	0.64	43 (10%)	11	14	78, 144, 238, 260	0
All	All	1249/1317 (94%)	0.47	93 (7%)	20	20	61, 127, 213, 260	0

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	71	LEU	6.7
1	A	374	GLU	6.6
1	C	408	ASP	6.6
1	B	275	SER	5.5
1	B	372	GLY	5.2
1	C	273	PHE	5.2
1	B	70	PHE	5.1
1	B	374	GLU	5.0
1	C	272	THR	5.0
1	B	72	TYR	5.0
1	B	373	LYS	4.9
1	A	275	SER	4.8
1	A	277	GLN	4.7
1	C	219	LEU	4.6
1	C	220	GLU	4.6
1	A	272	THR	4.6
1	C	251	HIS	4.6
1	A	130	THR	4.4
1	C	130	THR	4.3
1	C	274	VAL	4.3
1	A	147	ASN	4.3
1	C	271	GLN	4.1
1	B	69	TYR	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	72	TYR	3.9
1	C	276	CYS	3.8
1	A	412	GLU	3.8
1	B	98	GLU	3.8
1	C	275	SER	3.8
1	B	67	GLN	3.7
1	C	90	ALA	3.6
1	C	410	PHE	3.6
1	B	276	CYS	3.6
1	B	427	VAL	3.6
1	C	177	GLY	3.5
1	C	137	GLU	3.4
1	A	262	LEU	3.3
1	A	259	ILE	3.3
1	A	263	GLY	3.1
1	C	409	ILE	3.1
1	B	430	LEU	3.1
1	C	200	ALA	3.1
1	C	352	LEU	3.1
1	C	239	GLU	3.1
1	C	372	GLY	3.0
1	C	374	GLU	3.0
1	A	274	VAL	3.0
1	C	98	GLU	2.9
1	A	271	GLN	2.9
1	B	287	PRO	2.9
1	B	118	LEU	2.9
1	B	232	PRO	2.8
1	A	323	ASN	2.8
1	B	179	GLN	2.8
1	A	276	CYS	2.8
1	C	348	ALA	2.8
1	A	319	VAL	2.8
1	B	101	PHE	2.8
1	A	408	ASP	2.7
1	C	375	LEU	2.7
1	A	416	TYR	2.7
1	C	89	PRO	2.7
1	B	262	LEU	2.7
1	A	98	GLU	2.6
1	A	249	GLN	2.6
1	C	118	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	150	PRO	2.6
1	C	431	LEU	2.5
1	A	417	GLU	2.5
1	C	250	ILE	2.5
1	C	373	LYS	2.5
1	C	198	PHE	2.5
1	B	263	GLY	2.5
1	C	252	SER	2.4
1	B	132	ASP	2.4
1	C	332	ASP	2.4
1	B	68	TYR	2.4
1	C	129	GLN	2.4
1	C	174	PHE	2.4
1	C	249	GLN	2.3
1	A	373	LYS	2.3
1	C	270	PHE	2.2
1	C	43	LYS	2.2
1	C	295	THR	2.2
1	B	63	THR	2.2
1	A	183	GLU	2.2
1	B	174	PHE	2.1
1	A	375	LEU	2.1
1	C	231	LEU	2.1
1	B	408	ASP	2.1
1	C	378	VAL	2.1
1	B	79	ASP	2.0
1	C	351	PHE	2.0
1	C	407	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	D	2	14/15	0.05	0.13	151,189,219,220	0
2	NAG	D	1	14/15	0.48	0.12	130,159,200,216	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	501	14/15	0.32	0.08	153,178,191,195	0
4	CL	C	503	1/1	0.37	0.18	155,155,155,155	0
3	NAG	B	503	14/15	0.51	0.11	149,181,192,207	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	C	502	14/15	0.54	0.10	177,199,217,218	0
3	NAG	C	501	14/15	0.62	0.20	152,175,191,192	0
4	CL	B	504	1/1	0.65	0.12	125,125,125,125	0
4	CL	A	502	1/1	0.66	0.17	111,111,111,111	0
5	BA	A	503	1/1	0.68	0.20	254,254,254,254	0

6.5 Other polymers [i](#)

There are no such residues in this entry.