



## wwPDB EM Validation Summary Report ⓘ

Mar 29, 2026 – 09:58 AM UTC

PDB ID : 6WLL / pdb\_00006wll  
EMDB ID : EMD-21833  
Title : Apo F. nucleatum glycine riboswitch models, 10.0 Angstrom resolution  
Authors : Kappel, K.; Zhang, K.; Su, Z.; Watkins, A.M.; Kladwang, W.; Li, S.; Pintilie, G.; Topkar, V.V.; Rangan, R.; Zheludev, I.N.; Yesselman, J.D.; Chiu, W.; Das, R.  
Deposited on : 2020-04-20  
Resolution : 10.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132  
MolProbity : **FAILED**  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49

i

ELECTRON MICROSCOPY

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric  | EM structures<br>(#Entries) | Similar EM resolution<br>(#Entries, resolution range(Å)) |
|---------|-----------------------------|----------------------------------------------------------|
| Q-score | 25397                       | 147 ( 9.50 - 10.50 )                                     |

## 2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 110240 atoms, of which 36820 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (171-MER).

| Mol | Chain | Residues | Atoms         |           |           |          |           |          | AltConf | Trace |
|-----|-------|----------|---------------|-----------|-----------|----------|-----------|----------|---------|-------|
| 1   | 1-A   | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 2-A   | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 3-A   | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 4-A   | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 5-A   | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 6-A   | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 7-A   | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 8-A   | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 9-A   | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 10-A  | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 11-A  | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 12-A  | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 13-A  | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 14-A  | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 15-A  | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 16-A  | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |
| 1   | 17-A  | 171      | Total<br>5512 | C<br>1646 | H<br>1841 | N<br>689 | O<br>1166 | P<br>170 | 0       | 0     |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Residues | Atoms |      |      |     |      |     | AltConf | Trace |
|-----|-------|----------|-------|------|------|-----|------|-----|---------|-------|
| 1   | 18-A  | 171      | Total | C    | H    | N   | O    | P   | 0       | 0     |
|     |       |          | 5512  | 1646 | 1841 | 689 | 1166 | 170 |         |       |
| 1   | 19-A  | 171      | Total | C    | H    | N   | O    | P   | 0       | 0     |
|     |       |          | 5512  | 1646 | 1841 | 689 | 1166 | 170 |         |       |
| 1   | 20-A  | 171      | Total | C    | H    | N   | O    | P   | 0       | 0     |
|     |       |          | 5512  | 1646 | 1841 | 689 | 1166 | 170 |         |       |

There are 2 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment        | Reference    |
|-------|---------|----------|--------|----------------|--------------|
| A     | -12     | G        | -      | expression tag | GB 929732263 |
| A     | -11     | G        | -      | expression tag | GB 929732263 |

MolProbity failed to run properly - this section is therefore empty.

### 3 Experimental information

| Property                             | Value                           | Source    |
|--------------------------------------|---------------------------------|-----------|
| EM reconstruction method             | SINGLE PARTICLE                 | Depositor |
| Imposed symmetry                     | POINT, Not provided             |           |
| Number of particles used             | 20269                           | Depositor |
| Resolution determination method      | FSC 0.143 CUT-OFF               | Depositor |
| CTF correction method                | PHASE FLIPPING ONLY             | Depositor |
| Microscope                           | FEI TALOS ARCTICA               | Depositor |
| Voltage (kV)                         | 200                             | Depositor |
| Electron dose ( $e^-/\text{\AA}^2$ ) | 30                              | Depositor |
| Minimum defocus (nm)                 | Not provided                    |           |
| Maximum defocus (nm)                 | Not provided                    |           |
| Magnification                        | Not provided                    |           |
| Image detector                       | GATAN K2 SUMMIT (4k x 4k)       | Depositor |
| Maximum map value                    | 0.128                           | Depositor |
| Minimum map value                    | -0.032                          | Depositor |
| Average map value                    | 0.001                           | Depositor |
| Map value standard deviation         | 0.008                           | Depositor |
| Recommended contour level            | 0.05                            | Depositor |
| Map size ( $\text{\AA}$ )            | 171.20001, 171.20001, 171.20001 | wwPDB     |
| Map dimensions                       | 160, 160, 160                   | wwPDB     |
| Map angles ( $^\circ$ )              | 90.0, 90.0, 90.0                | wwPDB     |
| Pixel spacing ( $\text{\AA}$ )       | 1.07, 1.07, 1.07                | Depositor |

## 4 Model quality [i](#)

### 4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles [i](#)

#### 4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 4.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 4.7 Other polymers [i](#)

There are no such residues in this entry.

## 4.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 5 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-21833. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 5.1 Orthogonal projections [i](#)

#### 5.1.1 Primary map



X



Y



Z

The images above show the map projected in three orthogonal directions.

### 5.2 Central slices [i](#)

#### 5.2.1 Primary map



X Index: 80



Y Index: 80



Z Index: 80

The images above show central slices of the map in three orthogonal directions.

## 5.3 Largest variance slices [i](#)

### 5.3.1 Primary map



X Index: 75



Y Index: 83



Z Index: 66


The images above show the largest variance slices of the map in three orthogonal directions.

## 5.4 Orthogonal standard-deviation projections (False-color) [i](#)

### 5.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 5.5 Orthogonal surface views [i](#)

### 5.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.05. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 5.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 6 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 6.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 6.2 Volume estimate [i](#)



The volume at the recommended contour level is 49 nm<sup>3</sup>; this corresponds to an approximate mass of 45 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 6.3 Rotationally averaged power spectrum ⓘ



\*Reported resolution corresponds to spatial frequency of 0.100 Å<sup>-1</sup>


## 7 Fourier-Shell correlation ⓘ

This section was not generated. No FSC curve or half-maps provided.


## 8 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-21833 and PDB model 6WLL. Per-residue inclusion information can be found in section ?? on page ??.


### 8.1 Map-model overlay [i](#)



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.05 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 8.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 8.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.05).

## 8.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.

8.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.05) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion                | Q-score                       |
|-------|-------------------------------|-------------------------------|
| All   | <div><div></div></div> 0.8050 | <div><div></div></div> 0.1510 |
| A     | <div><div></div></div> 0.8110 | <div><div></div></div> 0.1510 |

