



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:26 PM UTC

PDB ID : 2WM4 / pdb_00002wm4
Title : X-ray structure of Mycobacterium tuberculosis cytochrome P450 CYP124 in complex with phytanic acid
Authors : Johnston, J.B.; Kells, P.M.; Podust, L.M.; Ortiz de Montellano, P.R.
Deposited on : 2009-06-30
Resolution : 2.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	435	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PUTATIVE CYTOCHROME P450 124.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	426	3360	2124	597	628	11	0	2	0

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	expression tag	UNP P0A516
A	-5	HIS	-	expression tag	UNP P0A516
A	-4	HIS	-	expression tag	UNP P0A516
A	-3	HIS	-	expression tag	UNP P0A516
A	-2	HIS	-	expression tag	UNP P0A516
A	-1	HIS	-	expression tag	UNP P0A516
A	0	HIS	-	expression tag	UNP P0A516

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is (3R,7S,11S)-3,7,11,15-tetramethylhexadecanoic acid (CCD ID: VGJ) (formula: $C_{20}H_{40}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			22	20	2		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	174	Total	O	0	0
			174	174		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PUTATIVE CYTOCHROME P450 124



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.60Å 73.40Å 109.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.94 – 2.11 43.94 – 2.11	Depositor EDS
% Data completeness (in resolution range)	96.8 (43.94-2.11) 97.0 (43.94-2.11)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.166 , 0.232 0.167 , 0.233	Depositor DCC
R_{free} test set	1197 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.613	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3600	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CA, VGJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	6/3446 (0.2%)	1.16	11/4700 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	176	ILE	CA-CB	6.57	1.62	1.54
1	A	346	ALA	CA-CB	6.56	1.63	1.53
1	A	326	ILE	CA-CB	6.51	1.62	1.55
1	A	58	ILE	CA-CB	6.03	1.60	1.54
1	A	94	ILE	CA-CB	5.97	1.61	1.54
1	A	301	ALA	CA-CB	-5.70	1.46	1.53

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	142	ARG	CA-C-N	-7.62	107.89	122.06
1	A	142	ARG	C-N-CA	-7.62	107.89	122.06
1	A	411	ARG	NE-CZ-NH1	-6.32	115.19	121.50
1	A	96	ILE	N-CA-C	6.22	117.00	110.72
1	A	60	LEU	N-CA-C	5.75	122.53	109.81
1	A	411	ARG	NE-CZ-NH2	5.50	124.15	119.20
1	A	271	THR	N-CA-CB	-5.50	101.20	110.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	36	ASP	N-CA-C	5.37	117.83	111.33
1	A	237	ASP	N-CA-C	-5.13	101.50	109.14
1	A	272	THR	N-CA-C	5.11	117.59	111.71
1	A	397	ARG	N-CA-C	-5.01	105.82	111.28

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	247	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3360	0	3240	76	0
2	A	43	0	30	6	0
3	A	22	0	39	4	0
4	A	1	0	0	0	0
5	A	174	0	0	10	0
All	All	3600	0	3309	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:PHE:HD2	1:A:213:MET:HE2	1.35	0.92
1:A:110:MET:HE3	1:A:260:PHE:CZ	2.08	0.89
1:A:212:PHE:CD2	1:A:213:MET:HE2	2.10	0.87
1:A:110:MET:CE	1:A:260:PHE:CZ	2.62	0.82
1:A:57:THR:HG22	1:A:68:GLY:HA2	1.62	0.81
1:A:235:HIS:O	1:A:236:ASP:CB	2.30	0.76
1:A:110:MET:HE2	1:A:121:LEU:HB3	1.67	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:LEU:HD22	1:A:266:VAL:CG1	2.18	0.74
1:A:271:THR:CG2	1:A:272:THR:H	2.02	0.73
1:A:271:THR:HG21	2:A:450:HEM:CHC	2.19	0.72
1:A:231:ARG:NH2	5:A:2100:HOH:O	2.21	0.72
1:A:271:THR:HG22	1:A:272:THR:H	1.57	0.69
1:A:97:ASN:ND2	3:A:1429:VGJ:H141	2.07	0.69
1:A:271:THR:CG2	1:A:272:THR:N	2.56	0.68
1:A:395:GLU:OE1	5:A:2161:HOH:O	2.11	0.68
1:A:146:HIS:NE2	1:A:398:ARG:NH2	2.42	0.67
1:A:234:HIS:HD2	1:A:244:GLU:OE1	1.78	0.67
1:A:105:GLU:OE1	5:A:2051:HOH:O	2.15	0.64
1:A:79:PHE:O	1:A:83:ARG:HG2	1.98	0.63
1:A:212:PHE:CD2	1:A:213:MET:CE	2.82	0.63
1:A:146:HIS:CE1	1:A:398:ARG:HH22	2.18	0.61
1:A:110:MET:HE1	1:A:260:PHE:CZ	2.34	0.61
1:A:144:ARG:HG2	1:A:174:GLN:OE1	2.00	0.61
1:A:69:HIS:HE1	5:A:2029:HOH:O	1.82	0.61
1:A:110:MET:HE2	1:A:121:LEU:CB	2.29	0.61
1:A:212:PHE:HD2	1:A:213:MET:CE	2.12	0.60
1:A:235:HIS:O	1:A:236:ASP:HB2	1.99	0.60
1:A:60:LEU:HD11	3:A:1429:VGJ:H162	1.84	0.60
1:A:57:THR:CG2	1:A:67:ASN:O	2.49	0.59
1:A:272:THR:HG21	1:A:388:GLU:OE1	2.02	0.59
1:A:274:ASN:ND2	1:A:420:ILE:H	2.00	0.59
1:A:271:THR:HG23	1:A:272:THR:N	2.18	0.59
1:A:387:ARG:NH2	1:A:390:ARG:HD3	2.20	0.57
1:A:271:THR:CG2	2:A:450:HEM:CHC	2.82	0.57
1:A:285:ARG:NH2	1:A:408:GLU:OE1	2.38	0.56
1:A:35:ASP:HA	1:A:411:ARG:NH1	2.21	0.56
1:A:253:SER:OG	1:A:256:GLU:HG3	2.06	0.55
1:A:110:MET:HE1	1:A:122:ARG:HA	1.88	0.55
1:A:146:HIS:HD2	5:A:2160:HOH:O	1.90	0.55
1:A:170:PRO:O	1:A:174:GLN:HG3	2.07	0.54
1:A:97:ASN:HD22	3:A:1429:VGJ:H141	1.71	0.54
1:A:95:THR:O	1:A:319:ARG:HD3	2.07	0.53
1:A:138:GLU:O	1:A:142:ARG:HB2	2.10	0.52
1:A:95:THR:OG1	1:A:97:ASN:ND2	2.44	0.51
1:A:198:LEU:HD22	1:A:266:VAL:HG13	1.93	0.50
1:A:130:THR:HG22	1:A:133:VAL:H	1.76	0.50
1:A:86[A]:ASP:OD2	5:A:2040:HOH:O	2.20	0.49
1:A:57:THR:HG21	1:A:67:ASN:O	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:THR:HB	1:A:96:ILE:CG2	2.43	0.48
1:A:69:HIS:HD2	1:A:340:SER:OG	1.95	0.48
1:A:235:HIS:O	1:A:236:ASP:HB3	2.14	0.47
1:A:69:HIS:CD2	1:A:340:SER:OG	2.68	0.47
1:A:130:THR:O	1:A:134:VAL:HG13	2.14	0.46
1:A:110:MET:HE1	1:A:260:PHE:HZ	1.79	0.46
2:A:450:HEM:HBC2	2:A:450:HEM:HHD	1.98	0.45
1:A:387:ARG:HH22	1:A:390:ARG:HD3	1.79	0.45
1:A:130:THR:CG2	1:A:133:VAL:H	2.30	0.45
1:A:288:GLU:HB3	5:A:2124:HOH:O	2.17	0.44
1:A:102:GLU:H	1:A:102:GLU:CD	2.26	0.44
1:A:75:TYR:CD2	1:A:75:TYR:C	2.96	0.44
1:A:57:THR:HB	1:A:96:ILE:HG21	1.99	0.44
1:A:399:GLN:HG3	5:A:2161:HOH:O	2.17	0.44
1:A:110:MET:HE3	1:A:260:PHE:CE2	2.52	0.44
1:A:343:TYR:CE1	1:A:372:GLY:HA2	2.53	0.43
1:A:378:PHE:O	1:A:379:CYS:C	2.61	0.43
1:A:98:ASP:HA	5:A:2048:HOH:O	2.19	0.43
1:A:96:ILE:HA	1:A:319:ARG:HD2	2.01	0.42
1:A:62:GLY:HA3	1:A:209:PHE:CG	2.54	0.42
3:A:1429:VGJ:H161	3:A:1429:VGJ:H111	1.90	0.42
1:A:51:PRO:HB2	1:A:329:ARG:HD3	2.01	0.41
1:A:212:PHE:CE2	1:A:213:MET:CE	3.04	0.41
1:A:7:ILE:HG21	1:A:7:ILE:HD13	1.62	0.41
1:A:407:GLU:HB2	1:A:424:PRO:HG2	2.02	0.41
2:A:450:HEM:HMB2	2:A:450:HEM:HBB2	2.02	0.41
1:A:359:PHE:CZ	1:A:361:LEU:HD21	2.55	0.41
1:A:144:ARG:HA	5:A:2080:HOH:O	2.21	0.40
2:A:450:HEM:HHD	2:A:450:HEM:CBC	2.51	0.40
2:A:450:HEM:HBB2	2:A:450:HEM:CMB	2.51	0.40
1:A:307:GLU:HA	1:A:307:GLU:OE1	2.22	0.40
1:A:32:TRP:CZ2	1:A:415:GLN:HG3	2.56	0.40
1:A:118:HIS:CD2	1:A:377:HIS:CE1	3.10	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	426/435 (98%)	410 (96%)	12 (3%)	4 (1%)	14 10

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	60	LEU
1	A	248	ASP
1	A	236	ASP
1	A	59	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	353/364 (97%)	329 (93%)	24 (7%)	14 11

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ILE
1	A	9	THR
1	A	14	THR
1	A	57	THR
1	A	94	ILE
1	A	103	LEU
1	A	130	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	134	VAL
1	A	142	ARG
1	A	144	ARG
1	A	198	LEU
1	A	229	ASP
1	A	231	ARG
1	A	236	ASP
1	A	254	SER
1	A	265	VAL
1	A	271	THR
1	A	272	THR
1	A	292	ARG
1	A	294	TRP
1	A	300	LEU
1	A	351	SER
1	A	387	ARG
1	A	398	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	HIS
1	A	97	ASN
1	A	99	GLN
1	A	119	GLN
1	A	234	HIS
1	A	274	ASN
1	A	278	HIS
1	A	364	ASN
1	A	383	ASN
1	A	399	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	VGJ	A	1429	-	21,21,21	0.80	0	24,25,25	1.07	2 (8%)
2	HEM	A	450	1	50,50,50	1.99	10 (20%)	67,82,82	1.68	11 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	VGJ	A	1429	-	-	10/22/22/22	-
2	HEM	A	450	1	-	0/14/54/54	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	450	HEM	C3D-C2D	7.04	1.52	1.36
2	A	450	HEM	FE-ND	5.16	2.10	1.94
2	A	450	HEM	FE-NA	4.75	2.10	1.95
2	A	450	HEM	FE-NC	4.68	2.10	1.95
2	A	450	HEM	CAB-C3B	3.07	1.55	1.47
2	A	450	HEM	CMC-C2C	2.37	1.55	1.50
2	A	450	HEM	FE-NB	2.29	2.01	1.94
2	A	450	HEM	CMD-C2D	2.23	1.55	1.50
2	A	450	HEM	C3C-C2C	-2.09	1.32	1.37
2	A	450	HEM	O1D-CGD	2.05	1.28	1.22

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	450	HEM	C3B-C2B-C1B	5.04	110.20	106.41
2	A	450	HEM	C4D-ND-C1D	4.58	110.62	105.21
2	A	450	HEM	CAD-C3D-C4D	4.35	132.27	124.70
2	A	450	HEM	CHD-C1D-ND	3.61	128.31	124.42
2	A	450	HEM	CHC-C4B-NB	3.01	127.66	124.42
2	A	450	HEM	CHB-C1B-NB	2.88	127.94	124.37
2	A	450	HEM	O2A-CGA-CBA	2.67	122.44	114.00
3	A	1429	VGJ	C18-C5-C4	-2.52	102.30	111.27
3	A	1429	VGJ	O1-C15-C14	2.48	121.72	114.00
2	A	450	HEM	C4C-C3C-C2C	2.29	108.80	106.81
2	A	450	HEM	C4D-C3D-C2D	-2.25	103.61	106.89
2	A	450	HEM	C2A-C1A-NA	-2.20	107.71	110.15
2	A	450	HEM	CAD-C3D-C2D	-2.05	124.02	127.87

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1429	VGJ	C11-C12-C13-C16
3	A	1429	VGJ	C5-C6-C7-C8
3	A	1429	VGJ	C9-C10-C11-C12
3	A	1429	VGJ	C11-C12-C13-C14
3	A	1429	VGJ	C13-C14-C15-O1
3	A	1429	VGJ	C3-C4-C5-C6
3	A	1429	VGJ	C18-C5-C6-C7
3	A	1429	VGJ	C6-C7-C8-C9
3	A	1429	VGJ	C13-C14-C15-O2
3	A	1429	VGJ	C20-C1-C2-C3


There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1429	VGJ	4	0
2	A	450	HEM	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	426/435 (97%)	-0.36	3 (0%) 84 86	13, 28, 41, 51	2 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	247	VAL	2.6
1	A	248	ASP	2.3
1	A	3	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	VGJ	A	1429	22/22	0.90	0.12	22,39,60,62	0
4	CA	A	1430	1/1	0.97	0.05	35,35,35,35	0
2	HEM	A	450	43/43	0.99	0.04	15,20,22,23	0


The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around VGJ A 1429:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.