



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:53 PM UTC

PDB ID : 3WML / pdb_00003wml
Title : Structure of phosphotriesterase mutant (S308L/Y309A) from Agrobacterium radiobacter
Authors : Jackson, C.J.; Carr, P.D.; Sugrue, E.
Deposited on : 2013-11-21
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2667 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphotriesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	0	0	0
			2506	1586	447	465	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	92	ALA	SER	conflict	UNP Q93LD7
A	265	ASP	ASN	conflict	UNP Q93LD7
A	308	LEU	SER	engineered mutation	UNP Q93LD7
A	309	ALA	TYR	engineered mutation	UNP Q93LD7

- Molecule 2 is FE (II) ION (CCD ID: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		

- Molecule 3 is COBALT (II) ION (CCD ID: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Co	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	155	Total	O	0	0
			155	155		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphotriesterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	109.00Å 109.00Å 61.77Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.36 – 1.99 29.36 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.36-1.99) 100.0 (29.36-1.99)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.157 , 0.187 0.173 , 0.196	Depositor DCC
R_{free} test set	1494 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 33.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.035 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2667	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, FE2, KCX, CO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.06	114/2544 (4.5%)	1.17	14/3455 (0.4%)

All (114) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	333	ILE	C-O	-9.47	1.16	1.24
1	A	342	PRO	C-O	-8.38	1.17	1.25
1	A	37	ILE	C-O	-8.08	1.16	1.24
1	A	298	VAL	C-O	-7.78	1.16	1.24
1	A	179	PHE	C-O	-7.55	1.15	1.24
1	A	49	ALA	C-O	-7.53	1.15	1.24
1	A	134	PRO	C-O	-7.35	1.16	1.24
1	A	136	LEU	C-O	-7.34	1.15	1.24
1	A	328	VAL	C-O	-7.26	1.18	1.24
1	A	41	ARG	C-O	-7.24	1.14	1.24
1	A	255	MET	C-O	-7.11	1.17	1.24
1	A	319	ARG	C-O	-7.03	1.16	1.24
1	A	72	PHE	C-O	-6.97	1.15	1.24
1	A	52	THR	C-O	-6.94	1.16	1.24
1	A	186	ALA	C-O	-6.85	1.16	1.24
1	A	92	ALA	C-O	-6.70	1.16	1.24
1	A	91	ARG	C-O	-6.69	1.16	1.24
1	A	194	THR	C-O	-6.69	1.15	1.23
1	A	113	LEU	C-O	-6.65	1.16	1.24
1	A	162	GLY	C-O	-6.63	1.15	1.24
1	A	191	SER	C-O	-6.60	1.16	1.24
1	A	80	ALA	C-O	-6.51	1.16	1.24
1	A	329	PRO	C-O	-6.51	1.16	1.24
1	A	353	ASN	C-O	-6.44	1.18	1.24
1	A	283	LEU	C-O	-6.39	1.16	1.24
1	A	45	PRO	C-O	-6.36	1.16	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	233	ASP	C-O	-6.36	1.15	1.24
1	A	204	ALA	C-O	-6.35	1.16	1.24
1	A	114	ALA	C-O	-6.33	1.16	1.24
1	A	294	ASP	C-O	-6.30	1.15	1.24
1	A	78	ALA	C-O	-6.26	1.16	1.24
1	A	108	ARG	C-O	-6.24	1.16	1.23
1	A	102	SER	C-O	-6.24	1.16	1.24
1	A	244	ALA	C-O	-6.24	1.16	1.24
1	A	149	PHE	C-O	-6.23	1.17	1.24
1	A	320	ILE	C-O	-6.16	1.17	1.24
1	A	115	GLU	C-O	-6.13	1.17	1.24
1	A	349	VAL	C-O	-6.10	1.17	1.24
1	A	88	ARG	C-O	-6.08	1.17	1.24
1	A	226	VAL	C-O	-6.05	1.17	1.24
1	A	84	VAL	C-O	-6.04	1.17	1.24
1	A	358	LEU	C-O	-6.03	1.17	1.24
1	A	277	TRP	C-O	-6.00	1.17	1.24
1	A	160	ASP	C-O	-5.99	1.16	1.24
1	A	159	GLU	C-O	-5.96	1.18	1.24
1	A	112	LEU	C-O	-5.94	1.17	1.24
1	A	300	HIS	C-O	-5.92	1.15	1.23
1	A	81	GLU	C-O	-5.90	1.17	1.24
1	A	122	VAL	C-O	-5.89	1.18	1.24
1	A	210	GLU	C-O	-5.85	1.17	1.24
1	A	43	PRO	C-O	-5.83	1.17	1.23
1	A	212	GLN	C-O	-5.82	1.17	1.24
1	A	344	GLU	C-O	-5.79	1.17	1.24
1	A	284	ILE	C-O	-5.78	1.17	1.24
1	A	322	PRO	C-O	-5.77	1.17	1.24
1	A	256	PRO	C-O	-5.77	1.17	1.24
1	A	321	ASN	C-O	-5.76	1.17	1.24
1	A	118	ARG	C-O	-5.75	1.17	1.24
1	A	190	ALA	C-O	-5.73	1.17	1.24
1	A	132	PHE	C-O	-5.72	1.16	1.24
1	A	287	LEU	C-O	-5.70	1.17	1.24
1	A	278	GLN	C-O	-5.69	1.17	1.24
1	A	198	VAL	C-O	-5.68	1.18	1.24
1	A	141	ARG	C-O	-5.67	1.16	1.23
1	A	280	ARG	C-O	-5.65	1.17	1.24
1	A	67	ARG	C-O	-5.61	1.17	1.24
1	A	139	ARG	C-O	-5.59	1.17	1.24
1	A	187	ALA	C-O	-5.59	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	95	VAL	C-O	-5.57	1.18	1.24
1	A	70	PRO	C-O	-5.54	1.17	1.24
1	A	235	ASP	C-O	-5.49	1.16	1.24
1	A	125	VAL	C-O	-5.48	1.18	1.24
1	A	116	VAL	C-O	-5.46	1.17	1.24
1	A	258	SER	C-O	-5.45	1.17	1.23
1	A	211	GLN	C-O	-5.42	1.17	1.24
1	A	336	LEU	C-O	-5.42	1.17	1.24
1	A	184	LEU	C-O	-5.41	1.17	1.24
1	A	153	GLU	C-O	-5.40	1.17	1.24
1	A	182	LEU	C-O	-5.38	1.17	1.24
1	A	77	LYS	C-O	-5.38	1.17	1.24
1	A	293	LYS	C-O	-5.38	1.17	1.24
1	A	350	THR	C-O	-5.35	1.17	1.24
1	A	285	LYS	C-O	-5.35	1.17	1.24
1	A	296	ILE	C-O	-5.34	1.18	1.24
1	A	315	ASP	C-O	-5.32	1.17	1.24
1	A	46	VAL	C-O	-5.30	1.18	1.24
1	A	36	LEU	C-O	-5.29	1.17	1.23
1	A	357	PHE	C-O	-5.24	1.18	1.24
1	A	334	PRO	C-O	-5.24	1.17	1.24
1	A	243	LEU	C-O	-5.23	1.18	1.24
1	A	214	ALA	C-O	-5.23	1.18	1.24
1	A	83	ALA	C-O	-5.22	1.18	1.24
1	A	86	GLY	C-O	-5.20	1.17	1.23
1	A	158	ILE	C-O	-5.20	1.18	1.24
1	A	53	LEU	C-O	-5.19	1.17	1.24
1	A	47	SER	C-O	-5.19	1.17	1.24
1	A	183	VAL	C-O	-5.19	1.18	1.24
1	A	248	TYR	C-O	-5.19	1.17	1.23
1	A	42	GLY	C-O	-5.18	1.17	1.24
1	A	66	LEU	C-O	-5.16	1.18	1.24
1	A	206	GLN	C-O	-5.13	1.17	1.24
1	A	111	ARG	C-O	-5.13	1.18	1.24
1	A	192	LEU	C-O	-5.11	1.18	1.24
1	A	82	LYS	C-O	-5.10	1.18	1.24
1	A	131	TRP	C-O	-5.10	1.18	1.23
1	A	230	HIS	C-O	-5.09	1.16	1.23
1	A	57	HIS	C-O	-5.08	1.18	1.23
1	A	178	PRO	C-O	-5.05	1.17	1.24
1	A	89	HIS	C-O	-5.05	1.18	1.24
1	A	138	MET	C-O	-5.04	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	189	ARG	C-O	-5.04	1.18	1.24
1	A	317	MET	C-O	-5.04	1.18	1.24
1	A	69	TRP	C-O	-5.00	1.18	1.23
1	A	217	GLU	C-O	-5.00	1.17	1.24

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	155	GLN	N-CA-C	8.32	120.43	111.36
1	A	154	ILE	N-CA-C	8.03	118.08	110.53
1	A	358	LEU	N-CA-C	6.84	118.82	111.36
1	A	161	THR	N-CA-C	6.11	120.81	113.23
1	A	229	GLY	N-CA-C	6.05	121.17	112.82
1	A	340	GLY	N-CA-C	5.84	123.67	115.43
1	A	272	PHE	N-CA-C	5.83	120.46	113.23
1	A	226	VAL	N-CA-C	5.43	115.94	108.12
1	A	332	VAL	N-CA-C	5.40	115.61	110.53
1	A	240	LEU	N-CA-C	5.20	116.63	111.07
1	A	304	PHE	CA-C-N	5.17	126.38	121.46
1	A	304	PHE	C-N-CA	5.17	126.38	121.46
1	A	200	THR	N-CA-C	5.16	117.66	109.50
1	A	257	TYR	N-CA-C	5.16	116.67	108.67

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2506	0	2517	23	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	4	0	6	0	0
5	A	155	0	0	6	0
All	All	2667	0	2523	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:361:THR:C	5:A:654:HOH:O	2.29	0.74
1:A:169:KCX:OQ2	1:A:201:HIS:HB2	1.93	0.68
1:A:230:HIS:HD2	1:A:253:ASP:OD1	1.79	0.65
1:A:101:VAL:O	1:A:169:KCX:HD3	1.98	0.64
1:A:361:THR:CA	5:A:654:HOH:O	2.46	0.63
1:A:333:ILE:HB	1:A:334:PRO:HD3	1.84	0.59
1:A:361:THR:HA	5:A:654:HOH:O	2.04	0.56
1:A:101:VAL:HB	1:A:169:KCX:HG2	1.89	0.54
1:A:55:HIS:HD2	5:A:542:HOH:O	1.90	0.53
1:A:131:TRP:CG	1:A:132:PHE:H	2.28	0.52
1:A:290:ARG:HG3	1:A:290:ARG:HH11	1.74	0.52
1:A:37:ILE:HG23	1:A:46:VAL:HG22	1.92	0.51
1:A:200:THR:O	1:A:229:GLY:HA3	2.12	0.48
1:A:106:ILE:HG22	1:A:106:ILE:O	2.14	0.47
1:A:117:SER:HB2	1:A:124:ILE:HD12	1.99	0.44
1:A:302:TRP:CH2	1:A:321:ASN:HB3	2.53	0.43
1:A:344:GLU:HG2	1:A:345:THR:N	2.34	0.43
1:A:272:PHE:CD1	1:A:272:PHE:N	2.84	0.43
1:A:57:HIS:HB2	1:A:303:LEU:HB3	2.01	0.42
1:A:255:MET:N	1:A:256:PRO:CD	2.82	0.42
1:A:327:PHE:CE1	1:A:331:ARG:HG3	2.54	0.42
1:A:169:KCX:OQ1	5:A:614:HOH:O	2.21	0.41
1:A:38:ASN:ND2	5:A:628:HOH:O	2.53	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/329 (99%)	316 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/261 (100%)	256 (98%)	5 (2%)	50	56

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	THR
1	A	200	THR
1	A	262	LEU
1	A	299	SER
1	A	303	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	A	55	HIS
1	A	155	GLN
1	A	230	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	KCX	A	169	2,1,3	10,11,12	0.69	0	6,12,14	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	169	2,1,3	-	7/9/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	169	KCX	N-CA-CB-CG
1	A	169	KCX	C-CA-CB-CG
1	A	169	KCX	OQ1-CX-NZ-CE
1	A	169	KCX	OQ2-CX-NZ-CE
1	A	169	KCX	CG-CD-CE-NZ
1	A	169	KCX	CE-CD-CG-CB
1	A	169	KCX	CA-CB-CG-CD

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	169	KCX	4	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	A	403	-	3,3,3	0.26	0	2,2,2	1.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	403	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/329 (99%)	-0.06	7 (2%) 63 63	11, 17, 36, 63	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	33	THR	5.4
1	A	264	GLY	3.6
1	A	262	LEU	3.3
1	A	261	GLY	3.2
1	A	34	GLY	2.9
1	A	337	ARG	2.4
1	A	361	THR	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	A	169	12/13	0.65	0.23	13,13,21,22	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	A	403	4/4	0.80	0.22	42,43,46,50	4
3	CO	A	402	1/1	0.99	0.03	14,14,14,14	0
2	FE2	A	401	1/1	1.00	0.04	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.