



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:02 PM UTC

PDB ID : 4WM8 / pdb_00004wm8
Title : Crystal Structure of Human Enterovirus D68
Authors : Liu, Y.; Sheng, J.; Fokine, A.; Meng, G.; Long, F.; Kuhn, R.J.; Rossmann, M.G.
Deposited on : 2014-10-08
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	12% 80% 13% 5%
2	B	248	8% 85% 11%
3	C	247	6% 85% 15%
4	D	68	13% 50% 7% 43%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6493 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	0	0	0
			2199	1404	378	408	9			

- Molecule 2 is a protein called VP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	238	Total	C	N	O	S	0	0	0
			1880	1199	328	341	12			


- Molecule 3 is a protein called VP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	247	Total	C	N	O	S	0	0	0
			1901	1197	317	368	19			

- Molecule 4 is a protein called VP4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	39	Total	C	N	O	0	0	0
			297	191	46	60			

- Molecule 5 is DECANOIC ACID (CCD ID: DKA) (formula: C₁₀H₂₀O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			12	10	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	67	Total	O	0	0
			67	67		
6	B	68	Total	O	0	0
			68	68		
6	C	63	Total	O	0	0
			63	63		
6	D	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: VP1



• Molecule 2: VP2




• Molecule 3: VP3



• Molecule 4: VP4





4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	327.00Å 347.80Å 357.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.25 – 2.00 40.25 – 2.00	Depositor EDS
% Data completeness (in resolution range)	55.6 (40.25-2.00) 42.9 (40.25-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.00Å)	Xtriage
Refinement program	CNS 1.3	Depositor
R, R_{free}	0.275 , 0.279 0.256 , 0.256	Depositor DCC
R_{free} test set	37646 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	28.7	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 26.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.65	EDS
Total number of atoms	6493	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DKA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/2250	0.90	10/3060 (0.3%)
2	B	0.37	0/1938	0.91	5/2654 (0.2%)
3	C	0.39	0/1946	0.96	10/2646 (0.4%)
4	D	0.34	0/304	0.83	0/411
All	All	0.37	0/6438	0.92	25/8771 (0.3%)

There are no bond length outliers.

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	101	VAL	N-CA-C	8.69	121.91	111.05
3	C	197	GLN	N-CA-C	-8.05	101.26	112.45
2	B	105	GLY	N-CA-C	-7.04	102.92	112.57
1	A	246	ILE	N-CA-C	6.53	118.80	108.87
3	C	105	TYR	N-CA-C	-6.34	105.29	113.16
2	B	58	THR	N-CA-C	6.28	118.93	111.71
1	A	224	ILE	N-CA-C	-6.27	99.55	108.58
2	B	92	PHE	N-CA-C	-6.21	104.43	111.14
1	A	290	MET	CA-C-N	6.20	127.59	119.84
1	A	290	MET	C-N-CA	6.20	127.59	119.84
3	C	22	ALA	N-CA-C	6.18	117.17	109.57
1	A	116	ASP	N-CA-C	-6.06	101.56	110.52
3	C	95	ARG	N-CA-C	5.92	117.41	111.07
1	A	284	ARG	N-CA-C	-5.91	100.36	109.52
1	A	227	GLU	N-CA-C	-5.64	103.08	110.53
2	B	201	PRO	N-CA-C	5.54	121.05	113.84
3	C	79	GLN	N-CA-C	5.53	118.47	110.28
3	C	52	MET	N-CA-C	5.51	118.22	109.96
1	A	109	LEU	N-CA-C	-5.38	106.48	113.16
3	C	136	PRO	N-CA-C	5.29	118.89	111.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	161	SER	N-CA-C	5.21	118.90	112.54
3	C	8	PRO	N-CA-C	-5.18	102.98	111.21
3	C	152	HIS	N-CA-C	5.06	115.98	108.60
1	A	187	MET	N-CA-C	5.05	118.98	112.41
2	B	127	ILE	N-CA-C	5.02	113.39	107.77

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2199	0	2199	40	0
2	B	1880	0	1815	22	0
3	C	1901	0	1843	35	0
4	D	297	0	286	3	0
5	A	12	0	19	0	0
6	A	67	0	0	0	0
6	B	68	0	0	0	0
6	C	63	0	0	0	0
6	D	6	0	0	0	0
All	All	6493	0	6162	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:151:ARG:O	2:B:151:ARG:HD3	1.80	0.82
2:B:214:VAL:HG22	3:C:64:MET:HE1	1.60	0.81
3:C:89:GLN:HE22	3:C:228:ARG:HH22	1.25	0.80
1:A:142:THR:H	1:A:226:ASN:HD21	1.34	0.75
1:A:278:ASN:HD21	3:C:62:ASN:HD22	1.37	0.72
1:A:278:ASN:HD21	3:C:62:ASN:ND2	1.88	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:53:PRO:HB3	2:B:240:ASN:HD21	1.56	0.70
3:C:89:GLN:NE2	3:C:228:ARG:HH22	1.89	0.70
1:A:275:ASN:HD21	3:C:95:ARG:HH12	1.41	0.69
1:A:278:ASN:ND2	3:C:62:ASN:HD22	1.91	0.67
2:B:198:MET:HE1	2:B:240:ASN:HD22	1.60	0.67
1:A:92:LYS:O	1:A:92:LYS:HD3	1.96	0.66
4:D:60:LEU:HD22	4:D:66:VAL:HG23	1.77	0.66
1:A:52:HIS:NE2	3:C:174:HIS:HD2	1.93	0.65
1:A:77:LYS:H	1:A:89:ASN:HD21	1.43	0.65
1:A:247:LYS:HE3	3:C:33:GLU:OE1	1.95	0.65
1:A:281:ILE:HD13	3:C:54:GLU:CD	2.22	0.64
1:A:276:THR:HG22	1:A:279:ALA:O	1.99	0.63
1:A:142:THR:H	1:A:226:ASN:ND2	1.98	0.60
1:A:224:ILE:HD11	1:A:235:VAL:HG21	1.84	0.58
2:B:220:ARG:HD3	3:C:206:SER:O	2.02	0.58
3:C:89:GLN:HE22	3:C:228:ARG:NH2	1.99	0.58
1:A:278:ASN:CG	3:C:62:ASN:HD22	2.12	0.57
1:A:116:ASP:OD2	1:A:245:HIS:HD2	1.87	0.56
2:B:30:ASN:HD21	4:D:58:GLU:HB2	1.70	0.56
1:A:226:ASN:H	1:A:226:ASN:HD22	1.53	0.55
1:A:201:ALA:HB1	1:A:213:PRO:HG3	1.88	0.55
2:B:129:GLU:OE1	2:B:204:HIS:HE1	1.88	0.55
2:B:214:VAL:HG22	3:C:64:MET:CE	2.34	0.55
2:B:220:ARG:HD2	3:C:207:SER:HA	1.89	0.55
1:A:276:THR:HG23	1:A:279:ALA:H	1.72	0.54
1:A:161:PHE:O	1:A:164:GLN:HG2	2.08	0.53
1:A:245:HIS:HE1	4:D:39:SER:O	1.93	0.52
1:A:184:ILE:HG23	1:A:187:MET:HE3	1.91	0.52
1:A:92:LYS:HD3	1:A:92:LYS:C	2.35	0.52
3:C:50:GLU:HA	3:C:217:ALA:HB2	1.90	0.52
1:A:72:LYS:HZ1	1:A:87:HIS:CE1	2.28	0.52
2:B:151:ARG:HD3	2:B:151:ARG:C	2.34	0.52
1:A:182:MET:HG3	3:C:24:VAL:CG2	2.40	0.52
2:B:53:PRO:HB3	2:B:240:ASN:ND2	2.24	0.50
2:B:167:ILE:HD13	3:C:64:MET:CE	2.41	0.50
1:A:278:ASN:ND2	3:C:62:ASN:ND2	2.55	0.49
2:B:199:ASP:OD2	2:B:204:HIS:HD2	1.94	0.49
1:A:281:ILE:HD12	1:A:281:ILE:N	2.27	0.49
1:A:278:ASN:OD1	3:C:62:ASN:HB3	2.15	0.47
2:B:246:ILE:HG22	2:B:246:ILE:O	2.15	0.47
1:A:184:ILE:CG2	1:A:187:MET:HE3	2.45	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:VAL:HG12	1:A:24:PRO:HD3	1.96	0.46
3:C:179:ASN:ND2	3:C:181:ASP:HB3	2.29	0.46
1:A:116:ASP:OD2	1:A:245:HIS:CD2	2.69	0.45
3:C:179:ASN:HD21	3:C:181:ASP:HB3	1.81	0.45
2:B:27:GLU:HB2	2:B:184:ASN:ND2	2.32	0.45
1:A:65:ARG:HH11	1:A:65:ARG:HG3	1.81	0.45
1:A:278:ASN:OD1	3:C:62:ASN:ND2	2.49	0.45
3:C:9:GLY:O	3:C:12:GLN:HG2	2.16	0.45
2:B:167:ILE:HD13	3:C:64:MET:HE3	1.99	0.44
3:C:128:LYS:HA	3:C:155:TRP:O	2.18	0.44
1:A:208:LEU:HD22	1:A:208:LEU:N	2.33	0.43
1:A:226:ASN:ND2	1:A:226:ASN:H	2.15	0.43
3:C:108:HIS:HD1	3:C:225:ARG:HH21	1.64	0.43
2:B:181:ARG:HG3	3:C:123:PHE:HA	1.99	0.43
3:C:177:MET:SD	3:C:186:ASN:ND2	2.92	0.43
1:A:197:TYR:O	1:A:210:GLY:HA2	2.19	0.43
2:B:215:VAL:HG23	3:C:213:ILE:HD11	2.01	0.42
1:A:76:TYR:HA	1:A:89:ASN:ND2	2.35	0.42
3:C:132:CYS:O	3:C:192:VAL:HA	2.20	0.41
1:A:30:GLU:HA	2:B:177:TRP:HB2	2.03	0.41
2:B:37:GLU:HG3	3:C:37:PRO:HB3	2.03	0.41
3:C:108:HIS:NE2	3:C:177:MET:HE3	2.36	0.41
2:B:95:ASN:O	2:B:99:HIS:HD2	2.03	0.41
1:A:23:VAL:O	1:A:23:VAL:HG23	2.20	0.41
3:C:89:GLN:NE2	3:C:89:GLN:HA	2.36	0.40
2:B:243:ARG:NH2	2:B:246:ILE:HD11	2.35	0.40
1:A:171:VAL:HG11	1:A:182:MET:HE1	2.03	0.40
1:A:69:VAL:HG12	1:A:103:LEU:CD2	2.51	0.40
3:C:50:GLU:CA	3:C:217:ALA:HB2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/297 (93%)	269 (98%)	6 (2%)	0	100	100
2	B	236/248 (95%)	222 (94%)	14 (6%)	0	100	100
3	C	245/247 (99%)	240 (98%)	5 (2%)	0	100	100
4	D	37/68 (54%)	34 (92%)	2 (5%)	1 (3%)	4	1
All	All	793/860 (92%)	765 (96%)	27 (3%)	1 (0%)	48	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	62	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/255 (95%)	242 (100%)	1 (0%)	84	89
2	B	205/212 (97%)	203 (99%)	2 (1%)	68	75
3	C	215/215 (100%)	214 (100%)	1 (0%)	81	87
4	D	32/55 (58%)	32 (100%)	0	100	100
All	All	695/737 (94%)	691 (99%)	4 (1%)	78	85

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	LYS
2	B	88	ASN
2	B	102	TYR
3	C	226	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	89	ASN
1	A	164	GLN
1	A	226	ASN
1	A	245	HIS
1	A	275	ASN
2	B	99	HIS
2	B	130	HIS
2	B	135	HIS
2	B	157	HIS
2	B	204	HIS
2	B	240	ASN
3	C	35	HIS
3	C	62	ASN
3	C	79	GLN
3	C	89	GLN
3	C	174	HIS
3	C	188	ASN
4	D	47	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	DKA	A	900	-	11,11,11	0.74	0	11,11,11	1.17	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DKA	A	900	-	-	2/9/9/9	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
5	A	900	DKA	C3-C2-C1	-2.06	109.13	114.51

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	900	DKA	O1-C1-C2-C3
5	A	900	DKA	O2-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/297 (94%)	1.02	35 (12%) 8 7	19, 26, 47, 66	0
2	B	238/248 (95%)	0.89	21 (8%) 15 14	19, 23, 33, 61	0
3	C	247/247 (100%)	0.89	14 (5%) 29 28	19, 24, 37, 48	0
4	D	39/68 (57%)	1.46	9 (23%) 2 2	22, 33, 82, 86	0
All	All	805/860 (93%)	0.97	79 (9%) 13 12	19, 25, 44, 86	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	29	ILE	4.1
1	A	139	PRO	4.0
1	A	207	GLY	3.9
4	D	62	ALA	3.8
1	A	163	TRP	3.8
1	A	66	ALA	3.7
1	A	175	ILE	3.6
1	A	1	VAL	3.6
1	A	295	VAL	3.6
3	C	217	ALA	3.6
3	C	24	VAL	3.5
4	D	67	LEU	3.4
2	B	186	ALA	3.4
4	D	64	ALA	3.4
2	B	218	GLY	3.4
4	D	52	PHE	3.3
3	C	180	SER	3.3
2	B	114	ALA	3.3
1	A	296	THR	3.2
1	A	67	ALA	3.1
2	B	22	ALA	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	204	SER	3.0
2	B	125	VAL	3.0
4	D	59	GLY	2.9
2	B	30	ASN	2.9
2	B	78	TRP	2.9
1	A	294	ILE	2.8
1	A	87	HIS	2.8
2	B	210	ALA	2.8
3	C	90	LEU	2.8
1	A	200	PHE	2.7
2	B	170	ALA	2.7
2	B	207	TRP	2.7
4	D	38	ALA	2.6
2	B	60	THR	2.5
1	A	220	LEU	2.5
1	A	217	ILE	2.5
3	C	246	TYR	2.4
1	A	192	ALA	2.4
2	B	225	VAL	2.4
2	B	79	TRP	2.4
4	D	31	PHE	2.4
1	A	211	ILE	2.4
1	A	93	TRP	2.4
3	C	231	PRO	2.3
1	A	208	LEU	2.3
3	C	222	PHE	2.3
1	A	97	THR	2.3
1	A	120	THR	2.3
2	B	247	THR	2.3
3	C	87	ASP	2.3
2	B	108	ILE	2.3
3	C	175	TYR	2.3
1	A	95	ILE	2.2
2	B	240	ASN	2.2
1	A	265	ALA	2.2
1	A	101	VAL	2.2
2	B	105	GLY	2.2
2	B	14	LEU	2.2
3	C	129	LEU	2.2
1	A	233	PHE	2.2
1	A	261	SER	2.2
1	A	209	TYR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	28	ALA	2.1
1	A	251	PRO	2.1
3	C	141	PRO	2.1
1	A	74	PHE	2.1
1	A	37	THR	2.1
1	A	154	THR	2.1
1	A	155	PRO	2.1
3	C	130	ILE	2.1
1	A	213	PRO	2.1
1	A	127	VAL	2.1
2	B	10	SER	2.1
1	A	153	LEU	2.0
2	B	192	TRP	2.0
2	B	107	LEU	2.0
4	D	60	LEU	2.0
3	C	138	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	DKA	A	900	12/12	0.96	0.18	62,67,75,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.