



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 10:31 PM UTC

PDB ID : 4WMD / pdb_00004wmd
Title : Crystal structure of catalytically inactive MERS-CoV 3CL protease (C148A)
in spacegroup C2221
Authors : Lountos, G.T.; Needle, D.; Waugh, D.S.
Deposited on : 2014-10-08
Resolution : 2.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4770 (2.60-2.56)
Clashscore	190562	5124 (2.60-2.56)
Ramachandran outliers	187476	5046 (2.60-2.56)
Sidechain outliers	187428	5046 (2.60-2.56)
RSRZ outliers	180081	4770 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	4% 86% 10% ..
1	B	306	5% 87% 12% .
1	C	306	% 88% 10% ..

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7289 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ORF1a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	301	Total	C	N	O	S	0	0	0
			2293	1457	380	432	24			
1	B	304	Total	C	N	O	S	0	1	0
			2319	1473	385	437	24			
1	C	303	Total	C	N	O	S	0	2	0
			2323	1474	388	437	24			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	148	ALA	CYS	engineered mutation	UNP W6A941
B	148	ALA	CYS	engineered mutation	UNP W6A941
C	148	ALA	CYS	engineered mutation	UNP W6A941

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		
2	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		
3	C	1	Total	C	O	0	0
			10	6	4		

- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 5 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			16	10	6		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	84	Total 84	O 84	0	0
6	B	102	Total 102	O 102	0	0
6	C	98	Total 98	O 98	0	0

- Molecule 1: ORF1a



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	81.02Å 168.52Å 250.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.16 – 2.58 46.16 – 2.58	Depositor EDS
% Data completeness (in resolution range)	99.4 (46.16-2.58) 93.2 (46.16-2.58)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.58Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1448)	Depositor
R, R_{free}	0.176 , 0.217 0.184 , 0.221	Depositor DCC
R_{free} test set	2000 reflections (3.72%)	wwPDB-VP
Wilson B-factor (Å ²)	45.9	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7289	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, PGE, 1PE, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/2347	0.88	8/3195 (0.3%)
1	B	0.54	0/2373	0.79	0/3231
1	C	0.54	0/2377	0.80	0/3236
All	All	0.55	0/7097	0.83	8/9662 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	74	ALA	CA-C-N	6.88	127.33	120.52
1	A	74	ALA	C-N-CA	6.88	127.33	120.52
1	A	75	PRO	CA-C-O	-6.82	113.15	122.15
1	A	283	GLY	N-CA-C	6.20	122.60	111.62
1	A	75	PRO	CA-C-N	5.85	132.23	121.70
1	A	75	PRO	C-N-CA	5.85	132.23	121.70
1	A	224	LYS	CA-C-N	-5.17	114.42	119.64
1	A	224	LYS	C-N-CA	-5.17	114.42	119.64

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	282	LEU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2293	0	2241	17	0
1	B	2319	0	2267	23	0
1	C	2323	0	2271	15	0
2	A	7	0	10	0	0
2	B	7	0	10	1	0
2	C	7	0	10	1	0
3	A	10	0	14	2	0
3	C	10	0	14	1	0
4	B	13	0	18	1	0
5	C	16	0	22	1	0
6	A	84	0	0	2	0
6	B	102	0	0	2	0
6	C	98	0	0	0	0
All	All	7289	0	6877	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:52:PRO:HD3	1:B:191:LYS:HG3	1.70	0.74
1:C:145:CYS:HA	3:C:403:PGE:H22	1.78	0.66
1:B:40:ARG:HA	1:B:90:LEU:HG	1.84	0.60
1:C:216:LEU:HD13	1:C:300:ILE:HG21	1.86	0.58
1:A:276:PHE:O	1:A:279:LYS:HB2	2.05	0.57
1:B:279:LYS:NZ	6:B:568:HOH:O	2.39	0.56
1:B:70:LYS:NZ	1:B:72:ILE:O	2.38	0.54
1:B:194:HIS:HD2	1:B:195:GLN:H	1.56	0.54
1:C:301:MET:HE2	1:C:303:VAL:HG21	1.89	0.54
1:C:105:THR:OG1	1:C:161[B]:ASN:ND2	2.32	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:402:PGE:H12	6:A:568:HOH:O	2.09	0.52
1:A:40:ARG:HA	1:A:90:LEU:HG	1.92	0.52
1:C:216:LEU:HD21	1:C:257:LYS:HG3	1.93	0.51
1:B:200:ASP:HB3	2:B:401:PEG:H22	1.93	0.51
1:B:194:HIS:CD2	1:B:195:GLN:N	2.80	0.50
1:A:283:GLY:HA2	1:A:284:SER:CB	2.40	0.50
1:A:111:PRO:HB3	1:A:135:PRO:HA	1.93	0.50
1:B:166:HIS:HE1	1:B:175:HIS:HB3	1.77	0.49
1:A:191:LYS:HG3	1:A:193:VAL:HG12	1.94	0.49
1:B:189:MET:N	1:B:195:GLN:OE1	2.23	0.48
1:A:280:GLN:HB3	1:A:283:GLY:HA3	1.95	0.48
3:A:402:PGE:H42	6:A:568:HOH:O	2.12	0.48
1:B:57:LEU:O	1:B:61:MET:HG2	2.13	0.48
5:C:401:1PE:H242	5:C:401:1PE:H252	1.53	0.48
1:A:278:GLY:HA3	1:A:279:LYS:HG2	1.94	0.48
1:C:111:PRO:HB3	1:C:135:PRO:HA	1.95	0.48
1:A:283:GLY:HA2	1:A:284:SER:HB3	1.96	0.47
1:A:72:ILE:HG12	1:A:125:PRO:HD2	1.97	0.47
1:B:70:LYS:NZ	6:B:536:HOH:O	2.46	0.47
1:C:43:MET:HB3	1:C:61:MET:HE3	1.97	0.46
1:B:190:ASP:OD1	1:B:190:ASP:N	2.46	0.46
1:B:189:MET:HE2	1:B:195:GLN:HG2	1.97	0.46
1:A:129:PHE:CG	1:B:4:VAL:HG21	2.52	0.45
1:C:227:ARG:HB3	1:C:263:GLU:HB3	1.98	0.45
1:B:74:ALA:HA	1:B:75:PRO:HD3	1.84	0.45
1:C:202:TYR:CE2	2:C:402:PEG:H22	2.51	0.45
1:A:166:HIS:HE1	1:A:175:HIS:HB3	1.81	0.45
1:B:6:MET:HE2	1:B:6:MET:HB2	1.85	0.44
1:B:12:ASP:HB3	1:B:160:ILE:HD11	1.99	0.44
1:C:40:ARG:HA	1:C:90:LEU:HG	2.00	0.44
1:A:176:THR:OG1	1:A:177:GLY:N	2.51	0.43
1:C:298:MET:HE2	1:C:303:VAL:O	2.18	0.43
1:C:286:MET:HE3	1:C:286:MET:HB3	1.66	0.43
1:A:74:ALA:HA	1:A:75:PRO:HD3	1.93	0.43
1:A:286:MET:HG3	1:B:285:THR:HG21	2.00	0.43
1:A:215:ILE:HD13	1:A:260:VAL:HG21	2.01	0.42
1:C:73:GLY:O	1:C:75:PRO:HD3	2.20	0.42
1:A:253:MET:HE2	1:A:253:MET:HB3	1.79	0.42
1:B:132:VAL:HG21	1:B:290:GLU:HA	2.00	0.42
1:B:116:SER:O	1:B:152:GLY:HA2	2.19	0.42
1:C:31:TRP:CE2	1:C:98:ASN:HB2	2.55	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:298:MET:CG	4:B:402:PG4:H32	2.50	0.41
1:B:284:SER:OG	1:B:286:MET:HE3	2.21	0.41
1:C:44:CYS:SG	1:C:52:PRO:HB3	2.61	0.41
1:B:167:GLN:HG3	1:B:184:MET:SD	2.61	0.40
1:A:44:CYS:SG	1:A:52:PRO:HB3	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/306 (98%)	283 (95%)	14 (5%)	2 (1%)	18	36
1	B	303/306 (99%)	290 (96%)	12 (4%)	1 (0%)	36	56
1	C	303/306 (99%)	298 (98%)	5 (2%)	0	100	100
All	All	905/918 (99%)	871 (96%)	31 (3%)	3 (0%)	36	56

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	76	ALA
1	A	284	SER
1	B	52	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/256 (98%)	243 (96%)	9 (4%)	31	56
1	B	255/256 (100%)	248 (97%)	7 (3%)	39	65
1	C	255/256 (100%)	245 (96%)	10 (4%)	28	53
All	All	762/768 (99%)	736 (97%)	26 (3%)	32	58

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	THR
1	A	47	ASP
1	A	59	ILE
1	A	62	THR
1	A	72	ILE
1	A	151	VAL
1	A	226	ASN
1	A	231	VAL
1	A	284	SER
1	B	26	THR
1	B	27	LEU
1	B	47	ASP
1	B	107	THR
1	B	151	VAL
1	B	196	VAL
1	B	197	GLN
1	C	27	LEU
1	C	59	ILE
1	C	69	GLN
1	C	72	ILE
1	C	196	VAL
1	C	277	GLN
1	C	280	GLN
1	C	286	MET
1	C	297	ASN
1	C	298	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	HIS
1	A	41	HIS
1	A	217	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	122	ASN
1	B	194	HIS
1	C	77	ASN
1	C	277	GLN
1	C	297	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PGE	C	403	-	9,9,9	1.02	0	8,8,8	0.49	0
2	PEG	C	402	-	6,6,6	0.65	0	5,5,5	0.42	0
2	PEG	A	401	-	6,6,6	0.67	0	5,5,5	0.36	0
5	1PE	C	401	-	15,15,15	0.79	0	14,14,14	0.49	0
3	PGE	A	402	-	9,9,9	1.06	0	8,8,8	0.78	0
4	PG4	B	402	-	12,12,12	1.00	0	11,11,11	0.56	0
2	PEG	B	401	-	6,6,6	0.62	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	C	403	-	-	3/7/7/7	-
2	PEG	C	402	-	-	3/4/4/4	-
2	PEG	A	401	-	-	3/4/4/4	-
5	1PE	C	401	-	-	8/13/13/13	-
3	PGE	A	402	-	-	4/7/7/7	-
4	PG4	B	402	-	-	5/10/10/10	-
2	PEG	B	401	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	401	1PE	C24-C14-OH5-C25
3	A	402	PGE	O2-C3-C4-O3
5	C	401	1PE	OH4-C13-C23-OH3
3	A	402	PGE	O1-C1-C2-O2
5	C	401	1PE	OH5-C14-C24-OH4
3	C	403	PGE	O2-C3-C4-O3
3	C	403	PGE	O1-C1-C2-O2
4	B	402	PG4	O2-C3-C4-O3
2	B	401	PEG	C4-C3-O2-C2
2	A	401	PEG	O2-C3-C4-O4
5	C	401	1PE	OH6-C15-C25-OH5
4	B	402	PG4	O1-C1-C2-O2
5	C	401	1PE	OH7-C16-C26-OH6
2	A	401	PEG	O1-C1-C2-O2
2	B	401	PEG	O1-C1-C2-O2
2	B	401	PEG	O2-C3-C4-O4
2	C	402	PEG	O1-C1-C2-O2
3	C	403	PGE	O3-C5-C6-O4
4	B	402	PG4	O4-C7-C8-O5
5	C	401	1PE	C12-C22-OH3-C23
5	C	401	1PE	C16-C26-OH6-C15
4	B	402	PG4	C3-C4-O3-C5
5	C	401	1PE	C23-C13-OH4-C24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	C	402	PEG	C1-C2-O2-C3
2	C	402	PEG	C4-C3-O2-C2
2	A	401	PEG	C4-C3-O2-C2
4	B	402	PG4	C8-C7-O4-C6
3	A	402	PGE	C1-C2-O2-C3
3	A	402	PGE	C4-C3-O2-C2

There are no ring outliers.

6 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	403	PGE	1	0
2	C	402	PEG	1	0
5	C	401	1PE	1	0
3	A	402	PGE	2	0
4	B	402	PG4	1	0
2	B	401	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	301/306 (98%)	-0.04	11 (3%) 45 40	35, 49, 78, 93	0
1	B	304/306 (99%)	0.03	15 (4%) 35 30	21, 50, 86, 106	1 (0%)
1	C	303/306 (99%)	-0.23	2 (0%) 84 82	21, 45, 71, 84	3 (0%)
All	All	908/918 (98%)	-0.08	28 (3%) 51 46	21, 49, 79, 106	4 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	GLY	5.0
1	B	196	VAL	4.5
1	B	171	ALA	4.2
1	B	304	VAL	4.2
1	B	194	HIS	3.7
1	C	303	VAL	3.6
1	A	76	ALA	3.6
1	B	46	ALA	3.6
1	A	301	MET	3.6
1	B	76	ALA	3.4
1	C	301	MET	3.2
1	A	75	PRO	3.2
1	B	193	VAL	2.9
1	A	278	GLY	2.7
1	A	283	GLY	2.7
1	B	195	GLN	2.6
1	B	303	VAL	2.5
1	B	75	PRO	2.4
1	A	286	MET	2.4
1	B	274	THR	2.4
1	B	64	HIS	2.4
1	B	172	ASN	2.3
1	A	1	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	48	GLN	2.2
1	B	74	ALA	2.1
1	A	72	ILE	2.1
1	A	46	ALA	2.1
1	A	64	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PGE	A	402	10/10	0.77	0.30	50,67,78,81	0
2	PEG	A	401	7/7	0.82	0.18	55,59,71,75	0
3	PGE	C	403	10/10	0.83	0.22	52,61,71,72	0
5	1PE	C	401	16/16	0.83	0.20	57,69,76,80	0
2	PEG	C	402	7/7	0.88	0.15	53,56,63,66	0
2	PEG	B	401	7/7	0.88	0.14	52,58,69,75	0
4	PG4	B	402	13/13	0.91	0.16	45,60,66,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.