



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2026 – 01:50 AM UTC

PDB ID : 5WM8 / pdb_00005wm8
Title : Structure of the 10R (+)-cis-BP-dG modified Rev1 ternary complex
Authors : Rechkoblit, O.; Kolbanovsky, A.; Landes, H.; Geacintov, N.E.; Aggarwal, A.K.
Deposited on : 2017-07-28
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	12	 92% 8%
2	T	17	 88% 12%
3	A	434	 5% 97% .

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 4514 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*GP*GP*GP*GP*TP*GP*TP*GP*GP*TP*AP*(DDG))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	P	12	Total	C	N	O	P	0	2	0
			274	130	51	80	13			

- Molecule 2 is a DNA chain called DNA (5'-D(P*AP*TP*CP*GP*CP*TP*AP*CP*CP*AP*CP*AP*CP*CP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	17	Total	C	N	O	P	0	0	1
			318	151	56	95	16			


- Molecule 3 is a protein called DNA repair protein REV1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	432	Total	C	N	O	S	0	9	0
			3484	2217	594	647	26			

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	P	1	Total	Mg	0	0
			1	1		
4	A	3	Total	Mg	0	0
			3	3		

- Molecule 5 is 1,2,3-TRIHIDROXY-1,2,3,4-TETRAHYDROBENZO[A]PYRENE (CCD ID: BAP) (formula: C₂₀H₁₆O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	T	1	Total	C	O	0	0
			23	20	3		

- Molecule 6 is 2'-DEOXYCYTIDINE-5'-TRIPHOSPHATE (CCD ID: DCP) (formula: $C_9H_{16}N_3O_{13}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			28	9	3	13	3		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 8 4 4	0	1
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 4 2 2	0	0

- Molecule 9 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total C O 7 4 3	0	0
9	A	1	Total C O 7 4 3	0	0

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	P	25	Total 25	O 25	0	0
10	T	19	Total 19	O 19	0	0
10	A	267	Total 267	O 267	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*GP*GP*GP*GP*TP*GP*TP*GP*GP*TP*AP*(DDG))-3')

Chain P:  92% 8%





- Molecule 2: DNA (5'-D(P*AP*TP*CP*GP*CP*TP*AP*CP*CP*AP*CP*AP*CP*CP*CP*C)-3')

Chain T:  88% 12%



- Molecule 3: DNA repair protein REV1

Chain A:  5% 97%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	62.55Å 180.36Å 54.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.00 – 1.92 45.00 – 1.93	Depositor EDS
% Data completeness (in resolution range)	95.0 (45.00-1.92) 95.0 (45.00-1.93)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.176 , 0.214 0.184 , 0.218	Depositor DCC
R_{free} test set	2290 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4514	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DDG, DCP, BAP, EDO, MG, PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	P	0.32	0/284	0.78	0/437
2	T	0.55	1/354 (0.3%)	0.82	0/542
3	A	0.65	0/3569	0.82	0/4818
All	All	0.62	1/4207 (0.0%)	0.82	0/5797

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	1	DC	O3'-P	-7.93	1.49	1.61

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	274	0	144	0	0
2	T	318	0	178	1	0
3	A	3484	0	3496	10	0
4	A	3	0	0	0	0
4	P	1	0	0	0	0
5	T	23	0	15	0	0
6	A	28	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	18	0	24	0	0
8	A	40	0	60	1	0
9	A	14	0	20	0	0
10	A	267	0	0	2	0
10	P	25	0	0	0	0
10	T	19	0	0	1	0
All	All	4514	0	3949	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:457:ILE:CD1	3:A:489:LEU:HD11	2.16	0.74
3:A:457:ILE:HD12	3:A:489:LEU:HD21	1.82	0.62
3:A:457:ILE:HD12	3:A:489:LEU:HD11	1.90	0.52
3:A:495:GLN:OE1	10:A:901:HOH:O	2.19	0.52
10:T:201:HOH:O	8:A:815:EDO:H22	2.10	0.50
3:A:669:MET:HE3	3:A:688:CYS:HB2	1.97	0.46
3:A:648[B]:GLN:NE2	10:A:905:HOH:O	2.48	0.46
2:T:2:DA:N1	3:A:307:ILE:HD11	2.31	0.45
3:A:457:ILE:HD13	3:A:489:LEU:HD11	1.96	0.45
3:A:364:ASP:O	3:A:365:CYS:C	2.63	0.42
3:A:455:LEU:HD13	3:A:457:ILE:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	439/434 (101%)	432 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	394/391 (101%)	391 (99%)	3 (1%)	73	71

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	362	ASP
3	A	382	CYS
3	A	722	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	495	GLN
3	A	499	GLN
3	A	619	GLN
3	A	733	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	DDG	P	12[B]	1	20,23,24	1.27	4 (20%)	27,33,36	2.05	8 (29%)
1	DDG	P	12[A]	1	20,23,24	1.27	4 (20%)	27,33,36	1.91	6 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DDG	P	12[B]	1	-	0/7/18/19	0/3/3/3
1	DDG	P	12[A]	1	-	0/7/18/19	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	P	12[A]	DDG	C6-N1	-2.68	1.33	1.38
1	P	12[B]	DDG	C6-N1	-2.68	1.33	1.38
1	P	12[A]	DDG	C5-C4	2.55	1.45	1.38
1	P	12[B]	DDG	C5-C4	2.55	1.45	1.38
1	P	12[A]	DDG	C4-N9	-2.37	1.32	1.38
1	P	12[B]	DDG	C4-N9	-2.37	1.32	1.38
1	P	12[A]	DDG	C5-N7	-2.20	1.34	1.39
1	P	12[B]	DDG	C5-N7	-2.20	1.34	1.39

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	P	12[A]	DDG	C5-C4-N3	-4.59	121.08	128.39
1	P	12[B]	DDG	C5-C4-N3	-4.59	121.08	128.39
1	P	12[A]	DDG	C2-N3-C4	4.31	119.72	112.30
1	P	12[B]	DDG	C2-N3-C4	4.31	119.72	112.30
1	P	12[A]	DDG	N9-C4-N3	3.46	132.88	125.95
1	P	12[B]	DDG	N9-C4-N3	3.46	132.88	125.95
1	P	12[A]	DDG	C6-C5-N7	3.04	135.83	130.29
1	P	12[B]	DDG	C6-C5-N7	3.04	135.83	130.29
1	P	12[A]	DDG	O6-C6-C5	-2.95	118.75	126.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	P	12[B]	DDG	O6-C6-C5	-2.95	118.75	126.53
1	P	12[B]	DDG	C2'-C3'-C4'	2.69	107.75	102.75
1	P	12[B]	DDG	O4'-C1'-C2'	2.62	109.51	106.41
1	P	12[A]	DDG	C5-C6-N1	2.04	118.45	113.25
1	P	12[B]	DDG	C5-C6-N1	2.04	118.45	113.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 4 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	GOL	A	805	-	5,5,5	0.42	0	5,5,5	0.68	0
8	EDO	A	810	-	3,3,3	0.45	0	2,2,2	0.52	0
8	EDO	A	815	-	3,3,3	0.51	0	2,2,2	0.17	0
7	GOL	A	806	-	5,5,5	0.29	0	5,5,5	0.23	0
8	EDO	A	812[B]	-	3,3,3	0.38	0	2,2,2	0.54	0
8	EDO	A	812[A]	-	3,3,3	0.32	0	2,2,2	0.48	0
8	EDO	A	811	-	3,3,3	0.37	0	2,2,2	0.46	0
7	GOL	A	807	-	5,5,5	0.45	0	5,5,5	0.50	0
8	EDO	A	816	-	3,3,3	0.44	0	2,2,2	0.33	0
9	PEG	A	817	-	6,6,6	0.47	0	5,5,5	0.26	0
9	PEG	A	818	-	6,6,6	0.48	0	5,5,5	0.25	0
8	EDO	A	813	-	3,3,3	0.41	0	2,2,2	0.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	EDO	A	808	-	3,3,3	0.45	0	2,2,2	0.08	0
5	BAP	T	101	2	27,27,27	2.25	10 (37%)	37,42,42	0.91	2 (5%)
8	EDO	A	814	-	3,3,3	0.44	0	2,2,2	0.16	0
8	EDO	A	809	-	3,3,3	0.43	0	2,2,2	0.29	0
6	DCP	A	804	4	28,29,29	1.39	4 (14%)	41,45,45	1.71	11 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	805	-	-	0/4/4/4	-
8	EDO	A	810	-	-	0/1/1/1	-
8	EDO	A	815	-	-	0/1/1/1	-
7	GOL	A	806	-	-	3/4/4/4	-
8	EDO	A	812[B]	-	-	0/1/1/1	-
8	EDO	A	812[A]	-	-	1/1/1/1	-
8	EDO	A	811	-	-	1/1/1/1	-
7	GOL	A	807	-	-	1/4/4/4	-
8	EDO	A	816	-	-	1/1/1/1	-
9	PEG	A	817	-	-	2/4/4/4	-
9	PEG	A	818	-	-	1/4/4/4	-
8	EDO	A	813	-	-	1/1/1/1	-
8	EDO	A	808	-	-	0/1/1/1	-
5	BAP	T	101	2	-	-	0/5/5/5
8	EDO	A	814	-	-	1/1/1/1	-
8	EDO	A	809	-	-	1/1/1/1	-
6	DCP	A	804	4	-	5/22/34/34	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	T	101	BAP	C6-C7	7.69	1.49	1.37
6	A	804	DCP	PB-O3A	4.74	1.64	1.59
5	T	101	BAP	C6-C5A	3.12	1.49	1.43
6	A	804	DCP	PA-O3A	3.05	1.62	1.59
5	T	101	BAP	C5B-C8A	2.94	1.48	1.42
5	T	101	BAP	C5A-C5B	2.76	1.48	1.42
6	A	804	DCP	PB-O3B	2.68	1.62	1.59
5	T	101	BAP	C7-C1'	-2.57	1.48	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	T	101	BAP	C8-C7	2.56	1.40	1.36
5	T	101	BAP	C3B-C1A	2.54	1.47	1.42
5	T	101	BAP	C3B-C3A	2.44	1.47	1.42
5	T	101	BAP	C10-C9	2.36	1.41	1.35
6	A	804	DCP	C6-C5	2.25	1.40	1.35
5	T	101	BAP	C2-C1	2.04	1.41	1.36

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	804	DCP	O3G-PG-O2G	3.52	121.01	107.80
6	A	804	DCP	O3A-PA-O1A	-3.39	100.51	110.70
6	A	804	DCP	O3A-PB-O1B	-3.39	100.51	110.70
6	A	804	DCP	O2A-PA-O5'	3.24	122.26	107.57
6	A	804	DCP	O2A-PA-O1A	2.92	126.05	112.44
5	T	101	BAP	O1'-C1'-C7	-2.84	105.01	110.42
6	A	804	DCP	O2B-PB-O1B	2.78	125.36	112.44
6	A	804	DCP	O3B-PB-O1B	-2.65	102.72	110.70
6	A	804	DCP	O5'-PA-O1A	-2.59	98.67	108.94
6	A	804	DCP	C2'-C1'-N1	-2.37	107.88	113.81
5	T	101	BAP	C5A-C5B-C8A	2.33	122.93	119.91
6	A	804	DCP	O3B-PG-O1G	-2.31	98.88	111.04
6	A	804	DCP	N4-C4-N3	2.08	121.63	117.91

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	804	DCP	PB-O3B-PG-O3G
7	A	806	GOL	C1-C2-C3-O3
7	A	806	GOL	O2-C2-C3-O3
8	A	809	EDO	O1-C1-C2-O2
8	A	812[A]	EDO	O1-C1-C2-O2
8	A	816	EDO	O1-C1-C2-O2
9	A	817	PEG	O2-C3-C4-O4
6	A	804	DCP	PA-O3A-PB-O1B
8	A	813	EDO	O1-C1-C2-O2
9	A	818	PEG	O2-C3-C4-O4
8	A	814	EDO	O1-C1-C2-O2
6	A	804	DCP	PA-O3A-PB-O2B
8	A	811	EDO	O1-C1-C2-O2
6	A	804	DCP	PB-O3B-PG-O1G

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	A	804	DCP	PB-O3B-PG-O2G
7	A	807	GOL	O1-C1-C2-O2
7	A	806	GOL	O1-C1-C2-C3
9	A	817	PEG	C1-C2-O2-C3


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	815	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	P	11/12 (91%)	0.10	0	100	100	17, 41, 54, 54	1 (9%)
2	T	17/17 (100%)	0.02	0	100	100	20, 41, 55, 59	0
3	A	432/434 (99%)	0.24	23 (5%)	32	36	11, 29, 51, 84	9 (2%)
All	All	460/463 (99%)	0.23	23 (5%)	34	38	11, 29, 54, 84	10 (2%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	738	VAL	6.1
3	A	348	THR	5.0
3	A	479	ILE	4.2
3	A	347	TYR	3.6
3	A	344	ILE	3.4
3	A	376	SER	3.1
3	A	457	ILE	3.0
3	A	378	SER	2.9
3	A	379	PHE	2.7
3	A	380	SER	2.6
3	A	307	ILE	2.5
3	A	377	SER	2.5
3	A	675	ALA	2.4
3	A	382	CYS	2.3
3	A	476	PRO	2.2
3	A	555	HIS	2.2
3	A	569	HIS	2.2
3	A	674	ASP	2.2
3	A	540	GLU	2.2
3	A	699	ILE	2.1
3	A	350	ILE	2.0
3	A	481	ASN	2.0
3	A	381	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	DDG	P	12[A]	21/22	0.97	0.05	21,24,31,33	10
1	DDG	P	12[B]	21/22	0.97	0.05	19,22,24,24	10

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	PEG	A	818	7/7	0.69	0.24	69,74,77,77	0
8	EDO	A	814	4/4	0.75	0.26	44,49,49,52	0
8	EDO	A	815	4/4	0.82	0.17	47,49,50,50	0
8	EDO	A	809	4/4	0.84	0.16	52,53,54,54	0
9	PEG	A	817	7/7	0.86	0.14	54,56,57,61	0
8	EDO	A	811	4/4	0.86	0.12	37,42,44,45	0
4	MG	A	801	1/1	0.87	0.12	29,29,29,29	0
4	MG	A	803	1/1	0.87	0.17	22,22,22,22	0
8	EDO	A	816	4/4	0.88	0.14	59,61,62,62	0
8	EDO	A	810	4/4	0.88	0.12	33,36,36,40	0
7	GOL	A	805	6/6	0.88	0.12	35,37,37,40	0
8	EDO	A	812[B]	4/4	0.89	0.17	45,46,47,48	4
8	EDO	A	812[A]	4/4	0.89	0.17	25,26,27,28	4
4	MG	P	101	1/1	0.90	0.09	30,30,30,30	0
7	GOL	A	806	6/6	0.91	0.11	48,50,51,52	0
7	GOL	A	807	6/6	0.91	0.09	36,40,41,44	0
8	EDO	A	813	4/4	0.91	0.13	30,35,36,41	0
8	EDO	A	808	4/4	0.91	0.12	31,34,35,40	0
5	BAP	T	101	23/23	0.93	0.10	30,40,43,44	0
4	MG	A	802	1/1	0.96	0.06	16,16,16,16	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	DCP	A	804	28/28	0.98	0.05	17,18,25,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.