



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 03:02 PM UTC

PDB ID : 5WMQ / pdb_00005wmq
Title : Crystal Structure of HLA-B8 in complex with ELR, an Influenza A virus peptide
Authors : Gras, S.; Rossjohn, J.
Deposited on : 2017-07-31
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	13% 89% 10% .
2	B	100	10% 87% 11% ..
3	C	9	78% 22%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3718 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, B-8 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	0	12	0
			2315	1436	418	452	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	45	MET	GLU	conflict	UNP P30460

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	4	0
			856	545	145	163	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called ELR peptide from IAV.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	1	0
			90	60	16	14			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	338	Total	O	0	0
			338	338		
4	B	104	Total	O	0	0
			104	104		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	15	Total	O	0	0
			15	15		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, B-8 alpha chain



- Molecule 2: Beta-2-microglobulin



- Molecule 3: ELR peptide from IAV



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.53Å 38.55Å 128.09Å 90.00° 101.78° 90.00°	Depositor
Resolution (Å)	20.50 – 1.40 20.50 – 1.40	Depositor EDS
% Data completeness (in resolution range)	97.0 (20.50-1.40) 97.0 (20.50-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.30 (at 1.40Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.199 , 0.216 0.208 , 0.229	Depositor DCC
R_{free} test set	4104 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	9.7	Xtriage
Anisotropy	0.537	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3718	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	1/2401 (0.0%)	1.06	4/3264 (0.1%)
2	B	0.75	0/883	1.07	4/1197 (0.3%)
3	C	0.83	0/95	0.98	0/126
All	All	0.79	1/3379 (0.0%)	1.06	8/4587 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	145	ARG	CZ-NH2	-5.28	1.26	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	37	ASP	CA-CB-CG	6.53	119.13	112.60
2	B	33	SER	N-CA-C	5.73	118.46	111.82
2	B	70	PHE	CA-CB-CG	5.48	119.28	113.80
1	A	13	SER	N-CA-C	-5.36	103.31	110.55
1	A	176	LYS	N-CA-C	5.31	117.76	111.33
1	A	129	ASP	CA-CB-CG	5.22	117.82	112.60
2	B	21	ASN	CA-CB-CG	5.09	117.69	112.60
2	B	96	ASP	CA-CB-CG	5.01	117.61	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2315	0	2171	19	0
2	B	856	0	820	9	0
3	C	90	0	96	2	0
4	A	338	0	0	8	0
4	B	104	0	0	0	0
4	C	15	0	0	0	0
All	All	3718	0	3087	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94[B]:THR:HG21	2:B:31[B]:HIS:HE1	1.40	0.87
1:A:93:HIS:HD2	1:A:119:ASP:OD2	1.74	0.69
1:A:12[A]:MET:HE3	4:A:611:HOH:O	1.93	0.68
2:B:17:ASN:HD21	2:B:97:ARG:HH22	1.41	0.67
1:A:94[B]:THR:HG21	2:B:31[B]:HIS:CE1	2.26	0.66
4:A:301:HOH:O	3:C:5:ARG:NH1	2.31	0.63
1:A:93:HIS:HE1	4:A:430:HOH:O	1.86	0.59
1:A:68:LYS:HG3	4:A:517:HOH:O	2.05	0.56
2:B:17:ASN:HD21	2:B:97:ARG:NH2	2.04	0.55
1:A:86:ASN:ND2	4:A:303:HOH:O	2.39	0.54
1:A:113:HIS:HD2	4:A:581:HOH:O	1.91	0.54
1:A:94[B]:THR:HG22	1:A:119:ASP:HA	1.91	0.53
2:B:17:ASN:ND2	2:B:97:ARG:HH22	2.08	0.50
1:A:54:GLN:HE22	1:A:174:ASN:HB3	1.77	0.50
1:A:35:ARG:CZ	2:B:53:ASP:HB3	2.44	0.48
1:A:127:ASN:HD21	1:A:134:THR:HB	1.78	0.48
2:B:42:ASN:ND2	2:B:77:GLU:H	2.12	0.48
1:A:102:ASP:OD2	1:A:113:HIS:HE1	1.97	0.47
4:A:321:HOH:O	3:C:2[A]:LEU:HD22	2.16	0.45
1:A:201:LEU:HD12	1:A:249:VAL:HG11	2.00	0.43
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.55	0.42
1:A:126:LEU:HG	1:A:130:LEU:HA	2.02	0.42
1:A:97[A]:SER:OG	4:A:301:HOH:O	2.21	0.41
1:A:111:ARG:HG2	1:A:113:HIS:CE1	2.56	0.41
2:B:42:ASN:HD21	2:B:77:GLU:H	1.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/276 (104%)	281 (98%)	5 (2%)	0	100	100
2	B	101/100 (101%)	100 (99%)	1 (1%)	0	100	100
3	C	8/9 (89%)	8 (100%)	0	0	100	100
All	All	395/385 (103%)	389 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/235 (105%)	240 (97%)	7 (3%)	38	9
2	B	98/95 (103%)	96 (98%)	2 (2%)	48	17
3	C	9/8 (112%)	9 (100%)	0	100	100
All	All	354/338 (105%)	345 (98%)	9 (2%)	39	11

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	71	THR
1	A	137	ASP
1	A	145	ARG
1	A	222	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	223	ASP
1	A	225	THR
2	B	47	GLU
2	B	70	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	86	ASN
1	A	93	HIS
1	A	113	HIS
1	A	127	ASN
1	A	141	GLN
1	A	155	GLN
1	A	218	GLN
1	A	260	HIS
1	A	262	GLN
2	B	17	ASN
2	B	42	ASN
2	B	89	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	276/276 (100%)	0.41	37 (13%)	7 7	4, 13, 49, 62	14 (5%)
2	B	99/100 (99%)	0.89	10 (10%)	12 12	5, 21, 40, 46	5 (5%)
3	C	9/9 (100%)	-0.78	0	100 100	3, 5, 9, 9	1 (11%)
All	All	384/385 (99%)	0.51	47 (12%)	8 9	3, 15, 42, 62	20 (5%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	ILE	6.1
1	A	276	PRO	4.7
1	A	274	TRP	4.1
1	A	218	GLN	3.9
1	A	223	ASP	3.7
1	A	268	LYS	3.7
1	A	226	GLN	3.5
1	A	44	ARG	3.4
1	A	196[A]	ASP	3.3
1	A	255	GLN	3.3
1	A	194	ILE	3.3
2	B	48	LYS	3.2
1	A	273	ARG	3.2
1	A	181	ARG	3.2
1	A	221	GLY	3.1
2	B	75	LYS	3.1
1	A	43	PRO	3.0
1	A	225	THR	2.8
1	A	45	MET	2.8
1	A	182	ALA	2.7
2	B	39	LEU	2.6
1	A	41	ALA	2.6
1	A	222	GLU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	47	GLU	2.6
2	B	20	SER	2.6
1	A	269	PRO	2.6
1	A	270	LEU	2.5
1	A	258	THR	2.4
1	A	260	HIS	2.4
2	B	19	LYS	2.4
1	A	272	LEU	2.3
2	B	72	PRO	2.3
1	A	224	GLN	2.3
1	A	15	PRO	2.3
1	A	249	VAL	2.3
1	A	217	TRP	2.3
1	A	215	LEU	2.3
1	A	230	LEU	2.3
1	A	42	SER	2.3
1	A	266	LEU	2.2
1	A	197	HIS	2.2
1	A	54	GLN	2.2
1	A	16	GLY	2.1
1	A	257	TYR	2.1
2	B	22	PHE	2.0
2	B	71	THR	2.0
1	A	261	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.