



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:13 PM UTC

PDB ID : 6WMH / pdb_00006wmh
Title : Next generation monomeric IgG4 Fc
Authors : Oganessian, V.Y.; Shan, L.; van Dyk, N.; Dall'Acqua, W.F.
Deposited on : 2020-04-21
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	
1	H	207	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3496 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Immunoglobulin heavy constant gamma 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	207	Total	C	N	O	S	0	0	0
			1674	1058	283	327	6			
1	A	206	Total	C	N	O	S	0	0	0
			1668	1055	282	325	6			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	252	TYR	MET	engineered mutation	UNP P01861
H	254	THR	SER	engineered mutation	UNP P01861
H	256	GLU	THR	engineered mutation	UNP P01861
H	297	ASP	ASN	engineered mutation	UNP P01861
H	351	PHE	LEU	engineered mutation	UNP P01861
H	354	GLU	SER	engineered mutation	UNP P01861
H	366	ARG	THR	engineered mutation	UNP P01861
H	395	LYS	PRO	engineered mutation	UNP P01861
H	405	ARG	PHE	engineered mutation	UNP P01861
H	407	GLU	TYR	engineered mutation	UNP P01861
A	252	TYR	MET	engineered mutation	UNP P01861
A	254	THR	SER	engineered mutation	UNP P01861
A	256	GLU	THR	engineered mutation	UNP P01861
A	297	ASP	ASN	engineered mutation	UNP P01861
A	351	PHE	LEU	engineered mutation	UNP P01861
A	354	GLU	SER	engineered mutation	UNP P01861
A	366	ARG	THR	engineered mutation	UNP P01861
A	395	LYS	PRO	engineered mutation	UNP P01861
A	405	ARG	PHE	engineered mutation	UNP P01861
A	407	GLU	TYR	engineered mutation	UNP P01861


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	H	98	Total 98	O 98	0	0
2	A	56	Total 56	O 56	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Immunoglobulin heavy constant gamma 4



- Molecule 1: Immunoglobulin heavy constant gamma 4



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	102.97Å 110.92Å 87.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 2.30 35.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.0 (35.00-2.30) 99.1 (35.00-2.30)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.201 , 0.243 0.208 , 0.242	Depositor DCC
R_{free} test set	1047 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å ²)	39.4	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 21.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3496	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.23	8/1711 (0.5%)	1.42	5/2324 (0.2%)
1	H	1.20	5/1717 (0.3%)	1.36	6/2332 (0.3%)
All	All	1.21	13/3428 (0.4%)	1.39	11/4656 (0.2%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	387	PRO	C-O	-8.72	1.14	1.23
1	H	275	PHE	C-O	6.84	1.32	1.24
1	A	335	THR	C-O	6.44	1.31	1.23
1	H	368	LEU	C-O	6.24	1.31	1.24
1	A	310	HIS	CE1-NE2	6.03	1.38	1.32
1	H	284	VAL	C-O	5.77	1.29	1.23
1	H	313	TRP	C-O	5.76	1.30	1.24
1	H	435	HIS	CE1-NE2	5.21	1.37	1.32
1	A	285	HIS	CE1-NE2	5.14	1.37	1.32
1	A	381	TRP	C-N	-5.12	1.26	1.33
1	A	381	TRP	CA-C	-5.09	1.46	1.52
1	A	363	VAL	C-O	5.08	1.30	1.24
1	A	382	GLU	N-CA	-5.05	1.39	1.45

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	381	TRP	CA-C-N	-5.98	112.42	122.29
1	A	381	TRP	C-N-CA	-5.98	112.42	122.29
1	H	401	ASP	CA-CB-CG	5.85	118.45	112.60
1	A	416	ARG	N-CA-CB	-5.74	101.67	110.16
1	A	341	GLY	CA-C-O	-5.72	115.56	121.91
1	A	382	GLU	CA-C-O	-5.56	114.61	121.06
1	H	307	THR	CA-CB-OG1	-5.06	102.01	109.60
1	H	344	ARG	CA-C-N	5.05	127.43	120.67
1	H	344	ARG	C-N-CA	5.05	127.43	120.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	323	VAL	CA-C-N	5.03	130.88	122.73
1	H	323	VAL	C-N-CA	5.03	130.88	122.73

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1668	0	1620	21	0
1	H	1674	0	1625	23	0
2	A	56	0	0	1	0
2	H	98	0	0	1	1
All	All	3496	0	3245	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:ASN:HD21	1:A:421:ASN:ND2	1.76	0.83
1:A:386:GLN:HB3	2:A:514:HOH:O	1.87	0.72
1:A:384:ASN:HD21	1:A:421:ASN:HD22	1.36	0.72
1:H:370:LYS:HE3	1:H:405:ARG:HH21	1.62	0.63
1:H:358:MET:HE3	1:H:363:VAL:HG11	1.82	0.60
1:H:384:ASN:HD21	1:H:421:ASN:ND2	2.01	0.58
1:H:370:LYS:HE3	1:H:405:ARG:NH2	2.19	0.57
1:H:384:ASN:HD21	1:H:421:ASN:HD22	1.53	0.57
1:H:429:HIS:CD2	1:H:431:ALA:H	2.24	0.55
1:A:249:ASP:OD1	1:A:255:ARG:NH1	2.40	0.55
1:H:332:ILE:HD11	2:H:587:HOH:O	2.07	0.55
1:H:273:VAL:HG13	1:H:323:VAL:HG13	1.90	0.54
1:A:429:HIS:CD2	1:A:431:ALA:H	2.26	0.53
1:H:429:HIS:HD2	1:H:431:ALA:H	1.55	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:406:LEU:C	1:H:406:LEU:HD12	2.35	0.52
1:A:438:GLN:NE2	1:A:440:SER:OG	2.42	0.51
1:H:249:ASP:OD1	1:H:255:ARG:NH1	2.40	0.51
1:A:429:HIS:HD2	1:A:431:ALA:H	1.59	0.51
1:A:322:LYS:HG3	1:A:333:GLU:HB3	1.92	0.51
1:A:328:LEU:HD22	1:A:332:ILE:HD11	1.93	0.50
1:H:344:ARG:CZ	1:H:403:SER:HB3	2.42	0.48
1:H:253:ILE:HA	1:H:310:HIS:CE1	2.49	0.47
1:A:328:LEU:HD22	1:A:332:ILE:CD1	2.44	0.47
1:A:425:CYS:O	1:A:438:GLN:HA	2.16	0.46
1:A:374:PRO:O	1:A:429:HIS:HE1	1.99	0.46
1:A:406:LEU:C	1:A:406:LEU:HD12	2.40	0.45
1:H:434:ASN:HD22	1:H:434:ASN:HA	1.57	0.45
1:A:442:SER:O	1:A:443:LEU:HD23	2.17	0.44
1:H:269:GLU:O	1:H:271:PRO:HD3	2.17	0.44
1:H:370:LYS:HD2	1:H:405:ARG:NH2	2.33	0.44
1:H:391:TYR:HA	1:H:409:ARG:O	2.18	0.43
1:H:263:VAL:O	1:H:301:ARG:HA	2.19	0.43
1:H:292:ARG:HA	1:H:302:VAL:HG22	1.99	0.43
1:H:438:GLN:NE2	1:H:440:SER:OG	2.49	0.43
1:H:370:LYS:CE	1:H:405:ARG:HH21	2.30	0.43
1:H:319:TYR:O	1:H:335:THR:HA	2.19	0.42
1:H:406:LEU:HD12	1:H:406:LEU:O	2.20	0.42
1:A:238:PRO:HB2	1:A:328:LEU:HD21	2.00	0.42
1:A:251:LEU:O	1:A:435:HIS:HD2	2.03	0.41
1:A:363:VAL:O	1:A:411:THR:HA	2.21	0.41
1:A:271:PRO:HB2	1:A:292:ARG:HD3	2.02	0.41
1:A:434:ASN:HD22	1:A:434:ASN:HA	1.60	0.41
1:A:358:MET:HG3	1:A:414:LYS:HE3	2.02	0.40
1:A:246:LYS:HA	1:A:247:PRO:HD3	1.98	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:596:HOH:O	2:H:597:HOH:O[6_555]	2.00	0.20

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	204/207 (99%)	201 (98%)	3 (2%)	0	100	100
1	H	205/207 (99%)	201 (98%)	4 (2%)	0	100	100
All	All	409/414 (99%)	402 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/195 (100%)	189 (97%)	5 (3%)	40	59
1	H	195/195 (100%)	188 (96%)	7 (4%)	31	47
All	All	389/390 (100%)	377 (97%)	12 (3%)	35	52

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	267	SER
1	H	273	VAL
1	H	322	LYS
1	H	344	ARG
1	H	358	MET
1	H	386	GLN
1	H	438	GLN
1	A	270	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	305	VAL
1	A	336	ILE
1	A	386	GLN
1	A	438	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	285	HIS
1	H	295	GLN
1	H	310	HIS
1	H	325	ASN
1	H	384	ASN
1	H	386	GLN
1	H	418	GLN
1	H	421	ASN
1	H	429	HIS
1	H	434	ASN
1	H	438	GLN
1	A	274	GLN
1	A	285	HIS
1	A	421	ASN
1	A	429	HIS
1	A	434	ASN
1	A	435	HIS
1	A	438	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	206/207 (99%)	0.25	14 (6%) 23 25	25, 47, 99, 173	0
1	H	207/207 (100%)	-0.24	3 (1%) 73 75	26, 38, 64, 113	0
All	All	413/414 (99%)	0.00	17 (4%) 41 43	25, 42, 89, 173	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	443	LEU	4.3
1	A	329	PRO	4.1
1	H	238	PRO	3.1
1	A	352	PRO	3.0
1	A	423	PHE	2.9
1	A	412	VAL	2.9
1	A	340	LYS	2.6
1	H	356	GLU	2.5
1	A	296	PHE	2.5
1	A	359	THR	2.3
1	A	238	PRO	2.3
1	A	272	GLU	2.2
1	A	326	LYS	2.2
1	A	360	LYS	2.1
1	A	327	GLY	2.1
1	A	357	GLU	2.0
1	H	329	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.