



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:09 PM UTC

PDB ID : 2WO8 / pdb_00002wo8
Title : MMP12 complex with a beta hydroxy carboxylic acid
Authors : Holmes, I.P.; Gaines, S.; Watson, S.P.; Lorthioir, O.; Walker, A.; Baddeley, S.J.; Herbert, S.; Egan, D.; Convery, M.A.; Singh, O.M.P.; Gross, J.W.; StreLOW, J.M.; Smith, R.H.; Amour, A.J.; Brown, D.; Martin, S.L.
Deposited on : 2009-07-22
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	164	4% 86% 14%
1	B	164	93% 6% .
1	C	164	3% 90% 9% ..
1	D	164	5% 90% 7% .

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5785 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MACROPHAGE METALLOELASTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	164	Total	C	N	O	S	0	2	1
			1288	823	223	236	6			
1	B	163	Total	C	N	O	S	0	0	0
			1281	819	222	236	4			
1	C	163	Total	C	N	O	S	0	0	0
			1277	816	221	236	4			
1	D	160	Total	C	N	O	S	0	0	0
			1255	803	218	231	3			


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		
2	D	2	Total	Zn	0	0
			2	2		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	B	2	Total	Ca	0	0
			2	2		
3	C	1	Total	Ca	0	0
			1	1		
3	D	2	Total	Ca	0	0
			2	2		

- Molecule 4 is (3S)-5-biphenyl-4-yl-3-hydroxypentanoic acid (CCD ID: 077) (formula: $C_{17}H_{18}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			20	17	3		
4	C	1	Total	C	O	0	0
			20	17	3		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0

- Molecule 6 is N²-[(2R)-2-[(1S)-1-[FORMYL(HYDROXY)AMINO]ETHYL]-5-PHENYL PENTANOYL]-N,3-DIMETHYL-L-VALINAMIDE (CCD ID: 023) (formula: C₂₁H₃₃N₃O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C N O 28 21 3 4	0	0
6	D	1	Total C N O 28 21 3 4	0	0

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	D	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	135	Total	O	0	0
			135	135		
8	B	166	Total	O	0	0
			166	166		
8	C	123	Total	O	0	0
			123	123		
8	D	113	Total	O	0	0
			113	113		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MACROPHAGE METALLOELASTASE

Chain A: 



- Molecule 1: MACROPHAGE METALLOELASTASE

Chain B: 




- Molecule 1: MACROPHAGE METALLOELASTASE

Chain C: 



- Molecule 1: MACROPHAGE METALLOELASTASE

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	46.04Å 65.26Å 67.81Å 66.47° 83.86° 71.37°	Depositor
Resolution (Å)	20.00 – 2.00 20.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.00) 87.6 (20.00-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 1.99Å)	Xtriage
Refinement program	REFMAC 5.4.0073	Depositor
R, R_{free}	0.184 , 0.246 (Not available) , 0.235	Depositor DCC
R_{free} test set	2279 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	20.0	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5785	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CA, ZN, 023, GOL, 077

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	1/1337 (0.1%)	0.94	0/1811
1	B	0.79	0/1320	1.02	3/1789 (0.2%)
1	C	0.76	0/1316	1.00	2/1785 (0.1%)
1	D	0.76	0/1294	1.03	1/1756 (0.1%)
All	All	0.77	1/5267 (0.0%)	1.00	6/7141 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	267	GLU	C-N	-7.21	1.23	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	208	GLY	N-CA-C	6.72	120.79	112.73
1	C	207	SER	N-CA-C	6.31	120.68	112.92
1	C	172	HIS	N-CA-C	5.97	117.78	107.93
1	B	146	PRO	N-CA-C	-5.91	108.54	114.68
1	B	205	THR	N-CA-C	-5.30	105.58	111.36
1	D	194	ASP	N-CA-C	5.13	117.33	110.35

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1288	0	1222	18	0
1	B	1281	0	1211	3	0
1	C	1277	0	1200	14	0
1	D	1255	0	1183	9	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
3	C	1	0	0	0	0
3	D	2	0	0	0	0
4	A	20	0	17	1	0
4	C	20	0	17	2	0
5	A	10	0	0	0	0
5	B	5	0	0	0	0
5	C	5	0	0	1	0
5	D	5	0	0	0	0
6	B	28	0	32	0	0
6	D	28	0	32	1	0
7	D	12	0	16	0	0
8	A	135	0	0	3	0
8	B	166	0	0	1	0
8	C	123	0	0	4	0
8	D	113	0	0	0	0
All	All	5785	0	4930	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125[B]:MET:HG3	1:A:130:VAL:HG23	1.34	1.06
1:D:264:ASP:HB2	1:D:265:PRO:HD2	1.51	0.91
1:A:125[B]:MET:HG3	1:A:130:VAL:CG2	2.13	0.76
1:A:125[B]:MET:CG	1:A:130:VAL:HG23	2.17	0.74
1:C:237:PHE:CD1	1:C:238:PRO:HD2	2.26	0.71
1:B:128:GLU:HG3	8:B:2022:HOH:O	1.91	0.69
1:C:145:THR:HB	1:C:146:PRO:HD2	1.80	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:ILE:HD12	8:A:2069:HOH:O	2.01	0.61
1:A:125[B]:MET:CG	1:A:130:VAL:CG2	2.75	0.60
1:A:156:MET:HA	1:A:156:MET:HE2	1.87	0.57
1:A:182:ALA:O	4:A:1271:077:H171	2.08	0.54
1:D:264:ASP:CB	1:D:265:PRO:HD2	2.31	0.54
1:A:264:ASP:HB2	1:A:265:PRO:CD	2.37	0.53
1:C:264:ASP:HB2	1:C:265:PRO:HD2	1.90	0.53
1:C:264:ASP:HB2	1:C:265:PRO:CD	2.39	0.53
1:C:154:THR:HG22	8:C:2015:HOH:O	2.09	0.53
1:C:208:GLY:N	5:C:1272:SO4:O4	2.32	0.52
1:A:145:THR:HB	1:A:146:PRO:CD	2.41	0.51
1:C:145:THR:HB	1:C:146:PRO:CD	2.41	0.50
1:C:171:PHE:CD1	1:C:172:HIS:HB2	2.49	0.48
1:A:264:ASP:HB2	1:A:265:PRO:HD2	1.96	0.48
1:A:248:PHE:O	1:A:249:ARG:HG3	2.14	0.48
1:A:113:TYR:CE1	1:A:150:SER:HB2	2.50	0.46
1:D:207:SER:HB3	1:D:242:TYR:CZ	2.51	0.46
1:A:243:VAL:O	1:A:244:ASP:C	2.59	0.46
1:C:178:GLY:HA2	8:C:2091:HOH:O	2.16	0.45
1:D:264:ASP:HB2	1:D:265:PRO:CD	2.35	0.45
1:A:182:ALA:HB1	1:A:196:HIS:O	2.17	0.45
1:C:245:ILE:C	1:C:245:ILE:HD12	2.43	0.44
1:D:108:VAL:HA	1:D:187:PRO:HG2	1.99	0.43
1:C:116:TYR:HA	1:C:159:ILE:O	2.17	0.43
1:C:154:THR:HG21	8:C:2054:HOH:O	2.18	0.43
1:A:108:VAL:HA	1:A:187:PRO:HG2	2.00	0.42
1:A:127:ARG:NH2	8:A:2024:HOH:O	2.52	0.42
1:A:187:PRO:HA	8:A:2080:HOH:O	2.19	0.42
1:D:111:LYS:H	1:D:111:LYS:HG2	1.37	0.42
1:D:108:VAL:CG1	1:D:261:LEU:HD22	2.50	0.41
1:C:171:PHE:CE1	1:C:172:HIS:HB2	2.55	0.41
1:C:215:THR:HG23	4:C:1271:077:H11	2.03	0.41
1:A:203:TRP:CZ3	1:A:210:THR:HG22	2.56	0.41
1:B:172:HIS:O	1:B:183:HIS:HE1	2.03	0.41
1:B:215:THR:O	1:B:219:GLU:HG2	2.20	0.41
4:C:1271:077:H1	8:C:2104:HOH:O	2.21	0.41
1:D:182:ALA:O	6:D:1266:023:H11	2.21	0.40
1:D:183:HIS:CD2	1:D:183:HIS:C	2.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/164 (99%)	156 (96%)	7 (4%)	0	100	100
1	B	161/164 (98%)	156 (97%)	5 (3%)	0	100	100
1	C	161/164 (98%)	157 (98%)	4 (2%)	0	100	100
1	D	158/164 (96%)	150 (95%)	8 (5%)	0	100	100
All	All	643/656 (98%)	619 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/133 (101%)	133 (99%)	1 (1%)	76	82
1	B	132/133 (99%)	130 (98%)	2 (2%)	57	64
1	C	131/133 (98%)	131 (100%)	0	100	100
1	D	129/133 (97%)	128 (99%)	1 (1%)	73	80
All	All	526/532 (99%)	522 (99%)	4 (1%)	73	80

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	165	ARG
1	B	233	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	241	LYS
1	D	241	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	172	HIS
1	B	172	HIS
1	B	259	GLN
1	C	246	ASN
1	D	112	HIS
1	D	259	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 25 ligands modelled in this entry, 14 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	GOL	D	1271	-	5,5,5	0.36	0	5,5,5	0.55	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	077	A	1271	2	21,21,21	1.02	2 (9%)	26,27,27	1.55	7 (26%)
5	SO4	C	1272	-	4,4,4	0.25	0	6,6,6	0.16	0
5	SO4	D	1273	-	4,4,4	0.22	0	6,6,6	0.55	0
5	SO4	A	1272	-	4,4,4	0.17	0	6,6,6	0.31	0
6	023	B	1268	2	27,28,28	0.80	0	30,38,38	1.68	6 (20%)
6	023	D	1266	2	27,28,28	0.85	0	30,38,38	1.45	5 (16%)
7	GOL	D	1272	-	5,5,5	0.49	0	5,5,5	0.72	0
5	SO4	B	1273	-	4,4,4	0.23	0	6,6,6	0.44	0
4	077	C	1271	2	21,21,21	0.95	0	26,27,27	1.35	2 (7%)
5	SO4	A	1273	-	4,4,4	0.24	0	6,6,6	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	D	1271	-	-	1/4/4/4	-
4	077	A	1271	2	-	6/13/13/13	0/2/2/2
7	GOL	D	1272	-	-	1/4/4/4	-
6	023	B	1268	2	-	0/33/36/36	0/1/1/1
4	077	C	1271	2	-	3/13/13/13	0/2/2/2
6	023	D	1266	2	-	0/33/36/36	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1271	077	O4-C18	-2.44	1.22	1.30
4	A	1271	077	C1-C3	2.05	1.42	1.38

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1268	023	O16-C11-N7	-4.71	119.27	124.50
4	C	1271	077	C16-C12-C20	-3.98	108.41	114.95
6	D	1266	023	O12-N7-C11	3.81	121.68	115.17
4	A	1271	077	C16-C12-C20	-3.33	109.48	114.95
6	B	1268	023	C25-N18-C14	-3.14	116.74	122.21
6	D	1266	023	C10-N5-C2	-3.02	115.87	121.98
6	D	1266	023	C25-N18-C14	-2.90	117.16	122.21
6	D	1266	023	C14-C10-N5	2.63	115.57	109.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1268	023	C9-C4-C1	2.63	118.92	114.28
4	A	1271	077	C5-C2-C6	2.55	123.44	120.54
6	B	1268	023	O19-C14-C10	2.42	124.82	120.85
4	A	1271	077	C5-C4-C1	-2.38	116.61	119.87
4	C	1271	077	O3-C18-C17	-2.36	115.59	122.84
6	D	1266	023	O16-C11-N7	-2.29	121.95	124.50
4	A	1271	077	O1-C20-C12	-2.24	103.37	109.35
4	A	1271	077	C4-C1-C3	2.20	122.96	120.24
6	B	1268	023	C27-C24-C17	-2.19	117.54	120.61
6	B	1268	023	C28-C26-C23	-2.15	117.58	120.24
4	A	1271	077	C3-C6-C8	2.12	124.91	121.25
4	A	1271	077	C14-C8-C6	-2.02	117.77	121.25

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1271	077	C18-C17-C20-O1
4	A	1271	077	C18-C17-C20-C12
7	D	1272	GOL	C1-C2-C3-O3
4	A	1271	077	C16-C12-C20-C17
4	A	1271	077	C16-C12-C20-O1
7	D	1271	GOL	C1-C2-C3-O3
4	A	1271	077	C60-C13-C16-C12
4	A	1271	077	C11-C13-C16-C12
4	C	1271	077	C16-C12-C20-C17
4	C	1271	077	C60-C13-C16-C12
4	C	1271	077	C11-C13-C16-C12


There are no ring outliers.

4 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1271	077	1	0
5	C	1272	SO4	1	0
6	D	1266	023	1	0
4	C	1271	077	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will


also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand 023 B 1268




Bond lengths



Bond angles




Torsions




Rings

Ligand 023 D 1266



Ligand 077 C 1271



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	164/164 (100%)	0.25	6 (3%) 45 44	19, 29, 45, 55	2 (1%)
1	B	163/164 (99%)	-0.20	0 100 100	15, 22, 35, 51	0
1	C	163/164 (99%)	0.16	5 (3%) 51 50	18, 27, 45, 53	0
1	D	160/164 (97%)	0.49	9 (5%) 30 29	15, 31, 50, 63	0
All	All	650/656 (99%)	0.17	20 (3%) 51 50	15, 27, 46, 63	2 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	154	THR	3.5
1	C	245	ILE	3.1
1	A	113	TYR	2.6
1	D	155	GLY	2.5
1	C	156	MET	2.5
1	C	172	HIS	2.5
1	A	191	ILE	2.5
1	C	113	TYR	2.4
1	D	110	ARG	2.4
1	C	171	PHE	2.4
1	D	191	ILE	2.3
1	D	189	SER	2.3
1	A	190	GLY	2.2
1	D	156	MET	2.2
1	D	263	GLY	2.2
1	A	268	ASN	2.1
1	A	246	ASN	2.1
1	D	147	LEU	2.0
1	D	114	ILE	2.0
1	A	171	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	A	1273	5/5	0.77	0.12	71,72,73,74	0
5	SO4	C	1272	5/5	0.83	0.15	77,78,79,81	0
5	SO4	B	1273	5/5	0.88	0.11	58,60,61,62	0
4	077	A	1271	20/20	0.91	0.10	29,31,34,36	0
4	077	C	1271	20/20	0.92	0.10	24,31,37,39	0
7	GOL	D	1272	6/6	0.93	0.09	23,29,30,31	0
6	023	D	1266	28/28	0.94	0.07	9,19,22,23	0
7	GOL	D	1271	6/6	0.95	0.07	25,25,28,30	0
6	023	B	1268	28/28	0.95	0.07	14,19,21,23	0
5	SO4	D	1273	5/5	0.97	0.07	26,28,31,31	0
5	SO4	A	1272	5/5	0.97	0.06	25,25,28,29	0
3	CA	D	1270	1/1	0.97	0.07	29,29,29,29	0
3	CA	B	1272	1/1	0.97	0.07	27,27,27,27	0
3	CA	C	1270	1/1	0.97	0.04	28,28,28,28	0
3	CA	A	1270	1/1	0.98	0.03	28,28,28,28	0
3	CA	B	1271	1/1	0.99	0.02	21,21,21,21	0
2	ZN	A	1269	1/1	0.99	0.02	29,29,29,29	0
2	ZN	C	1268	1/1	0.99	0.03	26,26,26,26	0
3	CA	D	1269	1/1	0.99	0.02	21,21,21,21	0
2	ZN	C	1269	1/1	0.99	0.02	27,27,27,27	0
2	ZN	D	1267	1/1	0.99	0.02	23,23,23,23	0
2	ZN	D	1268	1/1	0.99	0.02	27,27,27,27	0
2	ZN	A	1268	1/1	0.99	0.02	28,28,28,28	0
2	ZN	B	1270	1/1	1.00	0.02	21,21,21,21	0
2	ZN	B	1269	1/1	1.00	0.01	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 077 A 1271:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around 077 C 1271:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 023 D 1266:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 023 B 1268:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.