



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 11:46 AM EDT

PDB ID : 2WO9 / pdb_00002wo9
Title : MMP12 complex with a beta hydroxy carboxylic acid
Authors : Holmes, I.P.; Gaines, S.; Watson, S.P.; Lorthioir, O.; Walker, A.; Baddeley, S.J.; Herbert, S.; Egan, D.; Convery, M.A.; Singh, O.M.P.; Gross, J.W.; StreLOW, J.M.; Smith, R.H.; Amour, A.J.; Brown, D.; Martin, S.L.
Deposited on : 2009-07-22
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	164	2% 90% 9%
1	B	164	0% 95% 5%
1	C	164	4% 95% 5%
1	D	164	4% 93% 3%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5940 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MACROPHAGE METALLOELASTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	2	0
			1280	817	222	238	3			
1	B	162	Total	C	N	O	S	0	2	0
			1276	817	221	234	4			
1	C	164	Total	C	N	O	S	0	0	0
			1285	820	223	238	4			
1	D	160	Total	C	N	O	S	0	2	0
			1257	803	218	233	3			


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		
2	D	2	Total	Zn	0	0
			2	2		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	B	3	Total	Ca	0	0
			3	3		
3	C	1	Total	Ca	0	0
			1	1		
3	D	2	Total	Ca	0	0
			2	2		

- Molecule 4 is (3S)-5-(4'-ACETYLBIIPHENYL-4-YL)-3-HYDROXPENTANOIC ACID (CCD ID: 068) (formula: C₁₉H₂₀O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			23	19	4		
4	C	1	Total	C	O	0	0
			23	19	4		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0

- Molecule 6 is N²-[(2R)-2-[(1S)-1-[FORMYL(HYDROXY)AMINO]ETHYL]-5-PHENYL PENTANOYL]-N,3-DIMETHYL-L-VALINAMIDE (CCD ID: 023) (formula: C₂₁H₃₃N₃O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C N O 28 21 3 4	0	0
6	D	1	Total C N O 28 21 3 4	0	0

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	165	Total	O	0	0
			165	165		
8	B	191	Total	O	0	0
			191	191		
8	C	160	Total	O	0	0
			160	160		
8	D	146	Total	O	0	0
			146	146		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MACROPHAGE METALLOELASTASE




- Molecule 1: MACROPHAGE METALLOELASTASE



- Molecule 1: MACROPHAGE METALLOELASTASE



- Molecule 1: MACROPHAGE METALLOELASTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	46.35Å 65.29Å 69.36Å 66.15° 83.84° 71.38°	Depositor
Resolution (Å)	19.88 – 1.70 19.88 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.88-1.70) 80.2 (19.88-1.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 1.69Å)	Xtriage
Refinement program	REFMAC 5.4.0073	Depositor
R, R_{free}	0.172 , 0.218 0.174 , 0.220	Depositor DCC
R_{free} test set	3153 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	17.0	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5940	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA, 068, 023, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/1324	0.90	2/1795 (0.1%)
1	B	0.89	1/1325 (0.1%)	0.92	2/1796 (0.1%)
1	C	0.79	0/1324	0.88	0/1796
1	D	0.78	0/1306	0.93	3/1772 (0.2%)
All	All	0.81	1/5279 (0.0%)	0.91	7/7159 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	227	GLY	N-CA	5.58	1.50	1.44

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	237	PHE	N-CA-C	-6.60	101.45	109.83
1	D	169	GLY	N-CA-C	6.30	124.06	115.30
1	B	208	GLY	N-CA-C	6.11	121.25	113.24
1	B	237	PHE	N-CA-C	-6.01	102.01	109.64
1	A	207[A]	SER	N-CA-C	5.93	120.09	112.26
1	A	207[B]	SER	N-CA-C	5.93	120.09	112.26
1	D	159	ILE	N-CA-C	5.14	115.26	107.75

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1280	0	1207	16	0
1	B	1276	0	1213	2	0
1	C	1285	0	1206	5	0
1	D	1257	0	1185	4	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	1	0	0	0	0
3	B	3	0	0	0	0
3	C	1	0	0	0	0
3	D	2	0	0	0	0
4	A	23	0	19	1	0
4	C	23	0	19	1	0
5	A	15	0	0	0	0
5	B	15	0	0	0	0
5	C	5	0	0	0	0
5	D	10	0	0	0	0
6	B	28	0	32	0	0
6	D	28	0	32	0	0
7	C	6	0	8	0	0
7	D	12	0	16	1	0
8	A	165	0	0	4	1
8	B	191	0	0	0	3
8	C	160	0	0	1	0
8	D	146	0	0	0	4
All	All	5940	0	4937	26	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:264:ASP:HB2	1:D:265:PRO:HD2	1.78	0.65
1:A:247:THR:CG2	1:A:247:THR:O	2.55	0.55
1:A:247:THR:O	1:A:247:THR:HG22	2.07	0.55
1:D:264:ASP:HB2	1:D:265:PRO:CD	2.38	0.53
1:A:177:LYS:HG2	1:B:126:ASN:HD22	1.74	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:GLU:O	1:A:199:GLU:HG3	2.08	0.52
1:C:145:THR:HB	1:C:146:PRO:HD2	1.93	0.51
1:A:154:THR:HG23	8:A:2081:HOH:O	2.11	0.50
1:C:267:GLU:OE2	8:C:2151:HOH:O	2.20	0.50
1:A:199:GLU:OE2	8:A:2116:HOH:O	2.20	0.47
1:A:145:THR:HB	1:A:146:PRO:HD2	1.98	0.46
1:A:106:GLY:N	8:A:2002:HOH:O	2.51	0.44
1:B:145:THR:HB	1:B:146:PRO:HD2	2.00	0.43
7:D:1271:GOL:H2	7:D:1272:GOL:H2	2.00	0.43
1:A:264:ASP:HB2	1:A:265:PRO:HD2	2.01	0.43
1:A:245:ILE:N	1:A:245:ILE:HD13	2.33	0.42
1:C:244:ASP:HB3	1:C:247:THR:OG1	2.19	0.42
1:A:182:ALA:O	4:A:1271:068:H171	2.19	0.42
1:A:248:PHE:CD2	1:A:248:PHE:C	2.96	0.42
1:A:264:ASP:HB2	1:A:265:PRO:CD	2.50	0.41
1:C:264:ASP:HB2	1:C:265:PRO:CD	2.50	0.41
1:C:140:VAL:HG13	1:C:250:LEU:HG	2.03	0.41
1:A:248:PHE:HA	8:A:2146:HOH:O	2.19	0.40
1:A:169:GLY:CA	1:D:106:GLY:HA2	2.52	0.40
1:A:169:GLY:HA2	1:D:106:GLY:HA2	2.02	0.40
4:C:1272:068:H191	4:C:1272:068:H13	1.77	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:B:2162:HOH:O	8:D:2032:HOH:O[1_454]	1.87	0.33
8:A:2147:HOH:O	8:D:2134:HOH:O[1_564]	2.00	0.20
8:B:2111:HOH:O	8:D:2032:HOH:O[1_454]	2.09	0.11
8:B:2177:HOH:O	8:D:2013:HOH:O[1_454]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	162/164 (99%)	159 (98%)	3 (2%)	0	100	100
1	B	162/164 (99%)	158 (98%)	4 (2%)	0	100	100
1	C	162/164 (99%)	158 (98%)	4 (2%)	0	100	100
1	D	160/164 (98%)	156 (98%)	4 (2%)	0	100	100
All	All	646/656 (98%)	631 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/133 (100%)	133 (100%)	0	100	100
1	B	133/133 (100%)	132 (99%)	1 (1%)	73	65
1	C	132/133 (99%)	132 (100%)	0	100	100
1	D	131/133 (98%)	130 (99%)	1 (1%)	73	65
All	All	529/532 (99%)	527 (100%)	2 (0%)	84	80

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	241	LYS
1	D	241	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	112	HIS
1	A	206	HIS
1	B	119	ASN
1	B	126	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	259	GLN
1	C	172	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 31 ligands modelled in this entry, 15 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	A	1272	-	4,4,4	0.27	0	6,6,6	0.28	0
5	SO4	A	1274	-	4,4,4	0.25	0	6,6,6	0.14	0
5	SO4	B	1275	-	4,4,4	0.23	0	6,6,6	0.11	0
7	GOL	D	1271	-	5,5,5	0.35	0	5,5,5	0.56	0
5	SO4	B	1273	-	4,4,4	0.24	0	6,6,6	0.20	0
6	023	D	1266	2	27,28,28	0.96	0	30,38,38	1.31	1 (3%)
5	SO4	D	1274	-	4,4,4	0.25	0	6,6,6	0.19	0
5	SO4	B	1274	-	4,4,4	0.25	0	6,6,6	0.20	0
5	SO4	C	1274	-	4,4,4	0.39	0	6,6,6	0.41	0
5	SO4	D	1273	-	4,4,4	0.20	0	6,6,6	0.17	0
6	023	B	1267	2	27,28,28	0.99	1 (3%)	30,38,38	1.28	2 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	C	1273	-	5,5,5	0.30	0	5,5,5	0.57	0
7	GOL	D	1272	-	5,5,5	0.58	0	5,5,5	0.61	0
4	068	C	1272	2	24,24,24	0.83	0	31,32,32	0.87	0
5	SO4	A	1273	-	4,4,4	0.21	0	6,6,6	0.13	0
4	068	A	1271	2	24,24,24	0.87	1 (4%)	31,32,32	0.91	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	D	1271	-	-	0/4/4/4	-
6	023	D	1266	2	-	0/33/36/36	0/1/1/1
6	023	B	1267	2	-	0/33/36/36	0/1/1/1
7	GOL	D	1272	-	-	0/4/4/4	-
4	068	C	1272	2	-	4/17/17/17	0/2/2/2
7	GOL	C	1273	-	-	4/4/4/4	-
4	068	A	1271	2	-	4/17/17/17	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1271	068	O19-C18	-2.61	1.22	1.30
6	B	1267	023	C24-C17	2.30	1.43	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1267	023	O16-C11-N7	-4.41	119.60	124.50
6	D	1266	023	O16-C11-N7	-4.26	119.76	124.50
6	B	1267	023	C10-C14-N18	-2.32	112.47	116.31
4	A	1271	068	C4-C3-C2	-2.04	111.60	114.95

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	1272	068	C18-C17-C2-C3
7	C	1273	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	C	1273	GOL	O1-C1-C2-O2
7	C	1273	GOL	C1-C2-C3-O3
7	C	1273	GOL	O2-C2-C3-O3
4	C	1272	068	C3-C4-C5-C6
4	A	1271	068	C3-C4-C5-C6
4	A	1271	068	C3-C4-C5-C10
4	C	1272	068	C3-C4-C5-C10
4	A	1271	068	C18-C17-C2-C3
4	C	1272	068	C2-C17-C18-O19
4	A	1271	068	C2-C17-C18-O19


There are no ring outliers.

4 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	D	1271	GOL	1	0
7	D	1272	GOL	1	0
4	C	1272	068	1	0
4	A	1271	068	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand 023 D 1266




Bond lengths



Bond angles




Torsions





Rings

Ligand 023 B 1267



Ligand 068 C 1272





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/164 (98%)	0.29	3 (1%) 66 71	13, 25, 40, 48	2 (1%)
1	B	162/164 (98%)	-0.12	1 (0%) 85 88	13, 22, 33, 49	2 (1%)
1	C	164/164 (100%)	0.25	7 (4%) 40 44	18, 26, 41, 53	0
1	D	160/164 (97%)	0.23	7 (4%) 39 42	18, 25, 40, 48	2 (1%)
All	All	648/656 (98%)	0.16	18 (2%) 55 59	13, 24, 40, 53	6 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	268	ASN	4.1
1	C	208	GLY	3.1
1	D	110	ARG	2.8
1	A	113	TYR	2.7
1	A	171	PHE	2.7
1	B	110	ARG	2.6
1	C	244	ASP	2.5
1	C	249	ARG	2.5
1	C	156	MET	2.4
1	A	156	MET	2.4
1	D	156	MET	2.4
1	D	263	GLY	2.4
1	D	154	THR	2.4
1	C	113	TYR	2.4
1	D	191	ILE	2.3
1	D	264	ASP	2.2
1	C	171	PHE	2.2
1	D	208	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	A	1273	5/5	0.78	0.13	58,60,61,62	0
7	GOL	C	1273	6/6	0.78	0.16	33,40,42,47	0
5	SO4	A	1274	5/5	0.80	0.12	50,50,51,51	0
5	SO4	A	1272	5/5	0.84	0.11	41,42,46,47	0
5	SO4	D	1274	5/5	0.88	0.10	50,50,55,55	0
5	SO4	B	1275	5/5	0.89	0.10	49,49,52,54	0
5	SO4	D	1273	5/5	0.92	0.07	50,51,52,53	0
4	068	A	1271	23/23	0.93	0.10	21,24,32,34	0
5	SO4	B	1274	5/5	0.93	0.08	52,53,55,56	0
4	068	C	1272	23/23	0.93	0.10	20,25,34,35	0
6	023	D	1266	28/28	0.96	0.07	14,19,22,25	0
6	023	B	1267	28/28	0.96	0.06	15,18,22,25	0
7	GOL	D	1271	6/6	0.96	0.09	22,24,27,28	0
3	CA	A	1270	1/1	0.98	0.05	23,23,23,23	0
5	SO4	C	1274	5/5	0.98	0.06	26,27,29,31	0
5	SO4	B	1273	5/5	0.98	0.06	23,23,25,27	0
3	CA	D	1270	1/1	0.98	0.08	26,26,26,26	0
7	GOL	D	1272	6/6	0.98	0.06	22,26,30,31	0
2	ZN	C	1269	1/1	0.99	0.04	25,25,25,25	0
2	ZN	D	1268	1/1	0.99	0.04	25,25,25,25	0
2	ZN	A	1268	1/1	0.99	0.04	26,26,26,26	0
3	CA	B	1270	1/1	0.99	0.03	21,21,21,21	0
3	CA	B	1271	1/1	0.99	0.14	28,28,28,28	0
3	CA	B	1272	1/1	0.99	0.09	27,27,27,27	0
3	CA	C	1271	1/1	0.99	0.03	23,23,23,23	0
2	ZN	A	1269	1/1	0.99	0.04	29,29,29,29	0
2	ZN	B	1269	1/1	1.00	0.04	22,22,22,22	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	B	1268	1/1	1.00	0.04	21,21,21,21	0
3	CA	D	1269	1/1	1.00	0.01	20,20,20,20	0
2	ZN	C	1270	1/1	1.00	0.03	24,24,24,24	0
2	ZN	D	1267	1/1	1.00	0.03	24,24,24,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 068 A 1271:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 068 C 1272:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 023 D 1266:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 023 B 1267:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.