



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 01:51 PM UTC

PDB ID : 3WO4 / pdb_00003wo4
Title : Crystal structure of the IL-18 signaling ternary complex
Authors : Tsutsumi, N.; Kimura, T.; Arita, K.; Ariyoshi, M.; Ohnishi, H.; Kondo, N.; Shirakawa, M.; Kato, Z.; Tochio, H.
Deposited on : 2013-12-19
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	157	 68% 31% .
2	B	312	 72% 21% . .
3	C	344	 3% 60% 16% . 22%
4	D	3	 67% 33%
5	E	5	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	3	 100%
7	G	2	 50% 50%
7	J	2	 100%
8	H	4	 75% 25%
9	I	4	 50% 50%

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 6030 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	157	Total	C	N	O	S	0	0	0
			1266	796	209	251	10			

- Molecule 2 is a protein called Interleukin-18 receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	301	Total	C	N	O	S	0	1	0
			2375	1499	406	456	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP Q13478
B	-1	PRO	-	expression tag	UNP Q13478


- Molecule 3 is a protein called Interleukin-18 receptor accessory protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	270	Total	C	N	O	S	0	0	0
			2022	1263	347	402	10			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP O95256
C	-1	PRO	-	expression tag	UNP O95256

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	F	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	G	2	Total	C	N	O	0	0	0
			28	16	2	10			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	J	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 8 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	H	4	Total	C	N	O	0	0	0
			49	28	2	19			

- Molecule 9 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
9	I	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	B	1	Total	C	N	O	0	0
			14	8	1	5		
10	B	1	Total	C	N	O	0	0
			14	8	1	5		
10	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 11 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	C	1	Total	Cl	0	0
			1	1		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	14	Total	O	0	0
			14	14		
12	B	7	Total	O	0	0
			7	7		
12	C	10	Total	O	0	0
			10	10		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-18



- Molecule 2: Interleukin-18 receptor 1



- Molecule 3: Interleukin-18 receptor accessory protein





- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 67% 33%



- Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 100%



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 100%



- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 50% 50%



- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J: 100%



- Molecule 8: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 75% 25%



- Molecule 9: α -D-mannopyranose-(1-3)- β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain I:



MAG1
MAG2
BMA3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.56Å 111.56Å 134.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.00 – 3.10 42.00 – 3.10	Depositor EDS
% Data completeness (in resolution range)	85.0 (42.00-3.10) 84.9 (42.00-3.10)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.39 (at 3.07Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.188 , 0.232 0.198 , 0.246	Depositor DCC
R_{free} test set	884 reflections (4.29%)	wwPDB-VP
Wilson B-factor (Å ²)	58.9	Xtriage
Anisotropy	0.580	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 100.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6030	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CL, FUC, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	0/1287	1.21	5/1726 (0.3%)
2	B	0.81	0/2433	1.23	6/3310 (0.2%)
3	C	0.83	0/2057	1.25	8/2800 (0.3%)
All	All	0.83	0/5777	1.24	19/7836 (0.2%)

There are no bond length outliers.

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	132	ASP	CA-CB-CG	7.84	120.44	112.60
2	B	164	GLU	CA-C-N	6.15	131.10	122.36
2	B	164	GLU	C-N-CA	6.15	131.10	122.36
3	C	341	ASN	CA-CB-CG	5.90	118.50	112.60
3	C	217	GLY	N-CA-C	5.77	118.41	110.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1266	0	1237	14	0
2	B	2375	0	2231	18	0
3	C	2022	0	1857	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	38	0	34	0	0
5	E	61	0	52	0	0
6	F	39	0	34	0	0
7	G	28	0	25	0	0
7	J	28	0	25	0	0
8	H	49	0	43	0	0
9	I	50	0	43	0	0
10	B	28	0	26	0	0
10	C	14	0	13	0	0
11	C	1	0	0	0	0
12	A	14	0	0	1	0
12	B	7	0	0	0	0
12	C	10	0	0	0	0
All	All	6030	0	5620	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:295:LEU:HD12	2:B:312:ILE:HD11	1.64	0.80
2:B:180:GLN:HG3	2:B:205:THR:HA	1.77	0.67
2:B:301:ALA:HA	2:B:306:THR:HG22	1.79	0.64
3:C:216:GLN:HG3	3:C:241:ARG:HA	1.84	0.59
2:B:54:LYS:HB3	2:B:98:PHE:HE1	1.68	0.58

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	155/157 (99%)	142 (92%)	12 (8%)	1 (1%)	21	52
2	B	300/312 (96%)	278 (93%)	18 (6%)	4 (1%)	9	35
3	C	264/344 (77%)	237 (90%)	25 (10%)	2 (1%)	16	47
All	All	719/813 (88%)	657 (91%)	55 (8%)	7 (1%)	12	41

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	116	PRO
2	B	72	SER
2	B	80	ASP
2	B	166	ASN
2	B	209	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/148 (98%)	118 (81%)	27 (19%)	1	8
2	B	263/288 (91%)	213 (81%)	50 (19%)	1	7
3	C	212/321 (66%)	179 (84%)	33 (16%)	2	12
All	All	620/757 (82%)	510 (82%)	110 (18%)	2	9

5 of 110 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	217	LEU
2	B	292	LEU
3	C	356	ARG
3	C	300	VAL
2	B	222	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	212	ASN
2	B	255	ASN
3	C	320	ASN
3	C	204	ASN
3	C	239	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

23 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	D	1	2,4	14,14,15	0.31	0	17,19,21	0.55	0
4	NAG	D	2	4	14,14,15	0.29	0	17,19,21	0.56	0
4	FUC	D	3	4	10,10,11	0.44	0	14,14,16	0.79	1 (7%)
5	NAG	E	1	5,2	14,14,15	0.28	0	17,19,21	0.73	1 (5%)
5	NAG	E	2	5	14,14,15	0.35	0	17,19,21	1.64	4 (23%)
5	BMA	E	3	5	11,11,12	0.33	0	15,15,17	0.86	1 (6%)
5	MAN	E	4	5	11,11,12	0.51	0	15,15,17	1.48	2 (13%)
5	MAN	E	5	5	11,11,12	0.39	0	15,15,17	1.01	1 (6%)
6	NAG	F	1	2,6	14,14,15	0.30	0	17,19,21	1.31	1 (5%)
6	NAG	F	2	6	14,14,15	0.38	0	17,19,21	1.68	4 (23%)
6	BMA	F	3	6	11,11,12	0.37	0	15,15,17	0.83	1 (6%)
7	NAG	G	1	2,7	14,14,15	0.31	0	17,19,21	1.00	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	NAG	G	2	7	14,14,15	0.34	0	17,19,21	0.38	0
8	NAG	H	1	2,8	14,14,15	0.30	0	17,19,21	0.75	0
8	NAG	H	2	8	14,14,15	0.32	0	17,19,21	0.51	0
8	BMA	H	3	8	11,11,12	0.34	0	15,15,17	0.60	0
8	FUC	H	4	8	10,10,11	0.55	0	14,14,16	0.95	2 (14%)
9	NAG	I	1	3,9	14,14,15	0.33	0	17,19,21	0.72	1 (5%)
9	NAG	I	2	9	14,14,15	0.32	0	17,19,21	0.73	0
9	BMA	I	3	9	11,11,12	0.30	0	15,15,17	0.49	0
9	MAN	I	4	9	11,11,12	0.40	0	15,15,17	0.79	1 (6%)
7	NAG	J	1	7,3	14,14,15	0.31	0	17,19,21	0.44	0
7	NAG	J	2	7	14,14,15	0.29	0	17,19,21	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	FUC	D	3	4	-	-	0/1/1/1
5	NAG	E	1	5,2	-	0/6/23/26	0/1/1/1
5	NAG	E	2	5	-	2/6/23/26	0/1/1/1
5	BMA	E	3	5	-	0/2/19/22	0/1/1/1
5	MAN	E	4	5	-	0/2/19/22	0/1/1/1
5	MAN	E	5	5	-	1/2/19/22	0/1/1/1
6	NAG	F	1	2,6	-	0/6/23/26	0/1/1/1
6	NAG	F	2	6	-	1/6/23/26	0/1/1/1
6	BMA	F	3	6	-	1/2/19/22	0/1/1/1
7	NAG	G	1	2,7	-	0/6/23/26	0/1/1/1
7	NAG	G	2	7	-	2/6/23/26	0/1/1/1
8	NAG	H	1	2,8	-	0/6/23/26	0/1/1/1
8	NAG	H	2	8	-	1/6/23/26	0/1/1/1
8	BMA	H	3	8	-	0/2/19/22	0/1/1/1
8	FUC	H	4	8	-	-	0/1/1/1
9	NAG	I	1	3,9	-	0/6/23/26	0/1/1/1
9	NAG	I	2	9	-	0/6/23/26	0/1/1/1
9	BMA	I	3	9	-	0/2/19/22	0/1/1/1
9	MAN	I	4	9	-	0/2/19/22	0/1/1/1
7	NAG	J	1	7,3	-	0/6/23/26	0/1/1/1
7	NAG	J	2	7	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	4	MAN	C1-O5-C5	4.92	118.78	112.19
6	F	2	NAG	C1-O5-C5	4.74	118.54	112.19
6	F	1	NAG	O5-C1-C2	-4.58	104.20	111.29
5	E	2	NAG	C1-O5-C5	4.54	118.27	112.19
5	E	5	MAN	C1-O5-C5	3.55	116.94	112.19

There are no chirality outliers.


5 of 8 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
7	G	2	NAG	O5-C5-C6-O6
6	F	3	BMA	O5-C5-C6-O6
5	E	5	MAN	O5-C5-C6-O6
6	F	2	NAG	O5-C5-C6-O6
7	G	2	NAG	C4-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
10	NAG	B	901	2	14,14,15	0.30	0	17,19,21	0.63	1 (5%)
10	NAG	C	901	3	14,14,15	0.30	0	17,19,21	0.53	0
10	NAG	B	902	2	14,14,15	0.32	0	17,19,21	0.61	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	B	901	2	-	1/6/23/26	0/1/1/1
10	NAG	C	901	3	-	0/6/23/26	0/1/1/1
10	NAG	B	902	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	B	901	NAG	C1-O5-C5	2.11	115.01	112.19
10	B	902	NAG	C1-O5-C5	2.07	114.96	112.19

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	B	901	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	157/157 (100%)	-0.41	0	100 100	30, 48, 80, 109	0
2	B	301/312 (96%)	-0.06	0	100 100	36, 74, 118, 151	1 (0%)
3	C	270/344 (78%)	0.19	10 (3%)	45 25	38, 80, 145, 164	0
All	All	728/813 (89%)	-0.04	10 (1%)	73 53	30, 68, 136, 164	1 (0%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	104	ILE	3.5
3	C	102	PRO	3.4
3	C	54	SER	3.3
3	C	33	THR	3.1
3	C	117	GLY	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MAN	E	4	11/12	0.16	0.15	171,175,182,184	0
8	BMA	H	3	11/12	0.25	0.13	132,138,143,144	0
6	BMA	F	3	11/12	0.28	0.13	143,146,148,149	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MAN	E	5	11/12	0.45	0.14	175,179,184,185	0
9	BMA	I	3	11/12	0.57	0.17	174,177,182,183	0
9	NAG	I	2	14/15	0.58	0.15	162,171,174,175	0
5	BMA	E	3	11/12	0.61	0.15	158,168,174,179	0
9	NAG	I	1	14/15	0.61	0.14	162,167,170,171	0
4	NAG	D	2	14/15	0.67	0.12	150,155,162,162	0
6	NAG	F	2	14/15	0.71	0.14	118,129,138,142	0
8	FUC	H	4	10/11	0.74	0.15	90,93,95,95	0
7	NAG	J	2	14/15	0.74	0.11	114,116,119,121	0
9	MAN	I	4	11/12	0.76	0.15	180,184,187,188	0
4	NAG	D	1	14/15	0.80	0.11	129,136,143,147	0
7	NAG	G	2	14/15	0.83	0.14	118,126,130,131	0
8	NAG	H	2	14/15	0.83	0.12	106,112,120,125	0
7	NAG	G	1	14/15	0.85	0.10	107,112,115,118	0
5	NAG	E	2	14/15	0.86	0.14	132,139,148,153	0
8	NAG	H	1	14/15	0.86	0.11	82,88,93,101	0
7	NAG	J	1	14/15	0.86	0.10	94,98,103,109	0
4	FUC	D	3	10/11	0.87	0.13	140,141,144,145	0
6	NAG	F	1	14/15	0.88	0.14	97,101,107,116	0
5	NAG	E	1	14/15	0.89	0.13	113,120,126,129	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain G:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
10	NAG	B	901	14/15	0.21	0.18	146,149,153,156	0
10	NAG	C	901	14/15	0.43	0.15	180,182,186,188	0
10	NAG	B	902	14/15	0.54	0.11	124,132,140,143	0
11	CL	C	908	1/1	0.97	0.15	55,55,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.