



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 04:42 PM EDT

PDB ID : 3WOQ / pdb_00003woq
Title : Crystal structure of the DAP BII hexapeptide complex III
Authors : Sakamoto, Y.; Suzuki, Y.; Iizuka, I.; Tateoka, C.; Roppongi, S.; Fujimoto, M.;
Nonaka, T.; Ogasawara, W.; Tanaka, N.
Deposited on : 2013-12-29
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	698	
1	B	698	
2	C	6	
2	D	6	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	802	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11667 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called dipeptidyl aminopeptidase BII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	697	Total	C	N	O	S	0	0	0
			5362	3394	935	1014	19			
1	B	697	Total	C	N	O	S	0	0	0
			5362	3394	935	1014	19			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	86	ALA	HIS	engineered mutation	UNP V5YM14
A	224	ALA	ASP	engineered mutation	UNP V5YM14
A	657	ALA	SER	engineered mutation	UNP V5YM14
B	86	ALA	HIS	engineered mutation	UNP V5YM14
B	224	ALA	ASP	engineered mutation	UNP V5YM14
B	657	ALA	SER	engineered mutation	UNP V5YM14

- Molecule 2 is a protein called Angiotensin IV.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	0	0	0
			55	40	8	7			
2	D	6	Total	C	N	O	0	0	0
			55	40	8	7			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		
4	B	2	Total	Zn	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	448	Total 448	O 448	0	0
5	B	322	Total 322	O 322	0	0
5	C	2	Total 2	O 2	0	0
5	D	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: dipeptidyl aminopeptidase BII




- Molecule 1: dipeptidyl aminopeptidase BII




- Molecule 2: Angiotensin IV





- Molecule 2: Angiotensin IV



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	121.70Å 121.70Å 218.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.92 – 1.82 39.92 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.3 (39.92-1.82) 99.3 (39.92-1.82)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.84 (at 1.82Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.185 , 0.227 0.194 , 0.232	Depositor DCC
R_{free} test set	7299 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.012	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 34.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11667	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.35 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.8092e-03.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.26	10/5479 (0.2%)	1.15	18/7427 (0.2%)
1	B	1.16	5/5479 (0.1%)	1.11	8/7427 (0.1%)
2	C	1.42	1/58 (1.7%)	1.13	0/79
2	D	1.34	1/58 (1.7%)	1.43	1/79 (1.3%)
All	All	1.21	17/11074 (0.2%)	1.13	27/15012 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	4	HIS	CA-C	7.53	1.60	1.53
1	A	68	GLY	N-CA	-7.49	1.37	1.45
1	A	290	GLN	C-O	-6.23	1.16	1.24
1	B	291	TRP	CA-C	6.08	1.58	1.52
2	D	4	HIS	CA-C	5.85	1.59	1.52
1	A	231	TYR	N-CA	5.73	1.52	1.45
1	A	334	ASN	CG-ND2	-5.69	1.21	1.33
1	A	92	ILE	N-CA	-5.68	1.40	1.46
1	A	691	THR	N-CA	-5.40	1.39	1.46
1	A	657	ALA	C-O	5.13	1.30	1.23
1	B	325	MET	C-O	-5.10	1.18	1.24
1	B	201	ALA	CA-C	-5.05	1.47	1.53
1	A	719	ASN	CA-C	-5.05	1.46	1.53
1	B	266	PHE	C-O	-5.05	1.17	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	347	ALA	CA-C	5.04	1.59	1.52
1	B	335	TYR	N-CA	5.04	1.52	1.46
1	A	176	LEU	N-CA	-5.00	1.40	1.46

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67	LEU	CA-C-N	10.55	132.72	122.27
1	A	67	LEU	C-N-CA	10.55	132.72	122.27
2	D	4	HIS	CB-CA-C	9.23	124.36	109.41
1	A	674	ASP	CA-C-N	6.60	126.73	122.18
1	A	674	ASP	C-N-CA	6.60	126.73	122.18
1	A	207	GLY	N-CA-C	6.26	120.47	112.83
1	B	690	MET	CG-SD-CE	-6.24	87.17	100.90
1	B	709	TYR	CA-C-N	-5.86	113.93	120.14
1	B	709	TYR	C-N-CA	-5.86	113.93	120.14
1	A	319	VAL	N-CA-C	5.79	116.57	110.72
1	A	202	PRO	CA-C-N	-5.75	114.03	119.89
1	A	202	PRO	C-N-CA	-5.75	114.03	119.89
1	B	579	LEU	N-CA-C	5.74	118.64	110.10
1	B	470	ASP	N-CA-C	-5.71	101.83	110.28
1	B	301	LYS	N-CA-C	5.41	117.18	111.28
1	A	709	TYR	CA-C-N	-5.39	114.17	119.99
1	A	709	TYR	C-N-CA	-5.39	114.17	119.99
1	A	261	LEU	CA-C-N	5.31	126.15	121.58
1	A	261	LEU	C-N-CA	5.31	126.15	121.58
1	A	109	ALA	CA-C-N	-5.26	113.50	119.28
1	A	109	ALA	C-N-CA	-5.26	113.50	119.28
1	B	708	VAL	CB-CA-C	-5.22	106.41	111.05
1	B	655	GLY	CA-C-O	5.20	124.09	119.56
1	A	536	LEU	N-CA-C	5.09	116.83	111.28
1	A	562	ASP	N-CA-C	-5.08	105.83	111.36
1	A	590	ALA	CA-C-N	5.04	124.53	119.19
1	A	590	ALA	C-N-CA	5.04	124.53	119.19

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	67	LEU	Peptide
1	A	68	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5362	0	5283	41	0
1	B	5362	0	5283	45	0
2	C	55	0	54	1	0
2	D	55	0	54	4	0
3	A	30	0	40	5	0
3	B	24	0	32	2	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	448	0	0	5	2
5	B	322	0	0	1	0
5	C	2	0	0	0	0
5	D	3	0	0	3	0
All	All	11667	0	10746	89	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:HIS:ND1	3:A:802:GOL:O1	2.14	0.81
1:A:595:MET:HE3	1:B:595:MET:CE	2.10	0.80
1:B:470:ASP:O	1:B:471:GLN:HB2	1.81	0.80
1:A:287:ASN:HD22	1:A:383:LEU:HD11	1.49	0.78
1:A:287:ASN:ND2	1:A:383:LEU:HD11	2.00	0.76
1:A:595:MET:HE3	1:B:595:MET:HE3	1.68	0.73
1:B:470:ASP:O	1:B:471:GLN:CB	2.38	0.72
1:A:354:GLU:HG2	5:A:1273:HOH:O	1.94	0.66
1:A:93:GLN:NE2	1:A:447:ARG:HE	1.97	0.63
1:A:515:LYS:HG2	5:A:1019:HOH:O	2.00	0.61
1:B:472:ARG:NH2	1:B:483:ASN:HD21	1.99	0.60
1:A:277:ASN:HD22	1:A:683:ASN:HD21	1.48	0.60
1:B:84:ASN:HD22	1:B:657:ALA:HB1	1.68	0.58
1:B:255:LYS:HE3	3:B:803:GOL:O3	2.03	0.58
2:D:4:HIS:CD2	2:D:5:PRO:HD2	2.39	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:HIS:HB3	1:A:606:VAL:HG22	1.86	0.57
1:B:354:GLU:O	1:B:358:VAL:HG23	2.04	0.57
2:D:4:HIS:C	5:D:103:HOH:O	2.48	0.56
1:A:306:MET:HE3	1:A:455:GLN:HB3	1.87	0.56
2:D:4:HIS:CB	5:D:103:HOH:O	2.53	0.56
2:D:4:HIS:HB3	5:D:103:HOH:O	2.05	0.56
1:B:718:MET:O	1:B:719:ASN:HB2	2.06	0.54
1:A:255:LYS:HE3	3:A:802:GOL:O3	2.08	0.54
1:A:100:ASN:ND2	5:A:1119:HOH:O	2.40	0.54
1:A:61:MET:HE1	1:A:581:ILE:HD11	1.90	0.54
1:B:411:LEU:HD21	1:B:441:LEU:HD11	1.89	0.53
1:A:217:MET:HE2	1:A:333:LYS:NZ	2.24	0.52
1:A:222:THR:H	1:A:645:ASN:HD21	1.57	0.52
1:B:222:THR:H	1:B:645:ASN:HD21	1.57	0.52
1:B:120:PRO:HD2	1:B:443:GLN:HE21	1.75	0.52
1:B:472:ARG:HH22	1:B:483:ASN:ND2	2.08	0.52
1:B:84:ASN:ND2	1:B:657:ALA:HB1	2.24	0.52
1:B:622:LEU:C	1:B:622:LEU:HD23	2.34	0.52
1:A:653:THR:H	1:A:656:ASN:HD22	1.57	0.51
1:B:93:GLN:NE2	1:B:447:ARG:HE	2.08	0.51
1:A:41:LEU:HD13	1:A:583:PHE:CG	2.46	0.51
1:B:472:ARG:HH22	1:B:483:ASN:HD21	1.58	0.51
1:B:41:LEU:HG	1:B:46:LEU:HD22	1.92	0.51
2:C:4:HIS:CD2	2:C:5:PRO:HD2	2.46	0.51
1:B:251:LYS:HB3	1:B:252:HIS:CD2	2.46	0.50
1:A:622:LEU:C	1:A:622:LEU:HD23	2.37	0.50
1:A:43:LYS:O	1:A:43:LYS:HD3	2.12	0.49
1:B:653:THR:H	1:B:656:ASN:HD22	1.60	0.49
1:B:225:PHE:CG	1:B:701:LEU:HG	2.48	0.48
1:A:84:ASN:HD22	1:A:657:ALA:HB1	1.77	0.48
1:A:281:LEU:HB2	1:A:284:GLU:HG3	1.96	0.48
1:A:149:LEU:HG	1:A:153:LYS:HE2	1.96	0.48
1:B:132:ASP:OD1	1:B:134:THR:OG1	2.19	0.48
1:B:197:ARG:HD2	1:B:231:TYR:CE1	2.49	0.47
1:A:370:GLN:N	1:A:371:PRO:CD	2.77	0.47
1:A:93:GLN:HE22	1:A:447:ARG:HE	1.59	0.47
1:B:220:ARG:HB3	1:B:645:ASN:HD22	1.80	0.47
1:A:257:ALA:O	3:A:805:GOL:H12	2.15	0.47
1:B:346:ASP:OD2	1:B:349:GLY:HA3	2.15	0.46
1:A:276:THR:HA	1:A:683:ASN:OD1	2.15	0.46
1:B:207:GLY:HA2	1:B:223:GLY:O	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:41:LEU:HD13	1:B:583:PHE:CG	2.51	0.45
1:B:431:GLU:HA	1:B:434:LEU:HD22	1.98	0.45
1:B:93:GLN:HE22	1:B:447:ARG:HE	1.63	0.45
1:B:299:HIS:HD2	1:B:459:TYR:OH	1.98	0.45
1:B:303:GLN:HE21	1:B:456:LEU:HD22	1.81	0.45
1:B:590:ALA:O	1:B:690:MET:HE1	2.17	0.45
1:A:197:ARG:O	1:A:228:TYR:HA	2.18	0.44
1:A:61:MET:CE	1:A:581:ILE:HD11	2.47	0.44
1:A:252:HIS:CE1	3:A:802:GOL:O1	2.71	0.43
1:B:412:TYR:CZ	1:B:416:ILE:HD11	2.52	0.43
1:B:109:ALA:HB1	1:B:114:ASP:HB2	2.00	0.43
1:A:217:MET:HE2	1:A:333:LYS:CE	2.48	0.43
1:B:599:PRO:O	1:B:646:TYR:HA	2.19	0.43
1:B:525:ILE:O	1:B:529:VAL:HG23	2.19	0.43
1:B:276:THR:HA	1:B:683:ASN:OD1	2.18	0.42
1:A:41:LEU:HG	1:A:46:LEU:HD22	2.02	0.42
1:B:412:TYR:CE2	1:B:529:VAL:HA	2.54	0.42
1:B:37:ILE:O	1:B:41:LEU:HB2	2.20	0.42
1:B:84:ASN:HD21	1:B:673:PHE:HA	1.85	0.41
1:A:387:LYS:NZ	5:A:1241:HOH:O	2.54	0.41
1:A:218:TRP:HA	1:A:219:PRO:C	2.45	0.41
1:A:473:VAL:HG13	1:A:534:THR:HG21	2.02	0.41
1:B:576:ASN:O	1:B:577:LEU:HB2	2.20	0.41
1:A:121:ASN:H	1:A:443:GLN:NE2	2.19	0.41
1:A:253:TRP:CE2	3:A:802:GOL:H32	2.55	0.41
1:A:306:MET:HE1	1:A:455:GLN:C	2.46	0.41
1:B:344:ARG:NH1	1:B:690:MET:HG3	2.36	0.41
1:B:421:PRO:HG2	1:B:424:GLU:OE1	2.21	0.41
1:B:718:MET:O	1:B:719:ASN:CB	2.70	0.40
3:B:804:GOL:C1	5:B:952:HOH:O	2.69	0.40
1:A:392:ARG:H	1:A:471:GLN:HE21	1.67	0.40
1:A:706:GLN:NE2	5:A:989:HOH:O	2.48	0.40
1:A:595:MET:HE3	1:B:595:MET:HE1	1.95	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1316:HOH:O	5:A:1318:HOH:O[8_665]	1.63	0.57
5:A:979:HOH:O	5:A:1318:HOH:O[8_665]	2.11	0.09

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	695/698 (100%)	680 (98%)	15 (2%)	0	100	100
1	B	695/698 (100%)	673 (97%)	22 (3%)	0	100	100
2	C	4/6 (67%)	2 (50%)	2 (50%)	0	100	100
2	D	4/6 (67%)	2 (50%)	2 (50%)	0	100	100
All	All	1398/1408 (99%)	1357 (97%)	41 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	538/539 (100%)	528 (98%)	10 (2%)	50	37
1	B	538/539 (100%)	522 (97%)	16 (3%)	36	17
2	C	6/6 (100%)	5 (83%)	1 (17%)	2	0
2	D	6/6 (100%)	6 (100%)	0	100	100
All	All	1088/1090 (100%)	1061 (98%)	27 (2%)	42	25

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	LEU
1	A	41	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	234	LYS
1	A	312	LYS
1	A	368	LYS
1	A	386	SER
1	A	427	SER
1	A	446	ARG
1	A	461	LEU
1	A	560	LEU
1	B	41	LEU
1	B	54	SER
1	B	59	ASP
1	B	60	PRO
1	B	138	LYS
1	B	149	LEU
1	B	251	LYS
1	B	313	GLN
1	B	339	LEU
1	B	365	GLN
1	B	373	LEU
1	B	434	LEU
1	B	461	LEU
1	B	483	ASN
1	B	492	LEU
1	B	549	ASN
2	C	4	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (31) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	ASN
1	A	93	GLN
1	A	190	ASN
1	A	277	ASN
1	A	287	ASN
1	A	290	GLN
1	A	303	GLN
1	A	313	GLN
1	A	350	GLN
1	A	385	GLN
1	A	443	GLN
1	A	471	GLN
1	A	549	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	585	ASN
1	A	645	ASN
1	A	656	ASN
1	B	84	ASN
1	B	93	GLN
1	B	190	ASN
1	B	252	HIS
1	B	290	GLN
1	B	299	HIS
1	B	303	GLN
1	B	385	GLN
1	B	443	GLN
1	B	540	GLN
1	B	585	ASN
1	B	645	ASN
1	B	656	ASN
1	B	683	ASN
2	D	4	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	804	-	5,5,5	0.34	0	5,5,5	0.61	0
3	GOL	B	802	-	5,5,5	0.30	0	5,5,5	0.41	0
3	GOL	B	804	-	5,5,5	0.47	0	5,5,5	0.68	0
3	GOL	B	803	-	5,5,5	0.73	0	5,5,5	0.40	0
3	GOL	A	802	-	5,5,5	0.36	0	5,5,5	1.17	1 (20%)
3	GOL	A	803	-	5,5,5	0.60	0	5,5,5	0.31	0
3	GOL	B	801	-	5,5,5	0.79	0	5,5,5	0.34	0
3	GOL	A	805	-	5,5,5	0.42	0	5,5,5	0.92	0
3	GOL	A	801	-	5,5,5	0.64	0	5,5,5	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	804	-	-	0/4/4/4	-
3	GOL	B	802	-	-	4/4/4/4	-
3	GOL	B	804	-	-	0/4/4/4	-
3	GOL	B	803	-	-	0/4/4/4	-
3	GOL	A	802	-	-	2/4/4/4	-
3	GOL	A	803	-	-	2/4/4/4	-
3	GOL	B	801	-	-	0/4/4/4	-
3	GOL	A	805	-	-	4/4/4/4	-
3	GOL	A	801	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	GOL	O1-C1-C2	-2.23	100.34	110.38

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	805	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	802	GOL	O1-C1-C2-O2
3	A	802	GOL	O1-C1-C2-C3
3	A	805	GOL	O1-C1-C2-C3
3	B	802	GOL	O1-C1-C2-C3
3	B	802	GOL	C1-C2-C3-O3
3	A	805	GOL	O2-C2-C3-O3
3	B	802	GOL	O2-C2-C3-O3
3	A	802	GOL	O1-C1-C2-O2
3	A	803	GOL	O2-C2-C3-O3
3	A	805	GOL	O1-C1-C2-O2
3	A	803	GOL	C1-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	804	GOL	1	0
3	B	803	GOL	1	0
3	A	802	GOL	4	0
3	A	805	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	697/698 (99%)	-0.16	5 (0%) 84 88	10, 22, 37, 55	0
1	B	697/698 (99%)	0.38	44 (6%) 26 30	12, 28, 49, 76	0
2	C	6/6 (100%)	0.96	2 (33%) 1 1	13, 16, 44, 47	0
2	D	6/6 (100%)	1.03	3 (50%) 0 0	15, 18, 47, 47	0
All	All	1406/1408 (99%)	0.12	54 (3%) 44 50	10, 25, 45, 76	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	478	ALA	4.2
1	B	469	ALA	3.9
2	C	4	HIS	3.4
1	B	377	ALA	3.4
1	B	482	GLY	3.3
1	B	549	ASN	3.1
1	B	364	GLY	3.1
1	B	358	VAL	3.1
1	B	369	GLY	3.1
1	B	555	VAL	3.1
1	B	373	LEU	2.9
1	B	143	ALA	2.9
1	B	481	GLY	2.9
1	B	352	LEU	2.8
1	B	362	LEU	2.8
1	B	492	LEU	2.8
1	B	366	GLY	2.8
1	B	473	VAL	2.8
1	B	375	ALA	2.7
1	B	374	ASP	2.6
1	B	467	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	364	GLY	2.6
1	A	68	GLY	2.5
1	B	367	ALA	2.5
1	B	368	LYS	2.5
2	D	4	HIS	2.5
1	B	474	ALA	2.5
1	B	466	LYS	2.5
1	B	483	ASN	2.4
1	B	487	ALA	2.4
1	B	50	PRO	2.4
1	B	486	ALA	2.3
2	D	6	PHE	2.3
1	B	476	VAL	2.3
1	B	361	TRP	2.2
1	A	367	ALA	2.2
1	B	485	ALA	2.2
2	C	6	PHE	2.2
1	B	470	ASP	2.2
1	B	568	GLY	2.2
2	D	5	PRO	2.2
1	B	551	ALA	2.2
1	B	620	LYS	2.2
1	B	370	GLN	2.2
1	B	371	PRO	2.2
1	B	423	ALA	2.2
1	B	563	TYR	2.1
1	B	465	ILE	2.1
1	A	240	ALA	2.1
1	B	356	ALA	2.1
1	B	360	GLY	2.1
1	B	135	LYS	2.1
1	A	721	GLY	2.1
1	B	359	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	802	6/6	0.73	0.19	54,58,61,61	0
3	GOL	B	803	6/6	0.83	0.14	37,47,51,52	0
3	GOL	B	804	6/6	0.83	0.18	46,50,53,54	0
3	GOL	A	804	6/6	0.87	0.14	38,41,48,50	0
3	GOL	A	803	6/6	0.91	0.14	40,43,45,46	0
3	GOL	A	805	6/6	0.92	0.11	24,36,37,47	0
3	GOL	A	802	6/6	0.94	0.10	28,32,36,37	0
3	GOL	B	801	6/6	0.96	0.07	17,21,22,23	0
3	GOL	A	801	6/6	0.96	0.06	16,18,20,22	0
4	ZN	B	806	1/1	0.96	0.06	37,37,37,37	0
4	ZN	A	807	1/1	0.97	0.06	29,29,29,29	0
4	ZN	B	805	1/1	0.97	0.16	45,45,45,45	0
4	ZN	A	806	1/1	0.97	0.14	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.