



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:40 AM UTC

PDB ID : 1WPY / pdb_00001wp
Title : Crystal Structure Of Biotin-(Acetyl-CoA-Carboxylase) ligase From Pyrococcus Horikoshii Ot3 in complex with biotin
Authors : Bagautdinov, B.; Kunishima, N.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2004-09-17
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	
1	B	235	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called biotin--[acetyl-CoA-carboxylase] ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	235	Total	C	N	O	Se	0	0	0
			1837	1185	311	336	5			
1	B	235	Total	C	N	O	Se	0	3	0
			1844	1190	311	338	5			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP O57883
A	44	MSE	MET	modified residue	UNP O57883
A	143	MSE	MET	modified residue	UNP O57883
A	176	MSE	MET	modified residue	UNP O57883
A	186	MSE	MET	modified residue	UNP O57883
B	1	MSE	MET	modified residue	UNP O57883
B	44	MSE	MET	modified residue	UNP O57883
B	143	MSE	MET	modified residue	UNP O57883
B	176	MSE	MET	modified residue	UNP O57883
B	186	MSE	MET	modified residue	UNP O57883

- Molecule 2 is BIOTIN (CCD ID: BTN) (formula: C₁₀H₁₆N₂O₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			16	10	2	3	1		
2	B	1	Total	C	N	O	S	0	0
			16	10	2	3	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	341	Total	O	0	0
			341	341		
3	B	297	Total	O	0	0
			297	297		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: biotin--[acetyl-CoA-carboxylase] ligase



- Molecule 1: biotin--[acetyl-CoA-carboxylase] ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	38.42Å 82.74Å 72.71Å 90.00° 103.17° 90.00°	Depositor
Resolution (Å)	33.56 – 1.60 33.56 – 1.60	Depositor EDS
% Data completeness (in resolution range)	96.4 (33.56-1.60) 96.8 (33.56-1.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.63 (at 1.60Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.194 , 0.213 0.194 , 0.213	Depositor DCC
R_{free} test set	2898 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	15.2	Xtriage
Anisotropy	0.401	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4351	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BTN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1861	0.85	6/2503 (0.2%)
1	B	0.39	0/1880	0.95	9/2529 (0.4%)
All	All	0.39	0/3741	0.90	15/5032 (0.3%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	71	PRO	N-CA-C	-10.21	94.87	111.11
1	B	74	ASP	N-CA-C	-9.25	102.12	112.57
1	B	70	VAL	N-CA-CB	-8.30	99.59	111.21
1	B	72	GLN	N-CA-C	7.59	126.97	110.80
1	A	174	ASN	CA-C-N	7.17	127.79	119.47
1	A	174	ASN	C-N-CA	7.17	127.79	119.47
1	B	70	VAL	CA-C-N	-6.54	113.22	119.89
1	B	70	VAL	C-N-CA	-6.54	113.22	119.89
1	B	72	GLN	CB-CG-CD	-5.53	103.20	112.60
1	A	17	GLN	N-CA-C	-5.44	105.43	111.36
1	B	99	ILE	N-CA-C	5.33	115.93	108.89
1	A	2	LEU	N-CA-C	5.31	117.98	111.82
1	B	72	GLN	N-CA-CB	-5.17	101.75	110.49
1	A	135	LYS	N-CA-C	-5.14	103.25	110.50
1	A	99	ILE	N-CA-C	5.07	115.89	108.58

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1837	0	1934	43	0
1	B	1844	0	1941	63	0
2	A	16	0	15	0	0
2	B	16	0	15	0	0
3	A	341	0	0	10	0
3	B	297	0	0	11	0
All	All	4351	0	3905	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:72:GLN:O	1:B:72:GLN:HG3	1.63	0.95
1:B:76:PRO:O	1:B:79[A]:VAL:HG12	1.70	0.90
1:B:79[B]:VAL:HG21	1:B:227:TYR:HE1	1.37	0.90
1:B:186:MSE:HE3	1:B:188:LEU:HD11	1.59	0.84
1:B:49:LEU:HB3	1:B:51:ARG:NH1	1.91	0.83
1:A:186:MSE:HE3	1:A:188:LEU:HD11	1.64	0.79
1:A:44:MSE:HG2	3:A:1615:HOH:O	1.84	0.78
1:B:79[B]:VAL:HG21	1:B:227:TYR:CE1	2.17	0.78
1:B:72:GLN:O	1:B:72:GLN:CG	2.35	0.75
1:A:70:VAL:HG11	3:A:1642:HOH:O	1.87	0.74
1:B:70:VAL:HG13	1:B:71:PRO:HD2	1.71	0.73
1:B:48:ARG:HD2	1:B:227:TYR:HE2	1.54	0.72
1:B:69:LYS:HA	3:B:1546:HOH:O	1.88	0.71
1:B:226:ILE:HD12	1:B:226:ILE:O	1.91	0.70
1:A:75:LEU:H	1:A:76:PRO:CD	2.05	0.69
1:B:75:LEU:HB3	1:B:76:PRO:HD3	1.76	0.67
1:A:175:PRO:HG3	3:A:1642:HOH:O	1.95	0.67
1:B:67:SER:HB2	3:B:1427:HOH:O	1.94	0.66
1:A:165:ASP:O	1:A:169:LEU:HD23	1.96	0.65
1:A:18:GLU:OE2	1:A:41:LYS:HE3	1.95	0.65
1:A:74:ASP:HA	1:A:77:LYS:HZ2	1.60	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:LYS:HA	3:B:1338:HOH:O	1.97	0.64
1:A:167:LEU:HG	3:A:1629:HOH:O	1.96	0.63
1:B:27:LYS:HD2	1:B:120:LYS:HE2	1.79	0.63
1:B:70:VAL:O	1:B:71:PRO:C	2.39	0.63
1:B:67:SER:N	1:B:68:PRO:HD3	2.13	0.63
1:A:70:VAL:HB	1:A:71:PRO:HD2	1.81	0.62
1:B:51:ARG:HG3	1:B:51:ARG:HH11	1.66	0.61
1:B:176:MSE:HG2	3:B:1569:HOH:O	2.01	0.61
1:B:204:ALA:HA	1:B:215:ILE:HD13	1.82	0.60
1:B:48:ARG:HA	1:B:116:LEU:HD13	1.83	0.59
1:B:70:VAL:C	1:B:71:PRO:O	2.39	0.59
1:B:73:LYS:HD3	1:B:73:LYS:H	1.69	0.58
1:B:96:ASP:HB3	3:B:1521:HOH:O	2.04	0.58
1:B:68:PRO:HG3	1:B:168:TYR:CE1	2.39	0.57
1:A:74:ASP:HA	1:A:77:LYS:NZ	2.20	0.57
1:A:68:PRO:C	1:A:69:LYS:HG3	2.30	0.56
1:B:49:LEU:HB3	1:B:51:ARG:HH11	1.71	0.54
1:B:79[A]:VAL:HG11	1:B:227:TYR:CE1	2.43	0.53
1:B:135:LYS:HE3	3:B:1463:HOH:O	2.08	0.53
1:B:227:TYR:CD2	1:B:228:GLY:N	2.77	0.53
1:A:186:MSE:HE1	1:A:207:ILE:HG12	1.91	0.52
1:A:77:LYS:HA	3:A:1359:HOH:O	2.08	0.52
1:A:198:GLY:O	1:A:199:SER:HB3	2.09	0.52
1:A:70:VAL:HB	1:A:71:PRO:CD	2.39	0.52
1:B:56:PRO:HG2	1:B:134:ASN:HB2	1.92	0.51
1:B:70:VAL:N	3:B:1546:HOH:O	2.45	0.50
1:A:232:LEU:C	1:A:232:LEU:HD23	2.36	0.50
1:A:186:MSE:HE1	1:A:207:ILE:HG23	1.93	0.50
1:A:191:ARG:HG2	1:A:235:LEU:HD12	1.92	0.50
1:B:68:PRO:HG3	1:B:168:TYR:HE1	1.75	0.49
1:A:5:LYS:HE2	3:A:1403:HOH:O	2.12	0.49
1:B:72:GLN:HA	3:B:1339:HOH:O	2.12	0.49
1:A:75:LEU:HD13	1:A:75:LEU:O	2.13	0.49
1:A:75:LEU:HB3	1:A:76:PRO:HD3	1.94	0.49
1:B:76:PRO:HD3	3:B:1338:HOH:O	2.11	0.49
1:A:166:ARG:NH2	3:A:1629:HOH:O	2.47	0.48
1:B:101:TRP:CE2	1:B:102:PRO:HB3	2.49	0.47
1:B:59:GLY:HA3	1:B:61:TRP:CZ2	2.49	0.47
1:B:227:TYR:HD2	1:B:228:GLY:N	2.12	0.47
1:B:49:LEU:O	1:B:51:ARG:HG3	2.14	0.47
1:B:79[B]:VAL:HG22	1:B:117:VAL:CG2	2.44	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:PRO:HG2	1:A:134:ASN:HB2	1.96	0.47
1:A:75:LEU:O	1:A:78:ILE:HG12	2.15	0.47
1:B:48:ARG:C	1:B:50:ASN:H	2.23	0.47
1:A:75:LEU:C	1:A:77:LYS:H	2.22	0.46
1:B:7:SER:HB2	1:B:169:LEU:HD21	1.98	0.46
1:B:68:PRO:HG3	1:B:168:TYR:OH	2.15	0.46
1:A:75:LEU:H	1:A:76:PRO:HD2	1.77	0.46
1:B:42:GLN:HB2	1:B:55:SER:HB2	1.97	0.46
1:A:206:ASP:CG	1:A:207:ILE:H	2.24	0.46
1:B:79[B]:VAL:CG2	1:B:227:TYR:HE1	2.18	0.46
1:A:1:MSE:HE3	1:B:157:PHE:CE2	2.51	0.45
1:B:74:ASP:HB3	1:B:77:LYS:HD2	1.97	0.45
1:B:179:LEU:O	1:B:183:ARG:HG3	2.15	0.45
1:A:206:ASP:CG	1:A:207:ILE:N	2.75	0.45
1:B:206:ASP:CG	1:B:207:ILE:H	2.25	0.45
1:B:12:ARG:NE	1:B:32:GLU:OE2	2.50	0.45
1:A:42:GLN:HB2	1:A:55:SER:HB2	2.00	0.44
1:A:101:TRP:HA	1:A:102:PRO:HA	1.80	0.44
1:B:17:GLN:O	1:B:40:ASP:HB2	2.18	0.44
1:B:183:ARG:HA	1:B:186:MSE:HE2	2.00	0.44
1:B:84:VAL:O	1:B:88:GLU:HG3	2.18	0.44
1:A:76:PRO:HB3	3:A:1484:HOH:O	2.18	0.43
1:A:75:LEU:N	1:A:76:PRO:CD	2.72	0.43
1:A:183:ARG:HG3	1:A:206:ASP:OD1	2.19	0.43
1:B:80:PHE:HB3	1:B:182:VAL:CG2	2.48	0.43
1:A:217:LEU:HD12	1:A:217:LEU:N	2.34	0.43
1:B:67:SER:N	1:B:68:PRO:CD	2.82	0.43
1:B:217:LEU:HD12	1:B:217:LEU:N	2.33	0.43
1:B:75:LEU:CB	3:B:1338:HOH:O	2.67	0.42
1:A:204:ALA:HA	1:A:215:ILE:HD13	2.01	0.42
1:B:75:LEU:C	1:B:75:LEU:HD23	2.45	0.41
1:B:227:TYR:CD2	1:B:227:TYR:C	2.97	0.41
1:A:75:LEU:HG	3:A:1552:HOH:O	2.20	0.41
1:A:205:GLU:O	1:A:206:ASP:HB2	2.20	0.41
1:A:62:LEU:C	1:A:62:LEU:HD12	2.46	0.41
1:B:69:LYS:HA	1:B:69:LYS:HD3	1.80	0.41
1:B:194:ILE:HG13	1:B:200:PHE:CZ	2.55	0.41
1:A:80:PHE:HB3	1:A:182:VAL:CG2	2.50	0.40
1:A:84:VAL:O	1:A:88:GLU:HG3	2.21	0.40
1:B:80:PHE:HB3	1:B:182:VAL:HG21	2.02	0.40
1:B:191:ARG:HD3	3:B:1393:HOH:O	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:LYS:NZ	3:A:1631:HOH:O	2.55	0.40
1:B:49:LEU:CB	1:B:51:ARG:HH11	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/235 (99%)	223 (96%)	9 (4%)	1 (0%)	30	14
1	B	236/235 (100%)	226 (96%)	7 (3%)	3 (1%)	9	2
All	All	469/470 (100%)	449 (96%)	16 (3%)	4 (1%)	14	3

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	75	LEU
1	B	49	LEU
1	B	68	PRO
1	B	230	VAL

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BTN	B	1302	-	17,17,17	0.95	1 (5%)	23,23,23	1.23	2 (8%)
2	BTN	A	1301	-	17,17,17	1.05	1 (5%)	23,23,23	1.19	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BTN	B	1302	-	-	2/7/28/28	0/2/2/2
2	BTN	A	1301	-	-	2/7/28/28	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1301	BTN	C2-S1	-2.69	1.78	1.82
2	B	1302	BTN	C2-S1	-2.32	1.79	1.82

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1302	BTN	C2-C4-C5	-3.16	105.03	108.89
2	A	1301	BTN	C2-C4-C5	-2.94	105.30	108.89
2	A	1301	BTN	C5-C6-S1	-2.87	102.11	106.06
2	B	1302	BTN	C5-C6-S1	-2.84	102.16	106.06

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1301	BTN	C9-C10-C11-O11
2	A	1301	BTN	C9-C10-C11-O12
2	B	1302	BTN	C9-C10-C11-O12
2	B	1302	BTN	C9-C10-C11-O11

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/235 (97%)	0.32	19 (8%) 17 17	7, 17, 41, 56	0
1	B	230/235 (97%)	0.51	28 (12%) 8 8	9, 18, 43, 54	3 (1%)
All	All	460/470 (97%)	0.42	47 (10%) 12 12	7, 18, 43, 56	3 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	227	TYR	6.3
1	B	49	LEU	5.3
1	B	71	PRO	5.1
1	A	74	ASP	5.0
1	B	122	ASP	5.0
1	A	75	LEU	4.7
1	B	70	VAL	4.3
1	B	76	PRO	4.2
1	B	51	ARG	4.1
1	B	68	PRO	4.0
1	B	75	LEU	4.0
1	B	121	GLY	4.0
1	A	70	VAL	3.8
1	B	230	VAL	3.8
1	B	119	GLY	3.5
1	B	138	ASN	3.5
1	A	139	GLY	3.4
1	A	76	PRO	3.4
1	A	71	PRO	3.3
1	B	69	LYS	3.2
1	B	120	LYS	3.2
1	B	229	ASP	3.2
1	A	49	LEU	3.1
1	A	73	LYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	73	LYS	2.9
1	B	50	ASN	2.9
1	A	235	LEU	2.8
1	B	67	SER	2.8
1	B	235	LEU	2.7
1	A	122	ASP	2.7
1	A	197	ASP	2.7
1	A	69	LYS	2.7
1	B	72	GLN	2.7
1	B	74	ASP	2.7
1	B	48	ARG	2.6
1	B	226	ILE	2.6
1	A	196	GLY	2.5
1	A	198	GLY	2.5
1	B	139	GLY	2.5
1	A	72	GLN	2.4
1	B	232	LEU	2.4
1	A	121	GLY	2.4
1	A	119	GLY	2.3
1	B	228	GLY	2.3
1	A	138	ASN	2.2
1	A	231	SER	2.2
1	B	197	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BTN	A	1301	16/16	0.95	0.08	13,16,25,27	0
2	BTN	B	1302	16/16	0.95	0.08	17,18,28,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.