



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 03:47 PM UTC

PDB ID : 3WPS / pdb_00003wps
Title : crystal structure of the GAP domain of MgcRacGAP(S387D)
Authors : Murayama, K.; Kato-murayama, M.; Shirouzu, M.; Kitamura, T.; Yokoyama, S.
Deposited on : 2014-01-15
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	208	5% 66% 28% • •
1	B	208	7% 62% 31% • •

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3279 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Rac GTPase-activating protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	201	Total	C	N	O	S	Se	0	0	0
			1588	1010	272	294	3	9			
1	B	201	Total	C	N	O	S	Se	0	0	0
			1588	1010	272	294	3	9			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	339	GLY	-	expression tag	UNP Q9H0H5
A	340	SER	-	expression tag	UNP Q9H0H5
A	341	SER	-	expression tag	UNP Q9H0H5
A	342	GLY	-	expression tag	UNP Q9H0H5
A	343	SER	-	expression tag	UNP Q9H0H5
A	344	SER	-	expression tag	UNP Q9H0H5
A	345	GLY	-	expression tag	UNP Q9H0H5
A	387	ASP	SER	engineered mutation	UNP Q9H0H5
A	518	SER	LEU	SEE REMARK 999	UNP Q9H0H5
B	339	GLY	-	expression tag	UNP Q9H0H5
B	340	SER	-	expression tag	UNP Q9H0H5
B	341	SER	-	expression tag	UNP Q9H0H5
B	342	GLY	-	expression tag	UNP Q9H0H5
B	343	SER	-	expression tag	UNP Q9H0H5
B	344	SER	-	expression tag	UNP Q9H0H5
B	345	GLY	-	expression tag	UNP Q9H0H5
B	387	ASP	SER	engineered mutation	UNP Q9H0H5
B	518	SER	LEU	SEE REMARK 999	UNP Q9H0H5

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	49	Total	O	0	0
			49	49		
3	B	44	Total	O	0	0
			44	44		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Rac GTPase-activating protein 1



• Molecule 1: Rac GTPase-activating protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	77.32Å 77.32Å 108.74Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.66 – 2.70 38.66 – 2.71	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.66-2.70) 99.9 (38.66-2.71)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.50 (at 2.72Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.210 , 0.259 0.207 , 0.247	Depositor DCC
R_{free} test set	960 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	45.7	Xtriage
Anisotropy	0.470	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for -h,-k,l 0.047 for h,-h-k,-l 0.023 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3279	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/1608	0.96	9/2163 (0.4%)
1	B	0.42	0/1608	0.98	7/2163 (0.3%)
All	All	0.43	0/3216	0.97	16/4326 (0.4%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	534	LEU	CA-C-N	8.99	129.07	119.90
1	B	534	LEU	C-N-CA	8.99	129.07	119.90
1	A	503	THR	N-CA-C	6.37	118.10	111.03
1	B	449	THR	N-CA-C	6.13	117.96	111.28
1	A	536	LEU	N-CA-C	-5.91	104.91	111.71
1	B	506	ALA	CB-CA-C	-5.84	109.82	116.54
1	B	362	ILE	CA-C-N	5.71	125.64	119.76
1	B	362	ILE	C-N-CA	5.71	125.64	119.76
1	A	534	LEU	CA-C-N	5.62	125.62	119.89
1	A	534	LEU	C-N-CA	5.62	125.62	119.89
1	A	511	ASN	CA-C-N	5.56	125.87	119.92
1	A	511	ASN	C-N-CA	5.56	125.87	119.92
1	A	405	THR	N-CA-C	5.47	117.10	111.03
1	A	377	GLY	N-CA-C	5.41	122.55	115.36
1	B	428	ASN	N-CA-C	5.17	119.30	113.15
1	A	506	ALA	CB-CA-C	-5.08	109.75	115.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1588	0	1625	49	0
1	B	1588	0	1625	60	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	49	0	0	2	0
3	B	44	0	0	2	0
All	All	3279	0	3250	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:408:LEU:HB2	1:B:411:LYS:HE3	1.46	0.96
1:A:492:ASP:H	1:A:495:ASN:HD22	1.25	0.85
1:A:396:LEU:O	1:A:400:PHE:HB2	1.82	0.79
1:B:405:THR:O	1:B:407:PRO:HD3	1.82	0.78
1:B:372:GLU:HG3	1:B:409:LEU:HB2	1.65	0.78
1:A:440:ARG:HD3	1:A:443:MSE:CE	2.15	0.77
1:B:443:MSE:HE2	1:B:523:ARG:HB2	1.66	0.76
1:A:440:ARG:HD3	1:A:443:MSE:HE2	1.70	0.73
1:B:475:PHE:CD1	1:B:543:MSE:HE1	2.24	0.71
1:A:406:VAL:HB	1:A:407:PRO:HD3	1.71	0.71
1:B:347:GLY:O	1:B:348:GLU:HG3	1.92	0.70
1:A:346:ILE:N	1:A:430:LYS:HA	2.08	0.68
1:A:385:ARG:HB2	1:A:491:MSE:CE	2.26	0.66
1:A:424:ASP:HA	1:A:427:ARG:HH12	1.60	0.66
1:A:424:ASP:HA	1:A:427:ARG:NH1	2.11	0.66
1:B:458:MSE:HE2	1:B:530:ARG:HB2	1.78	0.66
1:B:545:VAL:O	1:B:546:GLU:HB2	1.95	0.65
1:B:492:ASP:OD1	1:B:495:ASN:ND2	2.31	0.64
1:A:477:MSE:CE	1:A:480:LEU:HD12	2.28	0.64
1:B:396:LEU:HD11	1:B:412:VAL:HG21	1.80	0.63
1:A:398:GLU:C	1:A:400:PHE:H	2.07	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:440:ARG:O	1:A:444:GLU:HG3	1.99	0.62
1:B:475:PHE:HD1	1:B:543:MSE:HE1	1.65	0.62
1:A:477:MSE:HG3	1:A:532:LEU:HD23	1.84	0.60
1:A:488:HIS:O	1:A:490:LYS:HG3	2.01	0.60
1:B:400:PHE:HE2	1:B:409:LEU:HD11	1.66	0.60
1:B:377:GLY:O	1:B:383:LEU:HD11	2.02	0.59
1:A:517:MSE:HE2	3:A:719:HOH:O	2.01	0.59
1:A:405:THR:O	1:A:408:LEU:HB2	2.02	0.59
1:B:475:PHE:HD1	1:B:543:MSE:CE	2.16	0.58
1:B:399:LYS:HZ1	1:B:411:LYS:HZ3	1.52	0.58
1:A:374:GLU:OE2	1:A:482:ARG:HD2	2.04	0.58
1:B:432:PRO:HB3	1:B:506:ALA:HB3	1.86	0.57
1:A:544:MSE:HE1	1:B:536:LEU:HD21	1.86	0.57
1:B:398:GLU:HG2	1:B:399:LYS:N	2.19	0.57
1:A:357:GLN:HA	1:A:361:MSE:HE3	1.87	0.56
1:B:470:ARG:HD2	1:B:542:PHE:CE1	2.40	0.56
1:B:492:ASP:H	1:B:495:ASN:HD22	1.53	0.56
1:B:512:PRO:HG2	1:B:517:MSE:HE2	1.88	0.56
1:B:415:ILE:HD12	1:B:418:ILE:HD12	1.88	0.56
1:A:372:GLU:HG3	1:A:409:LEU:HB3	1.89	0.55
1:B:431:GLU:HB2	1:B:432:PRO:HD2	1.89	0.55
1:B:420:SER:O	1:B:424:ASP:HB2	2.08	0.54
1:A:507:HIS:CG	1:A:512:PRO:HG2	2.42	0.54
1:A:385:ARG:HB2	1:A:491:MSE:HE1	1.89	0.54
1:B:443:MSE:CE	1:B:523:ARG:HB2	2.35	0.53
1:B:357:GLN:HG3	3:B:739:HOH:O	2.08	0.52
1:B:439:ASN:O	1:B:443:MSE:HG3	2.09	0.52
1:A:477:MSE:HE2	1:A:480:LEU:HD12	1.92	0.52
1:A:390:ASP:O	1:A:394:LYS:HG3	2.09	0.52
1:B:522:LYS:HD3	3:B:744:HOH:O	2.10	0.52
1:B:545:VAL:O	1:B:546:GLU:CB	2.58	0.51
1:B:359:SER:HB2	1:B:546:GLU:OE2	2.11	0.51
1:B:466:PRO:HB2	1:B:469:ASN:OD1	2.11	0.50
1:B:458:MSE:O	1:B:462:VAL:HG23	2.12	0.50
1:B:372:GLU:HG3	1:B:409:LEU:CB	2.40	0.49
1:B:499:VAL:O	1:B:502:PRO:HD2	2.11	0.49
1:B:477:MSE:O	1:B:481:GLN:HG3	2.13	0.49
1:A:395:GLU:O	1:A:398:GLU:N	2.45	0.49
1:A:442:PHE:CE1	1:A:458:MSE:HE1	2.47	0.49
1:A:442:PHE:CD1	1:A:458:MSE:HE1	2.47	0.49
1:B:538:TYR:O	1:B:541:GLN:HG2	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:470:ARG:HD2	1:B:542:PHE:CZ	2.48	0.48
1:A:443:MSE:HE3	1:A:523:ARG:NH1	2.28	0.48
1:B:427:ARG:NH1	1:B:502:PRO:O	2.47	0.48
1:B:399:LYS:NZ	1:B:411:LYS:HZ3	2.11	0.48
1:B:377:GLY:C	1:B:383:LEU:HD11	2.39	0.48
1:A:440:ARG:HA	1:A:443:MSE:HE2	1.95	0.47
1:B:545:VAL:O	1:B:545:VAL:HG12	2.15	0.47
1:A:440:ARG:CZ	1:A:443:MSE:HE1	2.45	0.47
1:B:399:LYS:NZ	1:B:411:LYS:NZ	2.62	0.47
1:A:546:GLU:OXT	1:A:546:GLU:HG2	2.15	0.46
1:A:380:GLU:O	1:A:489:THR:HA	2.15	0.46
1:A:443:MSE:HE3	1:A:523:ARG:CZ	2.46	0.46
1:A:477:MSE:O	1:A:481:GLN:HG3	2.15	0.46
1:A:481:GLN:O	1:A:485:GLN:HG3	2.15	0.46
1:B:524:GLN:HB2	1:B:525:PRO:HD3	1.98	0.46
1:B:383:LEU:O	1:B:384:TYR:HB2	2.16	0.45
1:A:484:ALA:HA	1:A:491:MSE:O	2.17	0.45
1:B:399:LYS:HG2	1:B:407:PRO:HG3	1.98	0.45
1:B:399:LYS:O	1:B:407:PRO:HG3	2.16	0.45
1:B:407:PRO:O	1:B:409:LEU:N	2.50	0.45
1:A:383:LEU:O	1:A:384:TYR:HB2	2.17	0.44
1:A:477:MSE:CE	1:A:477:MSE:HA	2.46	0.44
1:A:450:ASP:HB3	1:A:453:ASN:HB2	1.98	0.44
1:B:408:LEU:C	1:B:410:SER:H	2.26	0.43
1:A:372:GLU:HG3	1:A:409:LEU:CB	2.48	0.43
1:B:433:LEU:HD21	1:B:473:LEU:HA	1.99	0.43
1:B:458:MSE:HG3	1:B:530:ARG:HB3	2.00	0.43
1:A:517:MSE:O	1:A:521:ILE:HG13	2.19	0.43
1:A:470:ARG:CD	1:A:542:PHE:CE1	3.02	0.42
1:B:459:TYR:CZ	1:B:534:LEU:HD22	2.54	0.42
1:A:477:MSE:HA	1:A:477:MSE:HE3	2.02	0.42
1:B:350:MSE:SE	1:B:430:LYS:HD3	2.69	0.42
1:B:359:SER:HA	1:B:360:PRO:C	2.44	0.42
1:B:474:ALA:HA	1:B:539:TRP:CE3	2.55	0.42
1:A:432:PRO:HB3	1:A:506:ALA:HB3	2.02	0.42
1:A:480:LEU:HD22	1:A:496:LEU:HD22	2.01	0.42
1:B:399:LYS:HZ1	1:B:411:LYS:NZ	2.16	0.41
1:A:398:GLU:O	1:A:400:PHE:N	2.53	0.41
1:A:439:ASN:OD1	1:A:440:ARG:NH1	2.54	0.41
1:B:376:ARG:HG3	1:B:376:ARG:HH11	1.85	0.41
1:A:544:MSE:HG3	3:A:738:HOH:O	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:403:VAL:O	1:B:403:VAL:HG12	2.21	0.41
1:B:402:ARG:HG3	1:B:403:VAL:N	2.35	0.40
1:B:524:GLN:N	1:B:525:PRO:CD	2.84	0.40
1:B:439:ASN:HB3	1:B:508:ALA:HB2	2.02	0.40
1:A:403:VAL:HG13	1:A:408:LEU:HG	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/208 (96%)	181 (91%)	15 (8%)	3 (2%)	8	22
1	B	199/208 (96%)	184 (92%)	13 (6%)	2 (1%)	12	32
All	All	398/416 (96%)	365 (92%)	28 (7%)	5 (1%)	9	25

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	408	LEU
1	B	408	LEU
1	A	399	LYS
1	B	545	VAL
1	A	356	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/174 (103%)	177 (99%)	2 (1%)	65	85
1	B	179/174 (103%)	174 (97%)	5 (3%)	38	68
All	All	358/348 (103%)	351 (98%)	7 (2%)	48	76

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	357	GLN
1	A	477	MSE
1	B	376	ARG
1	B	398	GLU
1	B	402	ARG
1	B	413	ASP
1	B	424	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	357	GLN
1	A	460	GLN
1	A	481	GLN
1	A	495	ASN
1	A	511	ASN
1	B	371	ASN
1	B	375	GLN
1	B	453	ASN
1	B	460	GLN
1	B	495	ASN
1	B	519	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	601	-	4,4,4	0.39	0	6,6,6	0.08	0
2	SO4	A	600	-	4,4,4	0.38	0	6,6,6	0.12	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	192/208 (92%)	0.10	10 (5%) 33 29	26, 41, 91, 107	0
1	B	192/208 (92%)	0.18	14 (7%) 21 18	28, 48, 89, 106	0
All	All	384/416 (92%)	0.14	24 (6%) 26 23	26, 45, 91, 107	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	346	ILE	7.2
1	B	404	LYS	4.5
1	A	406	VAL	4.5
1	B	405	THR	4.0
1	B	346	ILE	3.8
1	A	403	VAL	3.7
1	B	406	VAL	3.5
1	A	404	LYS	3.5
1	B	408	LEU	3.2
1	B	347	GLY	3.2
1	A	347	GLY	3.0
1	A	348	GLU	2.9
1	B	403	VAL	2.8
1	B	348	GLU	2.8
1	B	546	GLU	2.8
1	A	401	LEU	2.6
1	B	401	LEU	2.5
1	B	400	PHE	2.5
1	A	546	GLU	2.4
1	B	399	LYS	2.4
1	A	408	LEU	2.3
1	B	397	LYS	2.2
1	B	391	ARG	2.2
1	A	400	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	B	601	5/5	0.77	0.19	118,119,119,120	0
2	SO4	A	600	5/5	0.79	0.17	105,105,106,106	0

6.5 Other polymers [i](#)

There are no such residues in this entry.