



# wwPDB X-ray Structure Validation Summary Report

Mar 13, 2026 – 11:49 PM UTC


PDB ID : 4WPH / pdb\_00004wph  
Title : Crystal structure of USP7 ubiquitin-like domains in compact conformation  
Authors : Pfoh, R.; Lacedao, I.; Saridakis, V.  
Deposited on : 2014-10-18  
Resolution : 2.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2995 (2.94-2.90)
Clashscore	190562	3213 (2.94-2.90)
Ramachandran outliers	187476	3128 (2.94-2.90)
Sidechain outliers	187428	3130 (2.94-2.90)
RSRZ outliers	180081	2995 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	 2% 82% 7% 10%
1	B	375	 2% 79% 8% 13%
2	C	11	 82% 9% 9%
2	D	11	 73% 9% 18%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5582 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	337	2748	1743	474	514	17	0	2	0
1	B	326	2635	1674	445	500	16	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	514	MET	-	initiating methionine	UNP Q93009
A	515	GLY	-	expression tag	UNP Q93009
A	516	SER	-	expression tag	UNP Q93009
A	517	SER	-	expression tag	UNP Q93009
A	518	HIS	-	expression tag	UNP Q93009
A	519	HIS	-	expression tag	UNP Q93009
A	520	HIS	-	expression tag	UNP Q93009
A	521	HIS	-	expression tag	UNP Q93009
A	522	HIS	-	expression tag	UNP Q93009
A	523	HIS	-	expression tag	UNP Q93009
A	524	SER	-	expression tag	UNP Q93009
A	525	SER	-	expression tag	UNP Q93009
A	526	GLY	-	expression tag	UNP Q93009
A	527	ARG	-	expression tag	UNP Q93009
A	528	GLU	-	expression tag	UNP Q93009
A	529	ASN	-	expression tag	UNP Q93009
A	530	LEU	-	expression tag	UNP Q93009
A	531	TYR	-	expression tag	UNP Q93009
A	532	PHE	-	expression tag	UNP Q93009
A	533	GLN	-	expression tag	UNP Q93009
A	534	GLY	-	expression tag	UNP Q93009
B	514	MET	-	initiating methionine	UNP Q93009
B	515	GLY	-	expression tag	UNP Q93009
B	516	SER	-	expression tag	UNP Q93009
B	517	SER	-	expression tag	UNP Q93009

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	518	HIS	-	expression tag	UNP Q93009
B	519	HIS	-	expression tag	UNP Q93009
B	520	HIS	-	expression tag	UNP Q93009
B	521	HIS	-	expression tag	UNP Q93009
B	522	HIS	-	expression tag	UNP Q93009
B	523	HIS	-	expression tag	UNP Q93009
B	524	SER	-	expression tag	UNP Q93009
B	525	SER	-	expression tag	UNP Q93009
B	526	GLY	-	expression tag	UNP Q93009
B	527	ARG	-	expression tag	UNP Q93009
B	528	GLU	-	expression tag	UNP Q93009
B	529	ASN	-	expression tag	UNP Q93009
B	530	LEU	-	expression tag	UNP Q93009
B	531	TYR	-	expression tag	UNP Q93009
B	532	PHE	-	expression tag	UNP Q93009
B	533	GLN	-	expression tag	UNP Q93009
B	534	GLY	-	expression tag	UNP Q93009

- Molecule 2 is a protein called ICP0.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	C	10	Total	C	N	O	S	0	0	0
			80	47	21	11	1			
2	D	9	Total	C	N	O	S	0	0	0
			76	45	20	10	1			

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Cl	0	0
			3	3		
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	21	Total	O	0	0
			21	21		
4	B	17	Total	O	0	0
			17	17		

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	C	1	Total	O	0	0
			1	1		



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.34Å 92.34Å 190.28Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.92 30.00 – 2.92	Depositor EDS
% Data completeness (in resolution range)	99.5 (30.00-2.92) 99.5 (30.00-2.92)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 2.90Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.233 , 0.262 0.232 , 0.261	Depositor DCC
$R_{free}$ test set	1041 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	65.6	Xtrriage
Anisotropy	0.189	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 55.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	5582	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.63	0/2806	0.84	0/3788
1	B	0.63	0/2689	0.82	1/3633 (0.0%)
2	C	0.81	0/80	0.92	0/103
2	D	0.68	0/76	0.90	0/97
All	All	0.63	0/5651	0.84	1/7621 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	778	SER	N-CA-C	6.00	118.45	110.53

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2748	0	2696	21	0
1	B	2635	0	2547	16	0
2	C	80	0	91	1	0
2	D	76	0	89	1	0
3	A	3	0	0	0	0
3	B	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	21	0	0	1	0
4	B	17	0	0	0	0
4	C	1	0	0	0	0
All	All	5582	0	5423	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 35 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:625:MET:HE1	1:A:661:PHE:HB2	1.69	0.74
1:B:623:TRP:CE2	1:B:686:MET:HE3	2.35	0.61
1:A:623:TRP:CE2	1:A:686:MET:HE3	2.36	0.61
1:A:583:MET:HE1	1:A:765:ILE:HG12	1.87	0.56
1:A:628:ARG:NH2	4:A:2111:HOH:O	2.38	0.56

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/375 (89%)	325 (98%)	8 (2%)	0	100	100
1	B	318/375 (85%)	312 (98%)	6 (2%)	0	100	100
2	C	8/11 (73%)	8 (100%)	0	0	100	100
2	D	7/11 (64%)	6 (86%)	1 (14%)	0	100	100
All	All	666/772 (86%)	651 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/343 (88%)	297 (98%)	6 (2%)	48	77
1	B	287/343 (84%)	279 (97%)	8 (3%)	38	70
2	C	8/9 (89%)	8 (100%)	0	100	100
2	D	8/9 (89%)	8 (100%)	0	100	100
All	All	606/704 (86%)	592 (98%)	14 (2%)	44	74

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	611	GLN
1	B	613	MET
1	B	835	MET
1	B	824	LYS
1	B	830	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	607	GLN
1	A	683	HIS
1	A	700	ASN
1	A	815	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	337/375 (89%)	0.12	8 (2%) 59 50	35, 73, 126, 181	2 (0%)
1	B	326/375 (86%)	0.02	7 (2%) 63 54	38, 71, 133, 157	0
2	C	10/11 (90%)	-0.02	0 100 100	54, 61, 78, 99	0
2	D	9/11 (81%)	0.57	0 100 100	88, 93, 112, 117	0
All	All	682/772 (88%)	0.08	15 (2%) 62 53	35, 73, 133, 181	2 (0%)

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	541	VAL	2.9
1	B	799	CYS	2.8
1	A	540	LEU	2.7
1	A	669	LEU	2.7
1	B	807	PRO	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	A	2001	1/1	0.78	0.10	89,89,89,89	0
3	CL	B	2001	1/1	0.79	0.13	90,90,90,90	0
3	CL	A	2003	1/1	0.83	0.07	82,82,82,82	0
3	CL	A	2002	1/1	0.94	0.11	74,74,74,74	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.