



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:41 PM UTC

PDB ID : 6WP2 / pdb\_00006wp2  
Title : The Crystal Structure of Apo Zinc-Bound Domain Swapped-Trimer Q108K:K40D:T53A:R58L:Q38F:Q4F:F57H Variant of HCRBPII  
Authors : Ghanbarpour, A.; Geiger, J.  
Deposited on : 2020-04-26  
Resolution : 2.48 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	133	
1	B	133	
1	C	133	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3328 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinol-binding protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	133	1101	699	189	208	5	0	3	0
1	B	133	1086	689	182	209	6	0	2	0
1	C	133	1088	690	184	209	5	0	2	0

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	4	PHE	GLN	engineered mutation	UNP P50120
A	38	PHE	GLN	engineered mutation	UNP P50120
A	40	ASP	LYS	engineered mutation	UNP P50120
A	53	ALA	THR	engineered mutation	UNP P50120
A	57	HIS	PHE	engineered mutation	UNP P50120
A	58	LEU	ARG	engineered mutation	UNP P50120
A	108	LYS	GLN	engineered mutation	UNP P50120
B	4	PHE	GLN	engineered mutation	UNP P50120
B	38	PHE	GLN	engineered mutation	UNP P50120
B	40	ASP	LYS	engineered mutation	UNP P50120
B	53	ALA	THR	engineered mutation	UNP P50120
B	57	HIS	PHE	engineered mutation	UNP P50120
B	58	LEU	ARG	engineered mutation	UNP P50120
B	108	LYS	GLN	engineered mutation	UNP P50120
C	4	PHE	GLN	engineered mutation	UNP P50120
C	38	PHE	GLN	engineered mutation	UNP P50120
C	40	ASP	LYS	engineered mutation	UNP P50120
C	53	ALA	THR	engineered mutation	UNP P50120
C	57	HIS	PHE	engineered mutation	UNP P50120
C	58	LEU	ARG	engineered mutation	UNP P50120
C	108	LYS	GLN	engineered mutation	UNP P50120

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by

depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total	Zn	0	0
			5	5		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	5	Total O 5 5	0	0
5	B	2	Total O 2 2	0	0
5	C	6	Total O 6 6	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Retinol-binding protein 2



- Molecule 1: Retinol-binding protein 2



- Molecule 1: Retinol-binding protein 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.51Å 74.45Å 94.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.17 – 2.48 47.17 – 2.48	Depositor EDS
% Data completeness (in resolution range)	99.2 (47.17-2.48) 99.8 (47.17-2.48)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.88 (at 2.48Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.222 , 0.312 0.224 , 0.314	Depositor DCC
$R_{free}$ test set	1572 reflections (9.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.4	Xtrriage
Anisotropy	0.780	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 34.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3328	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.40	1/1133 (0.1%)	0.65	1/1527 (0.1%)
1	B	0.42	0/1114	0.62	0/1503
1	C	0.39	0/1116	0.65	0/1507
All	All	0.40	1/3363 (0.0%)	0.64	1/4537 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	100	GLU	CG-CD	-5.22	1.39	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	100	GLU	CA-CB-CG	-7.01	100.08	114.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1101	0	1063	26	0
1	B	1086	0	1041	41	0
1	C	1088	0	1043	38	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	5	0	0	0	0
2	C	1	0	0	0	0
3	A	12	0	16	0	0
3	B	12	0	16	3	0
3	C	6	0	8	0	0
4	C	4	0	3	0	0
5	A	5	0	0	0	0
5	B	2	0	0	0	0
5	C	6	0	0	2	0
All	All	3328	0	3190	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:ILE:HG12	1:A:51:THR:HG22	1.59	0.84
1:B:7:THR:HG22	1:B:41:VAL:HG12	1.62	0.81
1:C:72:GLU:OE2	5:C:301:HOH:O	2.07	0.71
1:B:20:MET:HE2	1:B:30:ARG:HG2	1.73	0.69
1:A:33:ALA:O	1:A:34:VAL:HB	1.93	0.67
1:C:42:ILE:HG12	1:C:51:THR:HG22	1.77	0.66
1:A:17:GLU:OE1	1:A:30[B]:ARG:NH1	2.29	0.65
1:B:8:TRP:CZ3	1:C:132:LYS:HB2	2.33	0.64
1:C:92:VAL:HG13	1:C:109:TRP:HB3	1.81	0.62
1:C:66:VAL:HG11	1:C:88:TRP:CD1	2.35	0.61
1:A:132:LYS:HB2	1:C:8:TRP:CZ3	2.34	0.61
1:B:1:THR:O	1:C:91:ASP:HB3	2.03	0.58
1:C:85:LEU:O	1:C:95:CYS:HA	2.04	0.58
1:A:86:VAL:HG22	1:A:95:CYS:HB2	1.85	0.57
1:B:10[B]:MET:HE2	1:B:13:ASN:HB2	1.86	0.56
1:C:101:LYS:HG2	1:C:121:CYS:SG	2.46	0.55
1:C:74:THR:CG2	1:C:80:ARG:H	2.19	0.55
1:B:10[B]:MET:SD	3:B:202:GOL:O1	2.58	0.55
1:A:9:GLU:OE2	1:B:131:LYS:HE2	2.06	0.55
1:C:20:MET:HG2	1:C:25:ILE:HD12	1.88	0.55
1:B:42:ILE:HG12	1:B:51:THR:HG22	1.90	0.54
1:B:9:GLU:HA	1:B:39:THR:HG22	1.90	0.54
1:A:54:THR:HG22	1:B:59:ASN:CG	2.32	0.53
1:B:10[A]:MET:HE2	1:C:128:GLN:HB3	1.91	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:74:THR:HG23	1:C:74:THR:O	2.08	0.52
1:B:10[B]:MET:HE3	1:B:38:PHE:HD2	1.76	0.51
1:B:29:THR:HA	1:B:32:ILE:HD12	1.93	0.51
1:B:106:TRP:CZ3	1:B:108:LYS:HE3	2.45	0.51
1:B:20:MET:HE1	1:B:30:ARG:HA	1.93	0.50
1:A:91:ASP:HB3	1:C:1:THR:O	2.12	0.50
1:C:1:THR:HG22	1:C:3:ASP:H	1.77	0.50
1:B:4:PHE:HA	1:C:110:ILE:HD11	1.94	0.50
1:A:8:TRP:HB3	1:B:130:PHE:HB3	1.93	0.49
1:A:1:THR:HG22	1:A:3:ASP:H	1.76	0.49
1:B:10[A]:MET:HB3	1:B:38:PHE:O	2.11	0.49
1:C:74:THR:O	1:C:79:ASN:N	2.46	0.49
1:A:128:GLN:HB3	1:C:10:MET:HE2	1.94	0.48
1:B:10[B]:MET:HB2	1:C:130:PHE:CE1	2.48	0.48
1:C:74:THR:HG22	1:C:80:ARG:O	2.14	0.47
1:A:20:MET:HG2	1:A:25:ILE:HD12	1.96	0.47
1:B:92:VAL:HG22	1:B:109:TRP:HB3	1.97	0.47
1:C:74:THR:HG23	1:C:80:ARG:H	1.80	0.47
1:C:86:VAL:HG22	1:C:95:CYS:HB2	1.97	0.47
1:C:65:THR:O	1:C:68:VAL:HG22	2.14	0.46
1:A:54:THR:HG22	1:B:59:ASN:OD1	2.15	0.46
1:B:115:LEU:HB3	1:B:130:PHE:HB2	1.97	0.46
1:B:27:PHE:CE1	1:B:31:LYS:HE2	2.51	0.46
1:A:130:PHE:HB3	1:C:8:TRP:HB3	1.98	0.45
1:C:115:LEU:HD12	1:C:116:TYR:H	1.81	0.45
1:A:74:THR:HG21	1:A:104:ARG:NH2	2.32	0.45
1:B:2:ARG:HG2	1:C:88:TRP:CH2	2.52	0.45
1:B:52:LYS:HD3	1:C:61:ASP:OD1	2.17	0.44
1:B:23:LEU:HD22	1:C:104:ARG:NH2	2.33	0.44
1:B:29:THR:HG23	5:C:302:HOH:O	2.17	0.44
1:B:31:LYS:HD3	1:B:31:LYS:HA	1.72	0.44
1:A:65:THR:HA	1:C:47:ASP:O	2.18	0.44
1:C:85:LEU:HD12	1:C:86:VAL:H	1.83	0.44
1:A:9:GLU:HB2	1:B:131:LYS:HG3	1.99	0.43
1:B:41:VAL:HG22	1:B:52:LYS:HB3	1.99	0.43
1:A:66:VAL:HG11	1:A:88:TRP:CD1	2.53	0.43
1:A:44:GLN:HB2	1:A:49:PHE:CE2	2.54	0.43
1:B:86:VAL:HG22	1:B:95:CYS:HB2	2.00	0.43
1:C:74:THR:HG22	1:C:80:ARG:H	1.83	0.43
1:B:10[A]:MET:HE3	1:B:10[A]:MET:HB2	1.94	0.43
1:A:73:TYR:CD1	1:A:75:LYS:HG3	2.54	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:4:PHE:CD1	1:B:4:PHE:N	2.87	0.42
1:B:10[A]:MET:HG2	3:B:202:GOL:HO1	1.84	0.42
1:B:44:GLN:HE22	3:B:201:GOL:H12	1.85	0.42
1:B:24:ASP:OD2	1:C:101:LYS:NZ	2.47	0.42
1:A:77:LEU:HD23	1:A:77:LEU:HA	1.80	0.42
1:C:35:ARG:NH2	1:C:76:SER:O	2.35	0.42
1:A:75:LYS:C	1:A:77:LEU:H	2.28	0.42
1:B:5:ASN:OD1	1:B:44:GLN:N	2.45	0.42
1:A:75:LYS:HE3	1:A:75:LYS:HB2	1.89	0.41
1:B:41:VAL:O	1:B:51:THR:HA	2.19	0.41
1:A:101:LYS:HA	1:A:101:LYS:HD2	1.73	0.41
1:A:51:THR:O	1:B:61:ASP:HA	2.21	0.41
1:C:44:GLN:HB3	1:C:45:ASP:H	1.69	0.41
1:C:29:THR:HA	1:C:32:ILE:HD12	2.03	0.41
1:B:41:VAL:CG2	1:B:52:LYS:HB3	2.51	0.41
1:C:85:LEU:HD12	1:C:86:VAL:N	2.36	0.41
1:B:4:PHE:HE1	1:C:88:TRP:HH2	1.68	0.40
1:A:41:VAL:O	1:A:51:THR:HA	2.21	0.40
1:C:78:ASP:OD2	1:C:104:ARG:NH2	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	134/133 (101%)	125 (93%)	7 (5%)	2 (2%)	<a href="#">8</a> <a href="#">14</a>
1	B	133/133 (100%)	127 (96%)	6 (4%)	0	<a href="#">100</a> <a href="#">100</a>
1	C	133/133 (100%)	125 (94%)	5 (4%)	3 (2%)	<a href="#">5</a> <a href="#">7</a>
All	All	400/399 (100%)	377 (94%)	18 (4%)	5 (1%)	<a href="#">9</a> <a href="#">16</a>

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	34	VAL
1	A	53	ALA
1	C	45	ASP
1	C	112	GLY
1	C	74	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/119 (99%)	118 (100%)	0	100	100
1	B	117/119 (98%)	115 (98%)	2 (2%)	53	75
1	C	117/119 (98%)	117 (100%)	0	100	100
All	All	352/357 (99%)	350 (99%)	2 (1%)	86	90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	10[A]	MET
1	B	10[B]	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	81	HIS
1	B	44	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	C	202	-	5,5,5	0.98	0	5,5,5	0.95	0
3	GOL	A	207	-	5,5,5	0.98	0	5,5,5	1.01	0
4	ACT	C	203	-	3,3,3	1.45	0	3,3,3	1.23	0
3	GOL	A	206	-	5,5,5	1.24	1 (20%)	5,5,5	1.00	0
3	GOL	B	201	2	5,5,5	0.94	0	5,5,5	1.14	0
3	GOL	B	202	-	5,5,5	0.56	0	5,5,5	1.49	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	202	-	-	4/4/4/4	-
3	GOL	A	207	-	-	0/4/4/4	-
3	GOL	A	206	-	-	2/4/4/4	-
3	GOL	B	201	2	-	1/4/4/4	-
3	GOL	B	202	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	206	GOL	C3-C2	2.23	1.60	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	B	202	GOL	C3-C2-C1	-2.51	102.58	111.80

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	202	GOL	O1-C1-C2-C3
3	A	206	GOL	O1-C1-C2-C3
3	B	202	GOL	C1-C2-C3-O3
3	C	202	GOL	C1-C2-C3-O3
3	B	202	GOL	O2-C2-C3-O3
3	C	202	GOL	O1-C1-C2-O2
3	C	202	GOL	O2-C2-C3-O3
3	B	201	GOL	O2-C2-C3-O3
3	A	206	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	201	GOL	1	0
3	B	202	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	133/133 (100%)	0.39	5 (3%) 44 40	27, 57, 72, 87	3 (2%)
1	B	133/133 (100%)	0.35	3 (2%) 61 58	28, 56, 73, 89	2 (1%)
1	C	133/133 (100%)	0.46	2 (1%) 72 69	27, 58, 77, 92	2 (1%)
All	All	399/399 (100%)	0.40	10 (2%) 58 55	27, 57, 75, 92	7 (1%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	33	ALA	3.6
1	A	60[A]	TYR	3.4
1	B	1	THR	3.1
1	A	1	THR	2.5
1	B	115	LEU	2.1
1	B	4	PHE	2.1
1	A	53	ALA	2.1
1	C	112	GLY	2.1
1	A	42	ILE	2.1
1	C	14	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	C	202	6/6	0.60	0.12	84,93,100,100	0
3	GOL	A	206	6/6	0.64	0.20	52,68,78,83	0
3	GOL	A	207	6/6	0.70	0.14	79,82,87,91	0
3	GOL	B	202	6/6	0.87	0.15	37,48,64,71	0
4	ACT	C	203	4/4	0.93	0.12	36,42,42,48	0
2	ZN	A	204	1/1	0.95	0.14	31,31,31,31	1
3	GOL	B	201	6/6	0.96	0.14	38,72,76,76	0
2	ZN	A	201	1/1	0.96	0.06	49,49,49,49	0
2	ZN	C	201	1/1	0.97	0.04	51,51,51,51	0
2	ZN	A	205	1/1	0.98	0.04	68,68,68,68	0
2	ZN	A	202	1/1	0.99	0.09	56,56,56,56	0
2	ZN	A	203	1/1	1.00	0.08	35,35,35,35	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around ZN A 204:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ZN A 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ZN C 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ZN A 205:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around ZN A 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.