



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:32 PM UTC

PDB ID : 2WQW / pdb_00002wqw
Title : DOUBLE-DISULFIDE CROSS-LINKED CRYSTAL DIMER of the *Listeria monocytogenes* InlB internalin domain
Authors : Ferraris, D.M.; Heinz, D.W.; Niemann, H.H.
Deposited on : 2009-08-27
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	5% 86% 13%
1	B	286	2% 90% 9%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called INTERNALIN B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	0	0	0
			2256	1436	378	440	2			
1	B	286	Total	C	N	O	S	0	0	0
			2256	1436	378	440	2			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	206	CYS	GLY	engineered mutation	UNP P25147
A	227	CYS	ALA	engineered mutation	UNP P25147
A	242	ALA	CYS	engineered mutation	UNP P25147
B	206	CYS	GLY	engineered mutation	UNP P25147
B	227	CYS	ALA	engineered mutation	UNP P25147
B	242	ALA	CYS	engineered mutation	UNP P25147

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 7 4 3	0	0
2	A	1	Total C O 7 4 3	0	0
2	A	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0
2	B	1	Total C O 7 4 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	65	Total O 65 65	0	0
3	B	77	Total O 77 77	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: INTERNALIN B



• Molecule 1: INTERNALIN B



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	186.99Å 186.99Å 115.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.88 – 2.24 46.88 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.88-2.24) 99.3 (46.88-2.24)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.199 , 0.239 0.208 , 0.240	Depositor DCC
R_{free} test set	1839 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	36.7	Xtriage
Anisotropy	0.596	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 26.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -1/3*h+1/3*k+4/3*l,-k,2/3*h+1/3*k+1/3*l 0.010 for -2/3*h-1/3*k-4/3*l,-1/3*h-2/3*k+4/3*l,-1/3*h+1/3*k+1/3*l 0.000 for -h,1/3*h-1/3*k-4/3*l,-1/3*h-2/3*k+1/3*l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4717	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.29	3/2292 (0.1%)	1.10	7/3110 (0.2%)
1	B	1.26	2/2292 (0.1%)	1.12	6/3110 (0.2%)
All	All	1.28	5/4584 (0.1%)	1.11	13/6220 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	288	PRO	C-O	6.25	1.31	1.24
1	A	233	ASP	CB-CG	-6.22	1.36	1.52
1	A	100	VAL	CB-CG2	-5.67	1.33	1.52
1	B	176	ILE	CA-CB	-5.28	1.47	1.54
1	B	84	ASN	C-O	-5.06	1.17	1.23

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	310	ARG	NE-CZ-NH2	-10.85	109.43	119.20
1	B	310	ARG	NE-CZ-NH1	-10.12	111.38	121.50
1	B	310	ARG	NE-CZ-NH2	9.58	127.82	119.20
1	A	310	ARG	CD-NE-CZ	9.18	137.25	124.40
1	A	310	ARG	NE-CZ-NH1	7.60	129.10	121.50
1	B	310	ARG	CD-NE-CZ	7.27	134.58	124.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	293	GLU	N-CA-C	5.78	117.90	108.76
1	A	179	ILE	N-CA-C	5.71	116.99	111.91
1	A	294	VAL	CG1-CB-CG2	-5.49	98.73	110.80
1	B	300	GLN	CA-C-N	-5.38	114.24	119.78
1	B	300	GLN	C-N-CA	-5.38	114.24	119.78
1	A	154	ILE	CB-CA-C	-5.07	103.77	111.33
1	A	154	ILE	N-CA-C	5.03	115.59	109.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	310	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2256	0	2322	20	0
1	B	2256	0	2322	11	0
2	A	21	0	30	1	0
2	B	42	0	60	2	0
3	A	65	0	0	1	0
3	B	77	0	0	2	0
All	All	4717	0	4734	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:ILE:HD11	1:A:97:LEU:CD2	2.24	0.67
1:A:318:PRO:C	1:A:319:LEU:HD12	2.20	0.67
1:B:116:LEU:HA	1:B:119:LEU:HD22	1.84	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:81:ILE:HD11	1:B:97:LEU:CD2	2.34	0.57
1:A:216:SER:OG	1:A:238:PHE:CE2	2.58	0.56
1:A:83:ALA:HB3	1:A:105:LEU:HD23	1.90	0.53
1:A:279:GLU:O	1:A:280:LYS:C	2.52	0.52
1:A:81:ILE:HD11	1:A:97:LEU:HD23	1.90	0.52
1:A:184:ARG:O	1:A:184:ARG:HG2	2.10	0.52
1:A:216:SER:OG	1:A:238:PHE:CD2	2.62	0.51
1:B:279:GLU:O	1:B:280:LYS:C	2.53	0.51
1:A:260:LYS:NZ	3:A:2051:HOH:O	2.43	0.49
1:B:184:ARG:HG2	1:B:184:ARG:O	2.11	0.48
1:B:83:ALA:HB3	1:B:105:LEU:HD23	1.95	0.48
1:A:161:VAL:O	1:A:184:ARG:NH1	2.45	0.48
1:A:318:PRO:O	1:A:319:LEU:HD12	2.13	0.48
1:A:299:TYR:CD2	1:A:299:TYR:C	2.93	0.47
1:B:161:VAL:O	1:B:184:ARG:NH1	2.48	0.47
2:B:1322:PEG:H32	3:B:2019:HOH:O	2.15	0.47
1:A:145:LYS:NZ	2:A:1324:PEG:O2	2.49	0.46
1:A:161:VAL:HG21	1:B:310:ARG:NH2	2.30	0.45
1:B:261:ASN:ND2	1:B:300:GLN:OE1	2.50	0.45
1:A:201:ILE:HD12	1:A:201:ILE:C	2.43	0.43
1:A:114:LYS:N	1:A:115:PRO:CD	2.82	0.42
1:A:263:ASP:OD1	1:A:265:SER:OG	2.33	0.42
1:A:202:VAL:N	1:A:203:PRO:CD	2.83	0.42
1:B:216:SER:OG	1:B:238:PHE:CE2	2.67	0.42
2:B:1324:PEG:O2	3:B:2075:HOH:O	2.21	0.41
1:A:290:PHE:CE1	1:A:321:GLU:O	2.72	0.41
1:B:81:ILE:HD11	1:B:97:LEU:HD23	2.01	0.41
1:B:201:ILE:HD12	1:B:201:ILE:C	2.46	0.41
1:A:122:LEU:HA	1:A:122:LEU:HD12	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/286 (99%)	271 (95%)	12 (4%)	1 (0%)	30	30
1	B	284/286 (99%)	268 (94%)	16 (6%)	0	100	100
All	All	568/572 (99%)	539 (95%)	28 (5%)	1 (0%)	43	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/265 (100%)	255 (96%)	10 (4%)	29	34
1	B	265/265 (100%)	252 (95%)	13 (5%)	22	23
All	All	530/530 (100%)	507 (96%)	23 (4%)	26	28

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	LYS
1	A	66	VAL
1	A	118	ASN
1	A	139	LYS
1	A	161	VAL
1	A	195	ASP
1	A	233	ASP
1	A	250	GLN
1	A	292	ASN
1	A	300	GLN
1	B	45	LYS
1	B	66	VAL
1	B	118	ASN
1	B	119	LEU
1	B	137	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	161	VAL
1	B	171	LEU
1	B	195	ASP
1	B	233	ASP
1	B	250	GLN
1	B	289	GLU
1	B	292	ASN
1	B	300	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	80	GLN
1	A	173	ASN
1	A	212	ASN
1	B	72	GLN
1	B	292	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PEG	B	1326	-	6,6,6	0.88	0	5,5,5	1.65	2 (40%)
2	PEG	A	1322	-	6,6,6	0.63	0	5,5,5	0.39	0
2	PEG	B	1322	-	6,6,6	0.74	0	5,5,5	0.64	0
2	PEG	B	1327	-	6,6,6	1.01	0	5,5,5	0.97	0
2	PEG	A	1323	-	6,6,6	0.72	0	5,5,5	0.61	0
2	PEG	B	1324	-	6,6,6	0.50	0	5,5,5	0.42	0
2	PEG	B	1323	-	6,6,6	0.62	0	5,5,5	0.63	0
2	PEG	B	1325	-	6,6,6	0.61	0	5,5,5	0.33	0
2	PEG	A	1324	-	6,6,6	1.00	0	5,5,5	1.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	B	1326	-	-	3/4/4/4	-
2	PEG	A	1322	-	-	2/4/4/4	-
2	PEG	B	1322	-	-	2/4/4/4	-
2	PEG	B	1327	-	-	3/4/4/4	-
2	PEG	A	1323	-	-	2/4/4/4	-
2	PEG	B	1324	-	-	1/4/4/4	-
2	PEG	B	1323	-	-	2/4/4/4	-
2	PEG	B	1325	-	-	3/4/4/4	-
2	PEG	A	1324	-	-	3/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1326	PEG	O2-C2-C1	2.38	120.60	110.11
2	B	1326	PEG	O2-C3-C4	2.19	119.78	110.11

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1326	PEG	C1-C2-O2-C3
2	A	1324	PEG	O1-C1-C2-O2
2	B	1322	PEG	O2-C3-C4-O4
2	B	1323	PEG	O1-C1-C2-O2
2	B	1325	PEG	O2-C3-C4-O4
2	B	1326	PEG	O1-C1-C2-O2
2	A	1322	PEG	O1-C1-C2-O2
2	B	1323	PEG	O2-C3-C4-O4
2	A	1324	PEG	O2-C3-C4-O4
2	A	1322	PEG	O2-C3-C4-O4
2	B	1322	PEG	O1-C1-C2-O2
2	A	1323	PEG	O2-C3-C4-O4
2	B	1327	PEG	C4-C3-O2-C2
2	B	1325	PEG	C1-C2-O2-C3
2	B	1324	PEG	O1-C1-C2-O2
2	B	1326	PEG	O2-C3-C4-O4
2	B	1327	PEG	O2-C3-C4-O4
2	A	1323	PEG	O1-C1-C2-O2
2	A	1324	PEG	C4-C3-O2-C2
2	B	1325	PEG	C4-C3-O2-C2
2	B	1327	PEG	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1322	PEG	1	0
2	B	1324	PEG	1	0
2	A	1324	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/286 (100%)	0.06	13 (4%) 38 36	22, 34, 69, 97	0
1	B	286/286 (100%)	-0.04	7 (2%) 59 61	23, 35, 60, 90	0
All	All	572/572 (100%)	0.01	20 (3%) 47 46	22, 35, 62, 97	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	288	PRO	4.3
1	A	290	PHE	3.9
1	A	320	LYS	3.4
1	A	286	HIS	3.3
1	A	289	GLU	3.0
1	A	292	ASN	2.9
1	B	36	GLU	2.6
1	B	321	GLU	2.6
1	B	320	LYS	2.4
1	A	319	LEU	2.4
1	B	288	PRO	2.4
1	A	287	LEU	2.4
1	A	250	GLN	2.2
1	A	238	PHE	2.2
1	A	251	SER	2.1
1	B	37	THR	2.1
1	A	321	GLU	2.1
1	B	251	SER	2.0
1	A	253	LEU	2.0
1	B	289	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PEG	B	1327	7/7	0.77	0.17	67,70,71,71	0
2	PEG	A	1324	7/7	0.78	0.22	66,71,73,74	0
2	PEG	B	1323	7/7	0.82	0.15	45,47,50,51	0
2	PEG	B	1322	7/7	0.86	0.13	59,59,61,62	0
2	PEG	B	1326	7/7	0.87	0.14	51,51,53,54	0
2	PEG	B	1324	7/7	0.89	0.11	46,51,59,61	0
2	PEG	A	1323	7/7	0.90	0.12	50,52,54,56	0
2	PEG	B	1325	7/7	0.90	0.12	51,54,55,55	0
2	PEG	A	1322	7/7	0.92	0.10	46,47,54,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.