



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:41 AM UTC

PDB ID : 3WQF / pdb_00003wqf
Title : D-threo-3-hydroxyaspartate dehydratase from *Delftia* sp. HT23 in the metal-free form
Authors : Yasutake, Y.; Matsumoto, Y.; Wada, M.
Deposited on : 2014-01-25
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	% 83% 13% ..
1	B	390	7% 80% 16% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PLP	A	401	-	X	-	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5893 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called D-threo-3-hydroxyaspartate dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	378	Total	C	N	O	S	0	0	0
			2813	1747	522	529	15			
1	B	379	Total	C	N	O	S	0	0	0
			2822	1752	524	531	15			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	GLY	-	expression tag	UNP B2DFG5
A	-8	HIS	-	expression tag	UNP B2DFG5
A	-7	HIS	-	expression tag	UNP B2DFG5
A	-6	HIS	-	expression tag	UNP B2DFG5
A	-5	HIS	-	expression tag	UNP B2DFG5
A	-4	HIS	-	expression tag	UNP B2DFG5
A	-3	HIS	-	expression tag	UNP B2DFG5
A	-2	ALA	-	expression tag	UNP B2DFG5
A	-1	MET	-	expression tag	UNP B2DFG5
A	0	SER	-	expression tag	UNP B2DFG5
B	-9	GLY	-	expression tag	UNP B2DFG5
B	-8	HIS	-	expression tag	UNP B2DFG5
B	-7	HIS	-	expression tag	UNP B2DFG5
B	-6	HIS	-	expression tag	UNP B2DFG5
B	-5	HIS	-	expression tag	UNP B2DFG5
B	-4	HIS	-	expression tag	UNP B2DFG5
B	-3	HIS	-	expression tag	UNP B2DFG5
B	-2	ALA	-	expression tag	UNP B2DFG5
B	-1	MET	-	expression tag	UNP B2DFG5
B	0	SER	-	expression tag	UNP B2DFG5

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	123	Total	O	0	0
			123	123		
3	B	105	Total	O	0	0
			105	105		

4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	157.84Å 157.84Å 157.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.92 – 2.30 49.92 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.92-2.30) 99.9 (49.92-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.28 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.198 , 0.239 0.194 , 0.235	Depositor DCC
R_{free} test set	2210 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for -l,-k,-h 0.007 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5893	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.24	6/2860 (0.2%)	1.20	8/3882 (0.2%)
1	B	1.27	3/2869 (0.1%)	1.19	5/3894 (0.1%)
All	All	1.26	9/5729 (0.2%)	1.20	13/7776 (0.2%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	172	HIS	CG-CD2	5.73	1.42	1.35
1	A	22	ARG	C-O	-5.50	1.17	1.24
1	A	321	HIS	CB-CG	-5.36	1.42	1.50
1	A	351	HIS	CG-CD2	5.23	1.41	1.35
1	B	107	LEU	N-CA	5.22	1.51	1.45
1	A	56	ALA	C-O	-5.21	1.17	1.24
1	B	81	ILE	C-O	5.19	1.29	1.24
1	A	140	HIS	CG-CD2	5.10	1.41	1.35
1	A	19	ARG	CZ-NH1	5.02	1.39	1.32

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	217	SER	N-CA-C	-6.91	98.85	109.14
1	A	24	ILE	N-CA-C	-6.41	104.25	110.72
1	A	321	HIS	CB-CA-C	-6.08	97.75	109.37
1	A	370	SER	N-CA-C	5.96	118.60	110.55
1	B	86	SER	N-CA-CB	5.67	118.71	109.69
1	B	294	GLN	N-CA-C	5.64	118.30	108.76
1	B	149	THR	CA-C-N	-5.59	112.84	119.05
1	B	149	THR	C-N-CA	-5.59	112.84	119.05
1	A	181	THR	CA-C-N	5.59	125.69	119.32
1	A	181	THR	C-N-CA	5.59	125.69	119.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	48	VAL	CA-C-N	-5.25	114.20	119.56
1	A	48	VAL	C-N-CA	-5.25	114.20	119.56
1	B	217	SER	N-CA-C	-5.04	101.08	108.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2813	0	2798	36	0
1	B	2822	0	2806	39	0
2	A	15	0	6	0	0
2	B	15	0	6	0	0
3	A	123	0	0	4	0
3	B	105	0	0	4	0
All	All	5893	0	5616	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:177:TYR:HA	1:B:248:ILE:HD11	1.44	0.98
1:B:25:ALA:O	1:B:29:GLN:HG3	1.66	0.95
1:A:30:ARG:HH11	1:A:30:ARG:CG	1.93	0.81
1:A:30:ARG:HH11	1:A:30:ARG:HA	1.49	0.77
1:B:177:TYR:CD1	1:B:177:TYR:N	2.54	0.75
1:A:30:ARG:HH11	1:A:30:ARG:HG3	1.51	0.75
1:A:30:ARG:HG3	1:A:30:ARG:NH1	2.02	0.75
1:B:177:TYR:N	1:B:177:TYR:HD1	1.85	0.74
1:B:177:TYR:HA	1:B:248:ILE:CD1	2.17	0.74
1:A:30:ARG:HH11	1:A:30:ARG:CA	2.02	0.72
1:B:307:ARG:HH11	1:B:307:ARG:HB2	1.52	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:ARG:HA	1:A:30:ARG:NH1	2.06	0.70
1:B:177:TYR:HD1	1:B:177:TYR:H	1.41	0.68
1:A:157:ARG:HD2	3:A:539:HOH:O	1.95	0.67
1:A:51:ALA:HA	1:A:54:GLN:HE21	1.58	0.66
1:B:71:GLU:CG	3:B:561:HOH:O	2.43	0.65
1:B:287:ARG:CZ	1:B:297:GLY:HA2	2.27	0.65
1:B:326:ARG:HD2	3:B:543:HOH:O	1.95	0.65
1:B:6:LEU:CD2	1:B:304:LEU:HD21	2.28	0.64
1:B:6:LEU:HD21	1:B:304:LEU:HD21	1.80	0.64
1:A:30:ARG:HH11	1:A:30:ARG:CB	2.11	0.63
1:B:51:ALA:HA	1:B:54:GLN:HE21	1.62	0.63
1:A:30:ARG:CG	1:A:30:ARG:NH1	2.59	0.62
1:B:218:THR:HB	1:B:219:PRO:HD3	1.85	0.57
1:B:97:GLN:O	1:B:101:ARG:HG3	2.05	0.56
1:B:291:ARG:HD2	1:B:291:ARG:H	1.71	0.56
1:B:303:ASP:OD2	1:B:307:ARG:NH1	2.40	0.54
1:A:254:GLU:H	1:A:254:GLU:CD	2.15	0.54
1:B:237:VAL:HG11	1:B:245:MET:HE3	1.89	0.53
1:B:132:TRP:CH2	1:B:212:VAL:HG11	2.43	0.53
1:A:270:ASP:HB2	3:A:617:HOH:O	2.08	0.53
1:A:338:ARG:HD2	3:A:516:HOH:O	2.07	0.53
1:B:287:ARG:NH1	1:B:297:GLY:HA2	2.24	0.53
1:A:68:LYS:HD3	1:A:377:LEU:HD21	1.91	0.52
1:B:274:ALA:HB3	1:B:324:LEU:HB3	1.91	0.52
1:B:71:GLU:HG2	3:B:561:HOH:O	2.05	0.51
1:B:72:GLN:OE1	1:B:377:LEU:HD13	2.10	0.51
1:A:326:ARG:HD2	1:A:332:GLU:OE1	2.10	0.51
1:B:141:ARG:HG2	1:B:142:SER:H	1.76	0.50
1:A:153:LEU:O	1:A:157:ARG:HG3	2.12	0.49
1:B:246:ARG:HD3	1:B:246:ARG:C	2.38	0.48
1:A:132:TRP:CH2	1:A:212:VAL:HG11	2.48	0.48
1:B:41:HIS:CE1	1:B:43:LYS:HB3	2.50	0.47
1:A:318:ASN:OD1	1:B:141:ARG:CD	2.63	0.47
1:A:332:GLU:OE2	1:A:338:ARG:NH2	2.42	0.47
1:A:11:PRO:HA	1:A:261:LEU:O	2.15	0.47
1:A:288:GLY:O	1:A:291:ARG:HG2	2.14	0.47
1:B:340:PRO:O	1:B:343:THR:OG1	2.27	0.46
1:B:338:ARG:HH11	1:B:338:ARG:HG3	1.80	0.46
1:B:314:LEU:HA	1:B:323:ILE:O	2.16	0.46
1:B:68:LYS:HG2	1:B:377:LEU:HD21	1.99	0.45
1:A:172:HIS:HB2	1:A:234:ARG:NH2	2.32	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:TYR:HA	1:A:248:ILE:HD11	1.98	0.45
1:A:266:GLY:HA3	3:B:545:HOH:O	2.16	0.45
1:A:67:LEU:HD23	1:A:67:LEU:HA	1.81	0.45
1:B:11:PRO:HA	1:B:261:LEU:O	2.16	0.44
1:A:149:THR:HG22	1:A:153:LEU:HG	1.99	0.44
1:B:3:ASP:O	1:B:371:VAL:HG22	2.18	0.44
1:A:28:GLN:HG2	1:A:57:ALA:O	2.18	0.44
1:A:46:LYS:NZ	3:A:582:HOH:O	2.50	0.43
1:A:246:ARG:C	1:A:246:ARG:HD3	2.43	0.43
1:B:49:PRO:O	1:B:52:ALA:HB3	2.19	0.43
1:A:138:ASP:OD1	1:A:140:HIS:HB3	2.19	0.43
1:A:134:GLU:OE2	1:A:142:SER:HB2	2.19	0.42
1:B:2:GLN:HG3	1:B:370:SER:HB2	2.02	0.42
1:B:281:MET:O	1:B:358:GLN:HG3	2.19	0.42
1:A:170:LEU:HA	1:A:214:SER:O	2.20	0.42
1:A:215:VAL:O	1:A:233:VAL:HA	2.20	0.42
1:A:323:ILE:N	1:A:323:ILE:CD1	2.82	0.41
1:B:296:PHE:O	1:B:297:GLY:C	2.62	0.41
1:A:140:HIS:CD2	1:A:174:GLY:HA3	2.56	0.41
1:B:17:LEU:HD23	1:B:17:LEU:HA	1.95	0.41
1:A:92:LEU:HA	1:A:92:LEU:HD23	1.91	0.41
1:B:170:LEU:HA	1:B:214:SER:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/390 (96%)	369 (98%)	5 (1%)	2 (0%)	24	31
1	B	377/390 (97%)	362 (96%)	14 (4%)	1 (0%)	36	46
All	All	753/780 (96%)	731 (97%)	19 (2%)	3 (0%)	30	38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	ARG
1	B	241	PHE
1	A	236	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/289 (96%)	273 (98%)	6 (2%)	45	65
1	B	280/289 (97%)	267 (95%)	13 (5%)	24	36
All	All	559/578 (97%)	540 (97%)	19 (3%)	32	49

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	30	ARG
1	A	60	SER
1	A	86	SER
1	A	291	ARG
1	A	323	ILE
1	B	2	GLN
1	B	3	ASP
1	B	30	ARG
1	B	45	SER
1	B	86	SER
1	B	93	PRO
1	B	176	SER
1	B	177	TYR
1	B	226	ARG
1	B	293	LYS
1	B	307	ARG
1	B	318	ASN
1	B	344	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	54	GLN
1	A	94	GLN
1	A	125	GLN
1	B	21	GLN
1	B	28	GLN
1	B	54	GLN
1	B	94	GLN
1	B	300	GLN
1	B	358	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLP	A	401	1	15,15,16	3.47	6 (40%)	21,22,23	2.20	9 (42%)
2	PLP	B	401	1	15,15,16	3.21	3 (20%)	21,22,23	1.39	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	401	1	-	3/6/6/8	0/1/1/1
2	PLP	B	401	1	-	3/6/6/8	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	PLP	C3-C2	9.31	1.50	1.41
2	B	401	PLP	C3-C2	8.48	1.49	1.41
2	B	401	PLP	C5-C4	7.43	1.48	1.40
2	A	401	PLP	C5-C4	6.27	1.47	1.40
2	A	401	PLP	C3-C4	4.65	1.49	1.40
2	B	401	PLP	C3-C4	4.19	1.48	1.40
2	A	401	PLP	C4A-C4	-2.93	1.45	1.51
2	A	401	PLP	P-O3P	-2.52	1.45	1.54
2	A	401	PLP	C6-N1	2.36	1.39	1.34

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	PLP	C5A-C5-C6	4.50	126.69	119.36
2	A	401	PLP	C3-C2-N1	-3.81	116.15	120.96
2	A	401	PLP	O3P-P-O4P	-3.41	97.78	106.67
2	A	401	PLP	C6-N1-C2	3.23	125.06	119.20
2	B	401	PLP	O3P-P-O4P	-3.19	98.34	106.67
2	A	401	PLP	O3P-P-O2P	3.00	119.06	107.80
2	A	401	PLP	C5A-C5-C4	-2.72	117.27	122.64
2	A	401	PLP	C4A-C4-C5	-2.62	118.24	120.94
2	B	401	PLP	O2P-P-O1P	2.57	120.87	110.83
2	A	401	PLP	C6-C5-C4	-2.53	116.03	118.10
2	B	401	PLP	C6-N1-C2	2.21	123.21	119.20
2	A	401	PLP	C2A-C2-N1	2.19	121.76	117.64

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	PLP	C5A-O4P-P-O2P

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	401	PLP	C5A-O4P-P-O1P
2	B	401	PLP	C5A-O4P-P-O2P
2	B	401	PLP	C5A-O4P-P-O3P
2	A	401	PLP	C5A-O4P-P-O3P
2	A	401	PLP	C5A-O4P-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	378/390 (96%)	-0.25	5 (1%) 75 76	19, 32, 60, 78	0
1	B	379/390 (97%)	0.51	27 (7%) 22 24	18, 45, 73, 89	0
All	All	757/780 (97%)	0.13	32 (4%) 40 42	18, 39, 68, 89	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	177	TYR	4.7
1	B	333	ALA	4.5
1	A	140	HIS	4.2
1	B	4	THR	4.2
1	B	140	HIS	3.3
1	A	177	TYR	3.3
1	B	12	ALA	3.1
1	B	252	ALA	3.1
1	B	2	GLN	3.1
1	B	367	ALA	3.0
1	B	310	PRO	2.7
1	A	141	ARG	2.7
1	B	57	ALA	2.6
1	B	330	ALA	2.5
1	B	345	LEU	2.5
1	B	139	GLY	2.4
1	A	334	ASP	2.4
1	B	335	ILE	2.4
1	B	291	ARG	2.3
1	B	253	ALA	2.3
1	B	327	ALA	2.3
1	B	25	ALA	2.2
1	B	326	ARG	2.2
1	B	374	TRP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	5	LEU	2.1
1	B	254	GLU	2.1
1	B	301	VAL	2.1
1	A	377	LEU	2.1
1	B	6	LEU	2.1
1	B	141	ARG	2.1
1	B	14	VAL	2.0
1	B	371	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PLP	B	401	15/16	0.95	0.08	33,38,45,46	0
2	PLP	A	401	15/16	0.98	0.05	26,28,30,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.