



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 01:48 AM UTC

PDB ID : 6WQN / pdb_00006wqn
EMDB ID : EMD-21872
Title : Structure of the 50S subunit of the ribosome from Methicillin Resistant Staphylococcus aureus in complex with the antibiotic, contezolid
Authors : Belousoff, M.J.
Deposited on : 2020-04-29
Resolution : 2.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	13054 (2.40 - 3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	116	
2	B	277	
3	C	118	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	105	
5	E	117	
6	F	91	
7	G	105	
8	H	107	
9	J	62	
10	K	72	
11	L	217	
12	M	58	
13	N	57	
14	O	49	
15	P	50	
16	Q	65	
17	R	37	
18	S	207	
19	V	145	
20	W	122	
21	X	146	
22	Y	144	
23	Z	122	
24	a	119	
25	1	2923	
26	2	115	
27	I	85	

2 Entry composition [i](#)

There are 28 unique types of molecules in this entry. The entry contains 80715 atoms, of which 15 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 50S ribosomal protein L19.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	A	112	Total	C	N	O	0	0
			907	572	182	153		

- Molecule 2 is a protein called 50S ribosomal protein L2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	274	Total	C	N	O	S	0	0
			2094	1303	415	371	5		

- Molecule 3 is a protein called 50S ribosomal protein L20.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	116	Total	C	N	O	S	0	0
			943	593	189	157	4		

- Molecule 4 is a protein called 50S ribosomal protein L21.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	100	Total	C	N	O	S	0	0
			785	499	139	146	1		

- Molecule 5 is a protein called 50S ribosomal protein L22.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	110	Total	C	N	O	S	0	0
			845	527	162	154	2		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	110	ALA	GLY	variant	UNP A0A077UKF9

- Molecule 6 is a protein called 50S ribosomal protein L23.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	81	Total	C	N	O	S	0	0
			654	410	116	125	3		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	87	ASP	ILE	variant	UNP W8TUB4

- Molecule 7 is a protein called 50S ribosomal protein L24.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	91	Total	C	N	O	S	0	0
			702	444	129	128	1		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	54	SER	GLY	variant	UNP W8TRD5

- Molecule 8 is a protein called 50S ribosomal protein L25.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	93	Total	C	N	O	S	0	0
			727	465	129	132	1		

- Molecule 9 is a protein called 50S ribosomal protein L28.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	J	59	Total	C	N	O	S	0	0
			463	287	99	76	1		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
J	62	ALA	-	insertion	UNP A0A077URJ8

- Molecule 10 is a protein called 50S ribosomal protein L29.

Mol	Chain	Residues	Atoms				AltConf	Trace
10	K	53	Total	C	N	O	0	0
			436	269	82	85		

- Molecule 11 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	L	215	Total	C	N	O	S	0	0
			1621	1015	299	303	4		

- Molecule 12 is a protein called 50S ribosomal protein L30.

Mol	Chain	Residues	Atoms				AltConf	Trace
12	M	56	Total	C	N	O	0	0
			432	269	82	81		

- Molecule 13 is a protein called 50S ribosomal protein L32.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	N	50	Total	C	N	O	S	0	0
			397	241	83	68	5		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
N	54	ALA	VAL	variant	UNP A0A077UWR7

- Molecule 14 is a protein called 50S ribosomal protein L33.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	O	47	Total	C	N	O	S	0	0
			390	233	79	73	5		

- Molecule 15 is a protein called 50S ribosomal protein L34.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	P	44	Total	C	N	O	S	0	0
			372	228	90	53	1		

- Molecule 16 is a protein called 50S ribosomal protein L35.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	Q	64	Total	C	N	O	S	0	0
			521	324	113	82	2		

- Molecule 17 is a protein called 50S ribosomal protein L36.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	R	37	Total	C	N	O	S	0	0
			296	186	60	45	5		

- Molecule 18 is a protein called 50S ribosomal protein L4.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	S	192	Total	C	N	O	S	0	0
			1472	924	271	275	2		

- Molecule 19 is a protein called 50S ribosomal protein L13.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	V	143	Total	C	N	O	S	0	0
			1138	710	209	217	2		

- Molecule 20 is a protein called 50S ribosomal protein L14.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	W	121	Total	C	N	O	S	0	0
			911	566	173	168	4		

- Molecule 21 is a protein called 50S ribosomal protein L15.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	X	144	Total	C	N	O		0	0
			1082	669	213	200			

- Molecule 22 is a protein called 50S ribosomal protein L16.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	Y	136	Total	C	N	O	S	0	0
			1089	698	206	181	4		

- Molecule 23 is a protein called 50S ribosomal protein L17.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	Z	120	Total	C	N	O	S	0	0
			951	584	182	184	1		

- Molecule 24 is a protein called 50S ribosomal protein L18.

Mol	Chain	Residues	Atoms				AltConf	Trace
24	a	110	Total	C	N	O	0	0
			857	536	165	156		

- Molecule 25 is a RNA chain called 23S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	1	2687	Total	C	N	O	P	0	0
			57631	25735	10578	18635	2683		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	1866	A	G	conflict	GB 1760383645

- Molecule 26 is a RNA chain called 5S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	2	111	Total	C	N	O	P	0	0
			2358	1056	422	770	110		


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
2	80	C	G	variant	GB 1750990749
2	109	C	G	variant	GB 1750990749
2	111	A	C	variant	GB 1750990749
2	112	G	A	variant	GB 1750990749

- Molecule 27 is a protein called 50S ribosomal protein L27.

Mol	Chain	Residues	Atoms				AltConf	Trace
27	I	78	Total	C	N	O	0	0
			597	367	116	114		

- Molecule 28 is Contezolid (CCD ID: ZC0) (formula: C₁₈H₁₅F₃N₄O₄) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						AltConf
			Total	C	F	H	N	O	
28	1	1	44	18	3	15	4	4	0




- Molecule 6: 50S ribosomal protein L23




- Molecule 7: 50S ribosomal protein L24




- Molecule 8: 50S ribosomal protein L25



- Molecule 9: 50S ribosomal protein L28




- Molecule 10: 50S ribosomal protein L29



- Molecule 11: 50S ribosomal protein L3





- Molecule 12: 50S ribosomal protein L30

Chain M: 72% 24% .



- Molecule 13: 50S ribosomal protein L32

Chain N: 77% 11% 12%



- Molecule 14: 50S ribosomal protein L33

Chain O: 55% 41% .




- Molecule 15: 50S ribosomal protein L34

Chain P: 68% 18% 12%



- Molecule 16: 50S ribosomal protein L35

Chain Q: 83% 15% .




- Molecule 17: 50S ribosomal protein L36

Chain R: 76% 24%



- Molecule 18: 50S ribosomal protein L4

Chain S:  65% 27% 7%



- Molecule 19: 50S ribosomal protein L13

Chain V:  79% 19% ..




- Molecule 20: 50S ribosomal protein L14

Chain W:  83% 16% ..




- Molecule 21: 50S ribosomal protein L15

Chain X:  73% 25% ..




- Molecule 22: 50S ribosomal protein L16

Chain Y:  69% 24% 6%




- Molecule 23: 50S ribosomal protein L17


Chain Z:  70% 27% ..




- Molecule 24: 50S ribosomal protein L18




- Molecule 25: 23S rRNA







● Molecule 26: 5S rRNA



● Molecule 27: 50S ribosomal protein L27



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	127000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	47	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.435	Depositor
Minimum map value	-0.224	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.012	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	372.32, 372.32, 372.32	wwPDB
Map dimensions	416, 416, 416	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.89500004, 0.89500004, 0.89500004	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZC0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/919	0.85	0/1228
2	B	0.24	0/2129	0.37	0/2858
3	C	0.08	0/955	0.26	0/1265
4	D	0.30	0/795	0.44	0/1062
5	E	0.30	0/853	0.42	0/1149
6	F	0.32	0/659	0.45	0/879
7	G	0.75	0/708	0.86	1/943 (0.1%)
8	H	0.62	0/735	0.75	0/986
9	J	0.36	0/469	0.51	0/625
10	K	0.09	0/437	0.31	0/583
11	L	0.43	0/1645	0.66	3/2208 (0.1%)
12	M	0.07	0/434	0.24	0/585
13	N	0.66	2/404 (0.5%)	0.68	1/537 (0.2%)
14	O	0.09	0/393	0.27	0/523
15	P	0.49	0/376	0.72	0/491
16	Q	0.35	0/526	0.53	0/690
17	R	0.75	0/299	0.96	0/393
18	S	0.22	0/1494	0.37	0/2018
19	V	0.30	0/1160	0.39	0/1563
20	W	0.51	0/918	0.69	0/1232
21	X	0.38	2/1096 (0.2%)	0.48	0/1461
22	Y	0.22	0/1113	0.36	0/1493
23	Z	0.08	0/955	0.28	0/1277
24	a	0.55	0/865	0.68	0/1154
25	1	0.09	0/64545	0.30	36/100652 (0.0%)
26	2	0.04	0/2636	0.09	0/4105
27	I	0.08	0/603	0.27	0/801
All	All	0.21	4/88121 (0.0%)	0.36	41/132761 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	N	5	LYS	C-N	8.48	1.45	1.33
21	X	71	ARG	C-N	6.63	1.42	1.33
13	N	6	ARG	C-N	6.58	1.42	1.33
21	X	70	ASN	C-N	5.86	1.42	1.33

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
25	1	254	A	C4'-C3'-O3'	10.59	128.88	113.00
11	L	159	ASP	N-CA-C	-10.55	99.78	111.07
25	1	2078	A	C4'-C3'-O3'	-8.37	96.84	109.40
25	1	1405	G	C4'-C3'-O3'	7.67	124.51	113.00
25	1	2850	G	C4'-C3'-O3'	7.66	124.48	113.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	907	0	981	24	0
2	B	2094	0	2205	47	0
3	C	943	0	1014	25	0
4	D	785	0	825	23	0
5	E	845	0	900	23	0
6	F	654	0	686	25	0
7	G	702	0	759	15	0
8	H	727	0	777	31	0
9	J	463	0	501	18	0
10	K	436	0	459	27	0
11	L	1621	0	1655	43	0
12	M	432	0	472	7	0
13	N	397	0	407	4	0
14	O	390	0	396	14	0
15	P	372	0	420	15	0
16	Q	521	0	586	7	0
17	R	296	0	340	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
18	S	1472	0	1520	46	0
19	V	1138	0	1130	24	0
20	W	911	0	970	25	0
21	X	1082	0	1119	33	0
22	Y	1089	0	1155	28	0
23	Z	951	0	999	28	0
24	a	857	0	903	32	0
25	1	57631	0	28986	612	0
26	2	2358	0	1198	54	0
27	I	597	0	607	8	0
28	1	29	15	0	0	0
All	All	80700	15	51970	1162	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 1162 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
25:1:1914:C:H2'	25:1:1915:G:H5''	1.48	0.94
25:1:1083:G:H2'	25:1:1084:U:H5''	1.53	0.91
25:1:1625:U:H2'	25:1:1626:A:H5''	1.53	0.90
20:W:112:MET:HE2	20:W:112:MET:HA	1.53	0.90
15:P:41:LYS:HD2	25:1:505:U:H5''	1.52	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	110/116 (95%)	103 (94%)	7 (6%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	272/277 (98%)	264 (97%)	8 (3%)	0	100	100
3	C	114/118 (97%)	111 (97%)	3 (3%)	0	100	100
4	D	98/105 (93%)	92 (94%)	6 (6%)	0	100	100
5	E	108/117 (92%)	106 (98%)	2 (2%)	0	100	100
6	F	77/91 (85%)	75 (97%)	2 (3%)	0	100	100
7	G	87/105 (83%)	85 (98%)	2 (2%)	0	100	100
8	H	91/107 (85%)	91 (100%)	0	0	100	100
9	J	57/62 (92%)	56 (98%)	1 (2%)	0	100	100
10	K	51/72 (71%)	48 (94%)	3 (6%)	0	100	100
11	L	213/217 (98%)	204 (96%)	9 (4%)	0	100	100
12	M	54/58 (93%)	54 (100%)	0	0	100	100
13	N	48/57 (84%)	48 (100%)	0	0	100	100
14	O	45/49 (92%)	44 (98%)	1 (2%)	0	100	100
15	P	42/50 (84%)	42 (100%)	0	0	100	100
16	Q	62/65 (95%)	62 (100%)	0	0	100	100
17	R	35/37 (95%)	35 (100%)	0	0	100	100
18	S	190/207 (92%)	183 (96%)	7 (4%)	0	100	100
19	V	141/145 (97%)	140 (99%)	1 (1%)	0	100	100
20	W	119/122 (98%)	116 (98%)	3 (2%)	0	100	100
21	X	142/146 (97%)	135 (95%)	7 (5%)	0	100	100
22	Y	134/144 (93%)	129 (96%)	5 (4%)	0	100	100
23	Z	118/122 (97%)	116 (98%)	2 (2%)	0	100	100
24	a	106/119 (89%)	103 (97%)	3 (3%)	0	100	100
27	I	76/85 (89%)	73 (96%)	3 (4%)	0	100	100
All	All	2590/2793 (93%)	2515 (97%)	75 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/102 (96%)	96 (98%)	2 (2%)	48	78
2	B	221/224 (99%)	216 (98%)	5 (2%)	44	76
3	C	96/98 (98%)	90 (94%)	6 (6%)	16	45
4	D	85/89 (96%)	84 (99%)	1 (1%)	63	86
5	E	89/94 (95%)	88 (99%)	1 (1%)	65	88
6	F	74/82 (90%)	73 (99%)	1 (1%)	59	85
7	G	77/91 (85%)	74 (96%)	3 (4%)	28	63
8	H	81/95 (85%)	79 (98%)	2 (2%)	42	74
9	J	49/51 (96%)	48 (98%)	1 (2%)	48	78
10	K	48/65 (74%)	45 (94%)	3 (6%)	16	45
11	L	170/175 (97%)	163 (96%)	7 (4%)	27	61
12	M	50/52 (96%)	49 (98%)	1 (2%)	48	78
13	N	45/49 (92%)	45 (100%)	0	100	100
14	O	45/47 (96%)	45 (100%)	0	100	100
15	P	39/45 (87%)	38 (97%)	1 (3%)	40	73
16	Q	55/56 (98%)	55 (100%)	0	100	100
17	R	35/35 (100%)	35 (100%)	0	100	100
18	S	158/170 (93%)	155 (98%)	3 (2%)	50	79
19	V	122/123 (99%)	120 (98%)	2 (2%)	55	83
20	W	99/100 (99%)	98 (99%)	1 (1%)	68	89
21	X	110/112 (98%)	109 (99%)	1 (1%)	70	90
22	Y	113/119 (95%)	109 (96%)	4 (4%)	32	66
23	Z	101/102 (99%)	100 (99%)	1 (1%)	68	89
24	a	87/95 (92%)	84 (97%)	3 (3%)	32	66
27	I	61/66 (92%)	59 (97%)	2 (3%)	33	67
All	All	2208/2337 (94%)	2157 (98%)	51 (2%)	44	76

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
11	L	117	ASP
18	S	58	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
27	I	16	SER
11	L	122	SER
12	M	29	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
19	V	97	ASN
21	X	78	ASN
20	W	72	ASN
21	X	81	GLN
11	L	148	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
25	1	2675/2923 (91%)	444 (16%)	48 (1%)
26	2	110/115 (95%)	30 (27%)	4 (3%)
All	All	2785/3038 (91%)	474 (17%)	52 (1%)

5 of 474 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
25	1	5	A
25	1	13	A
25	1	28	A
25	1	35	G
25	1	42	G

5 of 52 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
25	1	1568	U
25	1	2337	A
26	2	57	G
25	1	1864	C
25	1	2320	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
28	ZC0	1	3001	-	30,32,32	3.02	10 (33%)	37,46,46	2.74	16 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
28	ZC0	1	3001	-	-	6/13/35/35	0/4/4/4

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
28	1	3001	ZC0	C20-N16	8.99	1.46	1.36
28	1	3001	ZC0	C04-C05	7.34	1.52	1.35
28	1	3001	ZC0	O19-C20	6.64	1.44	1.35
28	1	3001	ZC0	C04-N03	4.91	1.50	1.36
28	1	3001	ZC0	O26-N25	-3.61	1.36	1.41

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
28	1	3001	ZC0	C17-C18-C22	-6.27	106.26	113.38
28	1	3001	ZC0	C17-N16-C20	-5.95	106.77	111.17
28	1	3001	ZC0	C18-C17-N16	5.38	106.93	101.85
28	1	3001	ZC0	C04-C05-C06	-4.83	117.53	120.89
28	1	3001	ZC0	C09-C08-C13	4.36	119.93	115.25

There are no chirality outliers.


5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
28	1	3001	ZC0	C09-C08-N03-C02
28	1	3001	ZC0	C13-C08-N03-C02
28	1	3001	ZC0	C10-C11-N16-C20
28	1	3001	ZC0	C12-C11-N16-C17
28	1	3001	ZC0	C12-C11-N16-C20

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-21872. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 208




Y Index: 208




Z Index: 208


6.2.2 Raw map



X Index: 208



Y Index: 208




Z Index: 208


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 216




Y Index: 186




Z Index: 174


6.3.2 Raw map



X Index: 223



Y Index: 186




Z Index: 174


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1087 nm³; this corresponds to an approximate mass of 982 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of $0.345 \text{ \AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.345 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.90	-	-
Author-provided FSC curve	2.86	3.17	2.91
Unmasked-calculated*	3.21	4.19	3.28

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.21 differs from the reported value 2.9 by more than 10 %


9 Map-model fit ⓘ

This section contains information regarding the fit between EMDB map EMD-21872 and PDB model 6WQN. Per-residue inclusion information can be found in section 3 on page 10.

9.1 Map-model overlay ⓘ

This section was not generated.

9.2 Q-score mapped to coordinate model ⓘ




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model ⓘ

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 97% of all backbone atoms, 97% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9660	 0.6010
1	 0.9680	 0.6040
2	 0.9400	 0.4950
A	 0.9660	 0.5960
B	 0.9690	 0.6230
C	 0.9920	 0.6480
D	 0.9530	 0.6090
E	 0.9830	 0.6390
F	 0.9560	 0.5870
G	 0.9290	 0.5440
H	 0.9030	 0.5040
I	 0.9900	 0.6410
J	 0.9460	 0.5890
K	 0.9340	 0.5600
L	 0.9750	 0.6200
M	 0.9650	 0.6180
N	 0.9690	 0.6140
O	 0.9440	 0.5800
P	 0.9890	 0.6630
Q	 0.9960	 0.6600
R	 0.9790	 0.6000
S	 0.9700	 0.6090
V	 0.9860	 0.6340
W	 0.9710	 0.6150
X	 0.9620	 0.6010
Y	 0.9740	 0.6160
Z	 0.9640	 0.6190
a	 0.9340	 0.5200

